

Ryuji Furihata

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2782234/publications.pdf>

Version: 2024-02-01

32
papers

699
citations

516710

16
h-index

580821

25
g-index

35
all docs

35
docs citations

35
times ranked

786
citing authors

#	ARTICLE	IF	CITATIONS
1	Hypnotic medication use among inpatients with schizophrenia and major depressive disorder: results of a nationwide study. <i>Sleep Medicine</i> , 2022, 89, 23-30.	1.6	16
2	Association between the examination rate of treatment-resistant schizophrenia and the clozapine prescription rate in a nationwide dissemination and implementation study. <i>Neuropsychopharmacology Reports</i> , 2022, 42, 3-9.	2.3	14
3	The characteristics of patients receiving psychotropic pro re nata medication at discharge for the treatment of schizophrenia and major depressive disorder: A nationwide survey from the EGUIDE project. <i>Asian Journal of Psychiatry</i> , 2022, 69, 103007.	2.0	12
4	Subjective assessment of participants in education programs on clinical practice guidelines in the field of psychiatry. <i>Neuropsychopharmacology Reports</i> , 2022, 42, 221-225.	2.3	12
5	A dissemination and education programme to improve the clinical behaviours of psychiatrists in accordance with treatment guidelines for schizophrenia and major depressive disorders: the Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE) project. <i>BIPsych Open</i> , 2022, 8, e83.	0.7	11
6	Prescription of Anticholinergic Drugs in Patients With Schizophrenia: Analysis of Antipsychotic Prescription Patterns and Hospital Characteristics. <i>Frontiers in Psychiatry</i> , 2022, 13, .	2.6	9
7	Clozapine Treatment Is Associated With Higher Prescription Rate of Antipsychotic Monotherapy and Lower Prescription Rate of Other Concomitant Psychotropics: A Real-World Nationwide Study. <i>International Journal of Neuropsychopharmacology</i> , 2022, 25, 818-826.	2.1	11
8	Improvements in the degree of understanding the treatment guidelines for schizophrenia and major depressive disorder in a nationwide dissemination and implementation study. <i>Neuropsychopharmacology Reports</i> , 2021, 41, 199-206.	2.3	17
9	Characteristics of discharge prescriptions for patients with schizophrenia or major depressive disorder: Real-world evidence from the Effectiveness of Guidelines for Dissemination and Education (EGUIDE) psychiatric treatment project. <i>Asian Journal of Psychiatry</i> , 2021, 63, 102744.	2.0	28
10	High-dose zolpidem abuse in a patient with insomnia comorbid with major depressive disorder. <i>Sleep and Biological Rhythms</i> , 2020, 18, 155-157.	1.0	3
11	Association between reduced serum BDNF levels and insomnia with short sleep duration among female hospital nurses. <i>Sleep Medicine</i> , 2020, 68, 167-172.	1.6	26
12	A composite measure of sleep health is associated with symptoms of depression among Japanese female hospital nurses. <i>Comprehensive Psychiatry</i> , 2020, 97, 152151.	3.1	27
13	Unmet needs of patients with major depressive disorder – Findings from the “Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE)” project: A nationwide dissemination, education, and evaluation study. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 667-669.	1.8	20
14	Association between Typus melancholicus and depressive symptoms: A Japanese general population survey. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 672-673.	1.8	2
15	Prescription patterns in patients with schizophrenia in Japan: First-quality indicator data from the survey of “Effectiveness of Guidelines for Dissemination and Education in psychiatric treatment (EGUIDE)” project. <i>Neuropsychopharmacology Reports</i> , 2020, 40, 281-286.	2.3	32
16	Identification of Chochukei (Zhang Zhongjing) as the revolutionist of decoctions. <i>Traditional & Kampo Medicine</i> , 2020, 7, 17-23.	0.6	0
17	Sleep disturbance is associated with not only shorter sleep duration, but also longer time in bed: a Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 2019, 17, 407-415.	1.0	6
18	Effect of <i>Enterococcus faecalis</i> 2001 on colitis and depressive-like behavior in dextran sulfate sodium-treated mice: involvement of the brain-gut axis. <i>Journal of Neuroinflammation</i> , 2019, 16, 201.	7.2	59

#	ARTICLE	IF	CITATIONS
19	Improvement of psychiatrists' clinical knowledge of the treatment guidelines for schizophrenia and major depressive disorders using the "Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE)" project: A nationwide dissemination, education, and evaluation study. <i>Psychiatry and Clinical Neurosciences</i> , 2019, 73, 642-648.	1.8	35
20	REM sleep behavior disorder predisposed by neuroleptic malignant syndrome: a case report. <i>Sleep and Biological Rhythms</i> , 2019, 17, 155-157.	1.0	2
21	Seasonal changes in sleep duration and sleep problems: A prospective study in Japanese community residents. <i>PLoS ONE</i> , 2019, 14, e0215345.	2.5	37
22	Unhealthy lifestyle factors and depressive symptoms: A Japanese general adult population survey. <i>Journal of Affective Disorders</i> , 2018, 234, 156-161.	4.1	26
23	Association of serum BDNF levels and the BDNF Val66Met polymorphism with the sleep pattern in healthy young adults. <i>PLoS ONE</i> , 2018, 13, e0199765.	2.5	9
24	Stressful events and coping strategies associated with symptoms of depression: A Japanese general population survey. <i>Journal of Affective Disorders</i> , 2018, 238, 482-488.	4.1	33
25	An Aggregate Measure of Sleep Health Is Associated With Prevalent and Incident Clinically Significant Depression Symptoms Among Community-Dwelling Older Women. <i>Sleep</i> , 2017, 40, .	1.1	78
26	An Epidemiological Study on Psychosocial Factors Associated with Depression among Japanese General Adult Population. <i>Journal of the Nihon University Medical Association</i> , 2016, 75, 81-87.	0.0	0
27	Napping and associated factors: a Japanese nationwide general population survey. <i>Sleep Medicine</i> , 2016, 20, 72-79.	1.6	38
28	Association of short sleep duration and short time in bed with depression: A Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 2015, 13, 136-145.	1.0	38
29	The Relationship between Exploratory Eye Movement Dysfunction and Psychiatric Symptoms in Schizophrenia. <i>Journal of the Nihon University Medical Association</i> , 2015, 74, 283-291.	0.0	0
30	The association between sleep problems and perceived health status: A Japanese nationwide general population survey. <i>Sleep Medicine</i> , 2012, 13, 831-837.	1.6	72
31	Self-help behaviors for sleep and depression: A Japanese nationwide general population survey. <i>Journal of Affective Disorders</i> , 2011, 130, 75-82.	4.1	26
32	Association of insomnia symptoms and non-restorative sleep with Typus melancholicus: a Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 0, , 1.	1.0	0