

Wataru Yamamoto

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2782789/wataru-yamamoto-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

26

papers

312

citations

10

h-index

17

g-index

28

ext. papers

441

ext. citations

3.8

avg, IF

2.85

L-index

#	Paper	IF	Citations
26	Prevalence and factors associated with sarcopenia in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2019 , 29, 589-595	3.3	52
25	Drug retention and discontinuation reasons between seven biologics in patients with rheumatoid arthritis -The ANSWER cohort study. <i>PLoS ONE</i> , 2018 , 13, e0194130	3.7	49
24	Periodontitis and Porphyromonas gingivalis in preclinical stage of arthritis patients. <i>PLoS ONE</i> , 2015 , 10, e0122121	3.7	36
23	Increase of hemoglobin levels by anti-IL-6 receptor antibody (tocilizumab) in rheumatoid arthritis. <i>PLoS ONE</i> , 2014 , 9, e98202	3.7	26
22	Drug tolerability and reasons for discontinuation of seven biologics in elderly patients with rheumatoid arthritis -The ANSWER cohort study. <i>PLoS ONE</i> , 2019 , 14, e0216624	3.7	23
21	Drug tolerability and reasons for discontinuation of seven biologics in 4466 treatment courses of rheumatoid arthritis-the ANSWER cohort study. <i>Arthritis Research and Therapy</i> , 2019 , 21, 91	5.7	23
20	Elderly onset of early rheumatoid arthritis is a risk factor for bone erosions, refractory to treatment: KURAMA cohort. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 1084-1093	2.3	21
19	Increased circulating adiponectin is an independent disease activity marker in patients with rheumatoid arthritis: A cross-sectional study using the KURAMA database. <i>PLoS ONE</i> , 2020 , 15, e0229998	3.7	14
18	Non-obese visceral adiposity is associated with the risk of atherosclerosis in Japanese patients with rheumatoid arthritis: a cross-sectional study. <i>Rheumatology International</i> , 2018 , 38, 1679-1689	3.6	11
17	Drug retention of 7 biologics and tofacitinib in biologics-naïve and biologics-switched patients with rheumatoid arthritis: the ANSWER cohort study. <i>Arthritis Research and Therapy</i> , 2020 , 22, 142	5.7	10
16	The association between serious infection and disease outcome in patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2016 , 35, 213-8	3.9	7
15	Drug retention of secondary biologics or JAK inhibitors after tocilizumab or abatacept failure as first biologics in patients with rheumatoid arthritis -the ANSWER cohort study. <i>Clinical Rheumatology</i> , 2020 , 39, 2563-2572	3.9	6
14	Urinary sodium-to-potassium ratio associates with hypertension and current disease activity in patients with rheumatoid arthritis: a cross-sectional study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 96	5.7	6
13	Habitual fish intake negatively correlates with prevalence of frailty among patients with rheumatoid arthritis. <i>Scientific Reports</i> , 2021 , 11, 5104	4.9	4
12	Drug retention of sarilumab, baricitinib, and tofacitinib in patients with rheumatoid arthritis: the ANSWER cohort study. <i>Clinical Rheumatology</i> , 2021 , 40, 2673-2680	3.9	4
11	Comparison of the efficacy and safety of biologic agents between elderly-onset and young-onset RA patients: the ANSWER cohort study. <i>Rheumatology International</i> , 2020 , 40, 1987-1995	3.6	3
10	Comparison of the drug retention and reasons for discontinuation of tumor necrosis factor inhibitors and interleukin-6 inhibitors in Japanese patients with elderly-onset rheumatoid arthritis-the ANSWER cohort study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 116	5.7	3

9	Comparison of efficacy between anti-IL-6 receptor antibody and other biological disease-modifying antirheumatic drugs in the patients with rheumatoid arthritis who have knee joint involvement: the ANSWER cohort, retrospective study. <i>Rheumatology International</i> , 2021 , 41, 1233-1241	3.6	2
8	The family history of rheumatoid arthritis in anti-cyclic citrullinated peptide antibody-positive patient is not a predictor of poor clinical presentation and treatment response with modern classification criteria and treatment strategy: the ANSWER cohort study. <i>Rheumatology International</i> , 2020 , 40, 217-225	3.6	2
7	Evaluation of risk factors for atherosclerosis using carotid ultrasonography in Japanese patients with rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 1312-1318	2.3	1
6	Factors affecting drug retention of Janus kinase inhibitors in patients with rheumatoid arthritis: the ANSWER cohort study.. <i>Scientific Reports</i> , 2022 , 12, 134	4.9	1
5	Pain catastrophizing hinders Disease Activity Score 28 - erythrocyte sedimentation rate remission of rheumatoid arthritis in patients with normal C-reactive protein levels. <i>International Journal of Rheumatic Diseases</i> , 2021 , 24, 1520-1529	2.3	1
4	Neutrophil count reduction 1 month after initiating tocilizumab can predict clinical remission within 1 year in rheumatoid arthritis patients. <i>Rheumatology International</i> , 2021 , 1	3.6	1
3	Differential efficacy of TNF inhibitors with or without the immunoglobulin fragment crystallizable (Fc) portion in rheumatoid arthritis: the ANSWER cohort study.. <i>Rheumatology International</i> , 2022 , 1	3.6	1
2	Influence of dietary habits on depression among patients with rheumatoid arthritis: A cross-sectional study using KURAMA cohort database. <i>PLoS ONE</i> , 2021 , 16, e0255526	3.7	0
1	Perceptions and behaviours related to COVID-19 in patients with rheumatoid arthritis: a cross-sectional study. <i>Clinical Rheumatology</i> , 2021 , 1	3.9	0