

Stphane Zaffran

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2783267/stephane-zaffran-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

90
papers

5,731
citations

30
h-index

75
g-index

107
ext. papers

6,537
ext. citations

7.3
avg, IF

5.45
L-index

#	Paper	IF	Citations
90	Identification of two variants in and genes in a patient with catecholaminergic polymorphic ventricular tachycardia suggesting new candidate disease genes and digenic inheritance.. <i>Clinical Case Reports (discontinued)</i> , 2022 , 10, e05339	0.7	1
89	Single Cell Approaches to Understand the Earliest Steps in Heart Development.. <i>Current Cardiology Reports</i> , 2022 , 1	4.2	0
88	Msx1 haploinsufficiency modifies the Pax9-deficient cardiovascular phenotype. <i>BMC Developmental Biology</i> , 2021 , 21, 14	3.1	3
87	Outflow Tract Formation-Embryonic Origins of Conotruncal Congenital Heart Disease. <i>Journal of Cardiovascular Development and Disease</i> , 2021 , 8,	4.2	2
86	Multiallelic rare variants support an oligogenic origin of sudden cardiac death in the young. <i>Herz</i> , 2021 , 46, 94-102	2.6	4
85	Side-dependent effect in the response of valve endothelial cells to bidirectional shear stress. <i>International Journal of Cardiology</i> , 2021 , 323, 220-228	3.2	2
84	Clinical insights into a tertiary care center cohort of patients with bicuspid aortic valve. <i>International Journal of Cardiovascular Imaging</i> , 2021 , 1	2.5	0
83	A roadmap for the Human Developmental Cell Atlas. <i>Nature</i> , 2021 , 597, 196-205	50.4	18
82	Piezo1 is required for outflow tract and aortic valve development. <i>Journal of Molecular and Cellular Cardiology</i> , 2020 , 143, 51-62	5.8	22
81	-dependent coordination of mouse cardiac progenitor cell patterning and differentiation. <i>ELife</i> , 2020 , 9,	8.9	16
80	Author response: Hox-dependent coordination of mouse cardiac progenitor cell patterning and differentiation 2020 ,		2
79	Identification of a peripheral blood gene signature predicting aortic valve calcification. <i>Physiological Genomics</i> , 2020 , 52, 563-574	3.6	2
78	A severe clinical phenotype of Noonan syndrome with neonatal hypertrophic cardiomyopathy in the second case worldwide with RAF1 S259Y neomutation. <i>Genetical Research</i> , 2019 , 101, e6	1.1	3
77	Human pre-valvular endocardial cells derived from pluripotent stem cells recapitulate cardiac pathophysiological valvulogenesis. <i>Nature Communications</i> , 2019 , 10, 1929	17.4	30
76	Transcriptome analysis of mouse and human sinoatrial node cells reveals a conserved genetic program. <i>Development (Cambridge)</i> , 2019 , 146,	6.6	30
75	Krox20 Regulates Endothelial Nitric Oxide Signaling in Aortic Valve Development and Disease. <i>Journal of Cardiovascular Development and Disease</i> , 2019 , 6,	4.2	5
74	Asb2/Filamin A Axis Is Essential for Actin Cytoskeleton Remodeling During Heart Development. <i>Circulation Research</i> , 2018 , 122, e34-e48	15.7	18

73	Giant congenital melanocytic nevus with vascular malformation and epidermal cysts associated with a somatic activating mutation in BRAF. <i>Pigment Cell and Melanoma Research</i> , 2018 , 31, 437-441	4.5	16
72	Myocardial Bmp2 gain causes ectopic EMT and promotes cardiomyocyte proliferation and immaturity. <i>Cell Death and Disease</i> , 2018 , 9, 399	9.8	15
71	Novel ALPK3 mutation in a Tunisian patient with pediatric cardiomyopathy and facio-thoraco-skeletal features. <i>Journal of Human Genetics</i> , 2018 , 63, 1077-1082	4.3	9
70	T-box genes and retinoic acid signaling regulate the segregation of arterial and venous pole progenitor cells in the murine second heart field. <i>Human Molecular Genetics</i> , 2018 , 27, 3747-3760	5.6	34
69	Krox20 defines a subpopulation of cardiac neural crest cells contributing to arterial valves and bicuspid aortic valve. <i>Development (Cambridge)</i> , 2018 , 145,	6.6	22
68	Analysis of HOXB1 gene in a cohort of patients with sporadic ventricular septal defect. <i>Molecular Biology Reports</i> , 2018 , 45, 1507-1513	2.8	
67	Ectopic expression of Hoxb1 induces cardiac and craniofacial malformations. <i>Genesis</i> , 2018 , 56, e23221	1.9	12
66	Hox and Tale transcription factors in heart development and disease. <i>International Journal of Developmental Biology</i> , 2018 , 62, 837-846	1.9	16
65	Bmp2 and Notch cooperate to pattern the embryonic endocardium. <i>Development (Cambridge)</i> , 2018 , 145,	6.6	20
64	Origines génétique et développementale de la bicuspidie aortique. <i>Archives Des Maladies Du Coeur Et Des Vaisseaux - Pratique</i> , 2017 , 2017, 22-26	0	
63	The alternatively spliced LRRFIP1 Isoform-1 is a key regulator of the Wnt/βcatenin transcription pathway. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017 , 1864, 1142-1152	4.9	8
62	Mechanisms of retinoic acid signaling during cardiogenesis. <i>Mechanisms of Development</i> , 2017 , 143, 9-19	1.7	52
61	Hoxa1 and Hoxb1 are required for pharyngeal arch artery development. <i>Mechanisms of Development</i> , 2017 , 143, 1-8	1.7	15
60	Reduced aggrecan expression affects cardiac outflow tract development in zebrafish and is associated with bicuspid aortic valve disease in humans. <i>International Journal of Cardiology</i> , 2017 , 249, 340-343	3.2	11
59	FOXC1 haploinsufficiency due to 6p25 deletion in a patient with rapidly progressing aortic valve disease. <i>American Journal of Medical Genetics, Part A</i> , 2017 , 173, 2489-2493	2.5	5
58	Actionable Genes, Core Databases, and Locus-Specific Databases. <i>Human Mutation</i> , 2016 , 37, 1299-1307	4.7	5
57	WES/WGS Reporting of Mutations from Cardiovascular "Actionable" Genes in Clinical Practice: A Key Role for UMD Knowledgebases in the Era of Big Databases. <i>Human Mutation</i> , 2016 , 37, 1308-1317	4.7	5
56	Krox20 heterozygous mice: A model of aortic regurgitation associated with decreased expression of fibrillar collagen genes. <i>Archives of Cardiovascular Diseases</i> , 2016 , 109, 188-98	2.7	4

55	Disruption of CXCR4 signaling in pharyngeal neural crest cells causes DiGeorge syndrome-like malformations. <i>Development (Cambridge)</i> , 2016 , 143, 582-8	6.6	27
54	An uncommon cause of tricuspid regurgitation: three-dimensional echocardiographic incremental value, surgical and genetic insights. <i>European Journal of Cardio-thoracic Surgery</i> , 2016 , 50, 180-2	3	
53	Hox Genes in Cardiovascular Development and Diseases. <i>Journal of Developmental Biology</i> , 2016 , 4,	3.5	14
52	Cardiac outflow morphogenesis depends on effects of retinoic acid signaling on multiple cell lineages. <i>Developmental Dynamics</i> , 2016 , 245, 388-401	2.9	21
51	Msx1CreERT2 knock-In allele: A useful tool to target embryonic and adult cardiac valves. <i>Genesis</i> , 2015 , 53, 337-45	1.9	7
50	Hoxb1 regulates proliferation and differentiation of second heart field progenitors in pharyngeal mesoderm and genetically interacts with Hoxa1 during cardiac outflow tract development. <i>Developmental Biology</i> , 2015 , 406, 247-58	3.1	33
49	Retinoic Acid Signaling and Heart Development 2015 , 353-369		2
48	Anterior Hox Genes in Cardiac Development and Great Artery Patterning. <i>Journal of Cardiovascular Development and Disease</i> , 2014 , 1, 3-13	4.2	4
47	Retinoids and Cardiac Development. <i>Journal of Developmental Biology</i> , 2014 , 2, 50-71	3.5	16
46	Loss of Krox20 results in aortic valve regurgitation and impaired transcriptional activation of fibrillar collagen genes. <i>Cardiovascular Research</i> , 2014 , 104, 443-55	9.9	16
45	Tbx1 coordinates addition of posterior second heart field progenitor cells to the arterial and venous poles of the heart. <i>Circulation Research</i> , 2014 , 115, 790-9	15.7	72
44	Genetic lineage tracing analysis of anterior Hox expressing cells. <i>Methods in Molecular Biology</i> , 2014 , 1196, 37-48	1.4	4
43	Early cardiac development: a view from stem cells to embryos. <i>Cardiovascular Research</i> , 2012 , 96, 352-62	9.9	88
42	ISL1 directly regulates FGF10 transcription during human cardiac outflow formation. <i>PLoS ONE</i> , 2012 , 7, e30677	3.7	35
41	New developments in the second heart field. <i>Differentiation</i> , 2012 , 84, 17-24	3.5	65
40	Fibroblast growth factor 10 gene regulation in the second heart field by Tbx1, Nkx2-5, and Islet1 reveals a genetic switch for down-regulation in the myocardium. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 18273-80	11.5	90
39	Value of in vivo T2 measurement for myocardial fibrosis assessment in diabetic mice at 11.75 T. <i>Investigative Radiology</i> , 2012 , 47, 319-23	10.1	29
38	Hox genes define distinct progenitor sub-domains within the second heart field. <i>Developmental Biology</i> , 2011 , 353, 266-74	3.1	106

37	A retinoic acid responsive <i>Hoxa3</i> transgene expressed in embryonic pharyngeal endoderm, cardiac neural crest and a subdomain of the second heart field. <i>PLoS ONE</i> , 2011 , 6, e27624	3.7	20
36	Decreased levels of embryonic retinoic acid synthesis accelerate recovery from arterial growth delay in a mouse model of DiGeorge syndrome. <i>Circulation Research</i> , 2010 , 106, 686-94	15.7	67
35	Endogenous retinoic acid regulates cardiac progenitor differentiation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 9234-9	11.5	80
34	Expression of <i>Slit</i> and <i>Robo</i> genes in the developing mouse heart. <i>Developmental Dynamics</i> , 2010 , 239, 3303-11	2.9	29
33	Conotruncal defects associated with anomalous pulmonary venous connections. <i>Archives of Cardiovascular Diseases</i> , 2009 , 102, 105-10	2.7	23
32	Genetics and embryological mechanisms of congenital heart diseases. <i>Archives of Cardiovascular Diseases</i> , 2009 , 102, 59-63	2.7	28
31	Atrial myocardium derives from the posterior region of the second heart field, which acquires left-right identity as <i>Pitx2c</i> is expressed. <i>Development (Cambridge)</i> , 2008 , 135, 1157-67	6.6	115
30	Myocardium at the base of the aorta and pulmonary trunk is prefigured in the outflow tract of the heart and in subdomains of the second heart field. <i>Developmental Biology</i> , 2008 , 313, 25-34	3.1	53
29	Retinoic acid deficiency alters second heart field formation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 2913-8	11.5	160
28	<i>Fgf10</i> and the Embryological Origin of Outflow Tract Myocardium 2007 , 81-83		
27	Correction: <i>Pax3</i> and <i>Pax7</i> have distinct and overlapping functions in adult muscle progenitor cells. <i>Journal of Cell Biology</i> , 2007 , 176, 125-125	7.3	78
26	The <i>Drosophila Hand</i> gene is required for remodeling of the developing adult heart and midgut during metamorphosis. <i>Developmental Biology</i> , 2007 , 311, 287-96	3.1	24
25	An <i>Nkx2-5/Bmp2/Smad1</i> negative feedback loop controls heart progenitor specification and proliferation. <i>Cell</i> , 2007 , 128, 947-59	56.2	418
24	Rotation of the myocardial wall of the outflow tract is implicated in the normal positioning of the great arteries. <i>Circulation Research</i> , 2006 , 98, 421-8	15.7	162
23	Congenital heart defects in <i>Fgfr2-IIIb</i> and <i>Fgf10</i> mutant mice. <i>Cardiovascular Research</i> , 2006 , 71, 50-60	9.9	75
22	Cardioblast-intrinsic Tinman activity controls proper diversification and differentiation of myocardial cells in <i>Drosophila</i> . <i>Development (Cambridge)</i> , 2006 , 133, 4073-83	6.6	70
21	Development of the Larval Visceral Musculature 2006 , 62-78		6
20	<i>Pax3</i> and <i>Pax7</i> have distinct and overlapping functions in adult muscle progenitor cells. <i>Journal of Cell Biology</i> , 2006 , 172, 91-102	7.3	500

19	Direct isolation of satellite cells for skeletal muscle regeneration. <i>Science</i> , 2005 , 309, 2064-7	33.3	821
18	The homeodomain of Tinman mediates homo- and heterodimerization of NK proteins. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 334, 361-9	3.4	16
17	Building the mammalian heart from two sources of myocardial cells. <i>Nature Reviews Genetics</i> , 2005 , 6, 826-35	30.1	914
16	Fgf10 expression identifies parabronchial smooth muscle cell progenitors and is required for their entry into the smooth muscle cell lineage. <i>Development (Cambridge)</i> , 2005 , 132, 2157-66	6.6	151
15	Right ventricular myocardium derives from the anterior heart field. <i>Circulation Research</i> , 2004 , 95, 261-8	15.7	292
14	La Souris comme modèle d'étude de la morphogénèse du cœur chez les mammifères : origine des myocytes et études d'explants cardiaques. <i>Société De Biologie Journal</i> , 2003 , 197, 187-194		1
13	Cell history determines the maintenance of transcriptional differences between left and right ventricular cardiomyocytes in the developing mouse heart. <i>Journal of Cell Science</i> , 2003 , 116, 5005-13	5.3	7
12	Early signals in cardiac development. <i>Circulation Research</i> , 2002 , 91, 457-69	15.7	246
11	The beta 3 tubulin gene is a direct target of bagpipe and biniou in the visceral mesoderm of Drosophila. <i>Mechanisms of Development</i> , 2002 , 114, 85-93	1.7	15
10	Cardiogenesis in the Drosophila model: control mechanisms during early induction and diversification of cardiac progenitors. <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 2002 , 67, 1-12	3.9	22
9	Pericardin, a Drosophila type IV collagen-like protein is involved in the morphogenesis and maintenance of the heart epithelium during dorsal ectoderm closure. <i>Development (Cambridge)</i> , 2002 , 129, 3241-53	6.6	48
8	biniou (FoxF), a central component in a regulatory network controlling visceral mesoderm development and midgut morphogenesis in Drosophila. <i>Genes and Development</i> , 2001 , 15, 2900-15	12.6	95
7	Molecular cloning and embryonic expression of dFKBP59, a novel Drosophila FK506-binding protein. <i>Gene</i> , 2000 , 246, 103-9	3.8	15
6	The NK-2 homeobox gene scarecrow (<i>scro</i>) is expressed in pharynx, ventral nerve cord and brain of Drosophila embryos. <i>Mechanisms of Development</i> , 2000 , 94, 237-41	1.7	42
5	The heterotrimeric protein Go is required for the formation of heart epithelium in Drosophila. <i>Journal of Cell Biology</i> , 1999 , 145, 1063-76	7.3	67
4	Cellular interactions during heart morphogenesis in the Drosophila embryo. <i>Biology of the Cell</i> , 1995 , 84, 13-24	3.5	28
3	Hox-Dependent Coordination of Cardiac Cell Patterning and Differentiation. <i>SSRN Electronic Journal</i> ,	1	1
2	Hox-dependent coordination of cardiac progenitor cell patterning and differentiation		2

1 Piezo1 is required for outflow tract and aortic valve development

3