

Il-ho Jang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2787498/publications.pdf>

Version: 2024-02-01

52
papers

2,861
citations

236925

25
h-index

214800

47
g-index

53
all docs

53
docs citations

53
times ranked

5081
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Revisiting the role of lysophosphatidic acid in stem cell biology. <i>Experimental Biology and Medicine</i> , 2021, 246, 1802-1809. | 2.4 | 4 |
| 2 | Facilitation of Bone Healing Processes Based on the Developmental Function of Meox2 in Tooth Loss Lesion. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8701. | 4.1 | 7 |
| 3 | Oncostatin M enhances osteogenic differentiation of dental pulp stem cells derived from supernumerary teeth. <i>Biochemical and Biophysical Research Communications</i> , 2020, 529, 169-174. | 2.1 | 4 |
| 4 | NELL2 Function in Axon Development of Hippocampal Neurons. <i>Molecules and Cells</i> , 2020, 43, 581-589. | 2.6 | 2 |
| 5 | Cancer upregulated gene 2 (CUG2), a novel oncogene, promotes stemness-like properties via the NPM1-TGF- β^2 signaling axis. <i>Biochemical and Biophysical Research Communications</i> , 2019, 514, 1278-1284. | 2.1 | 13 |
| 6 | Inhibitory Effect of KP-A038 on Osteoclastogenesis and Inflammatory Bone Loss Is Associated With Downregulation of Blimp1. <i>Frontiers in Pharmacology</i> , 2019, 10, 367. | 3.5 | 15 |
| 7 | N-Acetylated Proline-Glycine-Proline Accelerates Cutaneous Wound Healing and Neovascularization by Human Endothelial Progenitor Cells. <i>Scientific Reports</i> , 2017, 7, 43057. | 3.3 | 28 |
| 8 | Functional expression and pharmaceutical efficacy of cardiac-specific ion channels in human embryonic stem cell-derived cardiomyocytes. <i>Scientific Reports</i> , 2017, 7, 13821. | 3.3 | 3 |
| 9 | Role of KrÄ¶ppel-Like Factor 4 in the Maintenance of Chemoresistance of Anaplastic Thyroid Cancer. <i>Thyroid</i> , 2017, 27, 1424-1432. | 4.5 | 22 |
| 10 | Trib2 regulates the pluripotency of embryonic stem cells and enhances reprogramming efficiency. <i>Experimental and Molecular Medicine</i> , 2017, 49, e401-e401. | 7.7 | 17 |
| 11 | Formyl Peptide Receptor 2 Is Involved in Cardiac Repair After Myocardial Infarction Through Mobilization of Circulating Angiogenic Cells. <i>Stem Cells</i> , 2017, 35, 654-665. | 3.2 | 33 |
| 12 | Identification of a novel angiogenic peptide from periostin. <i>PLoS ONE</i> , 2017, 12, e0187464. | 2.5 | 12 |
| 13 | Role of TAZ in Lysophosphatidic Acid-Induced Migration and Proliferation of Human Adipose-Derived Mesenchymal Stem Cells. <i>Biomolecules and Therapeutics</i> , 2017, 25, 354-361. | 2.4 | 5 |
| 14 | The anti-microbial peptide SR-0379 stimulates human endothelial progenitor cell-mediated repair of peripheral artery diseases. <i>BMB Reports</i> , 2017, 50, 504-509. | 2.4 | 3 |
| 15 | Selective Fluidization of Synaptosomal Plasma Membrane Vesicles by 17 β -Estradiol. <i>Biomedical Science Letters</i> , 2017, 23, 17-24. | 0.3 | 0 |
| 16 | FOXP1 functions as an oncogene in promoting cancer stem cell-like characteristics in ovarian cancer cells. <i>Oncotarget</i> , 2016, 7, 3506-3519. | 1.8 | 65 |
| 17 | Autotaxin Regulates Maintenance of Ovarian Cancer Stem Cells through Lysophosphatidic Acid-Mediated Autocrine Mechanism. <i>Stem Cells</i> , 2016, 34, 551-564. | 3.2 | 90 |
| 18 | Biomedical therapy using synthetic WKYMMV hexapeptide. <i>Organogenesis</i> , 2016, 12, 53-60. | 1.2 | 9 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Crucial role of HMGA1 in the self-renewal and drug resistance of ovarian cancer stem cells. <i>Experimental and Molecular Medicine</i> , 2016, 48, e255-e255. | 7.7 | 51 |
| 20 | Phospholipid End-Capped Bioreducible Polyurea Micelles as a Potential Platform for Intracellular Drug Delivery of Doxorubicin in Tumor Cells. <i>ACS Biomaterials Science and Engineering</i> , 2016, 2, 1883-1893. | 5.2 | 10 |
| 21 | Synthesis and Characterization of Water-Soluble Conjugated Oligoelectrolytes for Near-Infrared Fluorescence Biological Imaging. <i>ACS Applied Materials & Interfaces</i> , 2016, 8, 15937-15947. | 8.0 | 29 |
| 22 | Hypoxia-NOTCH1-SOX2 signaling is important for maintaining cancer stem cells in ovarian cancer. <i>Oncotarget</i> , 2016, 7, 55624-55638. | 1.8 | 84 |
| 23 | A FRET Assay for Celiac Disease. <i>Biomedical Science Letters</i> , 2016, 22, 160-166. | 0.3 | 0 |
| 24 | Stimulation of cutaneous wound healing by an FPR2-specific peptide agonist WKYMVm. <i>Wound Repair and Regeneration</i> , 2015, 23, 575-582. | 3.0 | 22 |
| 25 | Injectable PLGA microspheres encapsulating WKYMVM peptide for neovascularization. <i>Acta Biomaterialia</i> , 2015, 25, 76-85. | 8.3 | 23 |
| 26 | Notch1 acts via Foxc2 to promote definitive hematopoiesis via effects on hemogenic endothelium. <i>Blood</i> , 2015, 125, 1418-1426. | 1.4 | 40 |
| 27 | Role of formyl peptide receptor 2 in homing of endothelial progenitor cells and therapeutic angiogenesis. <i>Advances in Biological Regulation</i> , 2015, 57, 162-172. | 2.3 | 12 |
| 28 | Periostin Accelerates Bone Healing Mediated by Human Mesenchymal Stem Cell-Embedded Hydroxyapatite/Tricalcium Phosphate Scaffold. <i>PLoS ONE</i> , 2015, 10, e0116698. | 2.5 | 32 |
| 29 | Isolation of Foreign Material-Free Endothelial Progenitor Cells Using CD31 Aptamer and Therapeutic Application for Ischemic Injury. <i>PLoS ONE</i> , 2015, 10, e0131785. | 2.5 | 21 |
| 30 | Therapeutic angiogenesis in a murine model of limb ischemia by recombinant periostin and its fasciclin I domain. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 1324-1332. | 3.8 | 22 |
| 31 | WKYMVm-Induced Activation of Formyl Peptide Receptor 2 Stimulates Ischemic Neovasculogenesis by Promoting Homing of Endothelial Colony-Forming Cells. <i>Stem Cells</i> , 2014, 32, 779-790. | 3.2 | 69 |
| 32 | Reptin Regulates Pluripotency of Embryonic Stem Cells and Somatic Cell Reprogramming Through Oct4-Dependent Mechanism. <i>Stem Cells</i> , 2014, 32, 3126-3136. | 3.2 | 10 |
| 33 | Tumor necrosis factor- α -activated mesenchymal stem cells promote endothelial progenitor cell homing and angiogenesis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2013, 1832, 2136-2144. | 3.8 | 112 |
| 34 | Signaling axis involving Hedgehog, Notch, and Scl promotes the embryonic endothelial-to-hematopoietic transition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E141-E150. | 7.1 | 58 |
| 35 | Efficient Production of Retroviruses Using PLGA/bPEI-DNA Nanoparticles and Application for Reprogramming Somatic Cells. <i>PLoS ONE</i> , 2013, 8, e76875. | 2.5 | 10 |
| 36 | Metabolic Regulation of Protein N-Alpha-Acetylation by Bcl-xL Promotes Cell Survival. <i>Cell</i> , 2011, 146, 607-620. | 28.9 | 185 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | Neonatal Recipients Offer Permissive Hematopoietic Microenvironment for Engraftment of Embryonic Murine Hematopoietic Stem Cells. <i>Blood</i> , 2011, 118, 2344-2344. | 1.4 | 0 |
| 38 | Cdk5 phosphorylates PLD2 to mediate EGF-dependent insulin secretion. <i>Cellular Signalling</i> , 2008, 20, 1787-1794. | 3.6 | 40 |
| 39 | BMP and Wnt Specify Hematopoietic Fate by Activation of the Cdx-Hox Pathway. <i>Cell Stem Cell</i> , 2008, 2, 72-82. | 11.1 | 192 |
| 40 | Modulation of murine embryonic stem cell-derived CD41+c-kit+ hematopoietic progenitors by ectopic expression of Cdx genes. <i>Blood</i> , 2008, 111, 4944-4953. | 1.4 | 48 |
| 41 | Prostaglandin E2 regulates vertebrate haematopoietic stem cell homeostasis. <i>Nature</i> , 2007, 447, 1007-1011. | 27.8 | 1,037 |
| 42 | BMP Signaling Via the Cdx-Hox Pathway Allocates Mesoderm to Hematopoietic vs Cardiac Fates.. <i>Blood</i> , 2006, 108, 4183-4183. | 1.4 | 0 |
| 43 | Prostaglandin E2 Is a Potent Regulator of Vertebrate Hematopoietic Stem Cell Homeostasis.. <i>Blood</i> , 2006, 108, 680-680. | 1.4 | 0 |
| 44 | Phosphatidylinositol (3,4,5)-trisphosphate specifically interacts with the phox homology domain of phospholipase D1 and stimulates its activity. <i>Journal of Cell Science</i> , 2005, 118, 4405-4413. | 2.0 | 53 |
| 45 | Munc-18-1 Inhibits Phospholipase D Activity by Direct Interaction in an Epidermal Growth Factor-reversible Manner. <i>Journal of Biological Chemistry</i> , 2004, 279, 16339-16348. | 3.4 | 33 |
| 46 | Sensitization of Epidermal Growth Factor-induced Signaling by Bradykinin Is Mediated by c-Src. <i>Journal of Biological Chemistry</i> , 2004, 279, 5852-5860. | 3.4 | 65 |
| 47 | Localization of VEGFR-2 and PLD2 in endothelial caveolae is involved in VEGF-induced phosphorylation of MEK and ERK. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2004, 286, H1881-H1888. | 3.2 | 62 |
| 48 | Regulation of phospholipase D2 by GTP-dependent interaction with dynamin. <i>Advances in Enzyme Regulation</i> , 2004, 44, 249-264. | 2.6 | 7 |
| 49 | Localization of Tie2 and phospholipase D in endothelial caveolae is involved in angiopoietin-1-induced MEK/ERK phosphorylation and migration in endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2003, 308, 101-105. | 2.1 | 38 |
| 50 | The Direct Interaction of Phospholipase C- β 1 with Phospholipase D2 Is Important for Epidermal Growth Factor Signaling. <i>Journal of Biological Chemistry</i> , 2003, 278, 18184-18190. | 3.4 | 48 |
| 51 | Localization of phospholipase C- β 1 signaling in caveolae: importance in EGF-induced phosphoinositide hydrolysis but not in tyrosine phosphorylation. <i>FEBS Letters</i> , 2001, 491, 4-8. | 2.8 | 35 |
| 52 | Phospholipase D1 Is Phosphorylated and Activated by Protein Kinase C in Caveolin-enriched Microdomains within the Plasma Membrane. <i>Journal of Biological Chemistry</i> , 2000, 275, 13621-13627. | 3.4 | 76 |