Nobuyuki Miyatake

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/279664/publications.pdf

Version: 2024-02-01

73 papers

454 citations

933447 10 h-index 940533 16 g-index

73 all docs

73 docs citations

times ranked

73

700 citing authors

#	Article	IF	CITATIONS
1	Preterm birth is strongly affected by the glucocorticoid dose during pregnancy in women complicated by systemic lupus erythematosus. Arthritis Research and Therapy, 2022, 24, 10.	3.5	13
2	Association Between Temporal Changes in Diet Quality and Concurrent Changes in Dietary Intake, Body Mass Index, and Physical Activity Among Japanese Adults: A Longitudinal Study. Frontiers in Nutrition, 2022, 9, 753127.	3.7	5
3	Rationale, Design and Baseline Characteristics of the Effect of Canagliflozin in Type 2 Diabetic Patients with Microalbuminuria in Japanese Population (<scp>CANPIONE</scp>) study. Diabetes, Obesity and Metabolism, 2022, , .	4.4	1
4	Relation between Prolonged Sedentary Bouts and Health-Related Quality of Life in Patients on Chronic Hemodialysis Acta Medica Okayama, 2022, 76, 113-119.	0.2	0
5	Relationship among psychological distress, food dependence, and the time discount rate: a pilot mediation analysis. Psychology, Health and Medicine, 2021, 26, 1005-1012.	2.4	O
6	Health-related quality of life in Japanese low-risk prostate cancer patients choosing active surveillance: 3-year follow-up from PRIAS-JAPAN. World Journal of Urology, 2021, 39, 2491-2497.	2.2	9
7	Relationship between the Number of Deaths Due to Renal Failure and Air Temperature Parameters in Hokkaido and Okinawa Prefectures, Japan. Epidemiologia, 2021, 2, 68-74.	2.2	0
8	Disaster Prevention Education along with Weekly Exercise Improves Self-Efficacy in Community-Dwelling Japanese People—A Randomized Control Trial. Medicina (Lithuania), 2021, 57, 231.	2.0	1
9	Reproducibility and prognostic significance of area of residual tumor (ART) in post-neoadjuvant resections of pancreatic ductal adenocarcinoma. Pancreatology, 2021, 21, 1506-1515.	1.1	3
10	Effect of a 1-year intervention comprising brief counselling sessions and low-dose physical activity recommendations in Japanese adults, and retention of the effect at 2Âyears: a randomized trial. BMC Sports Science, Medicine and Rehabilitation, 2021, 13, 133.	1.7	5
11	A Text Mining-Based Survey of Pre-Impressions of Medical Staff toward COVID-19 Vaccination in a Designated Medical Institution for Class II Infectious Diseases. Vaccines, 2021, 9, 1282.	4.4	4
12	The Association between Chronic Hemodialysis and Toe Pinch Force in Japanese Patients: Cross-Sectional Study. Healthcare (Switzerland), 2021, 9, 1745.	2.0	1
13	Does physical activity attenuate the association between birth weight and glycated hemoglobin in nondiabetic Japanese women?. Journal of Developmental Origins of Health and Disease, 2020, 11, 379-383.	1.4	1
14	Determination of metallization with energy-dispersive X-ray fluorescent spectrometry in experimental electric injury. Legal Medicine, 2020, 47, 101768.	1.3	0
15	Face-to-Face Exercises Performed by Instructors to Improve the Mental Health of Japanese in the Community—A Randomized Control Trial. Medicina (Lithuania), 2020, 56, 404.	2.0	O
16	Seasonal Changes in Continuous Sedentary Behavior in Community—Dwelling Japanese Adults: A Pilot Study. Medicines (Basel, Switzerland), 2020, 7, 48.	1.4	1
17	Changes and Variations in Death Due to Senility in Japan. Healthcare (Switzerland), 2020, 8, 443.	2.0	3
18	Factors affecting the Apgar score of offsprings born to mothers suffering from systemic lupus erythematosus. Medicine (United States), 2020, 99, e22843.	1.0	4

#	Article	IF	CITATIONS
19	Locomotive Syndrome is a Risk Factor for the Dropout of Continuous Positive Airway Pressure Treatment in Patients with Obstructive Sleep Apnea Syndrome. Healthcare (Switzerland), 2020, 8, 177.	2.0	1
20	An exploratory study of relationship between lower-limb muscle mass and diabetic polyneuropathy in patients with type 2 diabetes. Journal of Diabetes and Metabolic Disorders, 2020, 19, 281-287.	1.9	1
21	Relationship between Air Temperature Parameters and the Number of Deaths Stratified by Cause in Gifu Prefecture, Japan. Healthcare (Switzerland), 2020, 8, 35.	2.0	6
22	Decrease in blood donation rates in Japan: a time series analysis. Revista Brasileira De Epidemiologia, 2020, 23, e200047.	0.8	1
23	Examination of the Causality Relationship between Psychological Distress and Social Participation in Elderly People: A Pilot Study. Acta Medica Okayama, 2020, 74, 59-63.	0.2	2
24	Relationships between Physical Activity, Sleeping Time, and Psychological Distress in Community-Dwelling Elderly Japanese. Medicina (Lithuania), 2019, 55, 318.	2.0	5
25	Behavior problems and dysfunctional parenting: Crossâ€sectional study in Japan. Pediatrics International, 2019, 61, 1096-1102.	0.5	7
26	Relationship between Psychological Distress and Continuous Sedentary Behavior in Healthy Older Adults. Medicina (Lithuania), 2019, 55, 324.	2.0	3
27	Influence of Smartphone Usage on Low Back Pain and/or Shoulder Pain in College Students in a Judo Therapist Training Course: A pilot study. Journal of Allied Health Sciences, 2019, 10, 1-9.	0.0	1
28	A Pilot Study of the Relationship between Diet and Mental Health in Community Dwelling Japanese Women. Medicina (Lithuania), 2019, 55, 513.	2.0	3
29	Long-Term Relationship between Psychological Distress and Continuous Sedentary Behavior in Healthy Older Adults: A Three Panel Study. Medicina (Lithuania), 2019, 55, 555.	2.0	2
30	Prediction of Chronic Lower Back Pain Using the Hierarchical Neural Network: Comparison with Logistic Regression—A Pilot Study. Medicina (Lithuania), 2019, 55, 259.	2.0	5
31	Relationship between Food Dependence and Nicotine Dependence in Smokers: A Cross-Sectional Study of Staff and Students at Medical Colleges. Medicina (Lithuania), 2019, 55, 202.	2.0	3
32	Effect of biologic disease-modifying anti-rheumatic drugs for patients with rheumatoid arthritis who hope to become mothers. Clinical Rheumatology, 2019, 38, 1453-1458.	2.2	12
33	Sub-analysis of the prevalence of locomotive syndrome and its relationship with health-related quality of life in patients with obstructive sleep apnea syndrome as classified by age and sex. Sleep and Biological Rhythms, 2019, 17, 149-153.	1.0	4
34	Relationship between Sedentary Behavior and All-cause Mortality in Japanese Chronic Hemodialysis Patients: A Prospective Cohort Study. Acta Medica Okayama, 2019, 73, 419-425.	0.2	7
35	"Active Guide" Brochure Reduces Sedentary Behavior of Elderly People: A Randomized Controlled Trial. Acta Medica Okayama, 2019, 73, 427-432.	0.2	2
36	Comparison of histological findings and the results of energy-dispersive X-ray spectrometry analysis in experimental electrical injury. Legal Medicine, 2018, 31, 20-23.	1.3	2

3

#	Article	IF	CITATIONS
37	Decrease in toe pinch force in male type 2 diabetic patients with diabetic nephropathy. Clinical and Experimental Nephrology, 2018, 22, 647-652.	1.6	3
38	A pilot study of short-term toe resistance training in patients with type 2 diabetes mellitus. Diabetology International, 2017, 8, 392-396.	1.4	4
39	A pilot study evaluating the factors associated with psychological distress of clinical psychologists in Kagawa Prefecture, Japan. Journal of Health Psychology Research, 2017, 29, 53-59.	0.0	0
40	The i> IL18 i> Promoter Polymorphism, rs1946518, Is Associated with the Risk of Periodontitis in Japanese Women: The Kyushu Okinawa Maternal and Child Health Study. Tohoku Journal of Experimental Medicine, 2017, 243, 159-164.	1.2	14
41	Greater Progression of Age-Related Aortic Stiffening in Adults with Poor Trunk Flexibility: A 5-Year Longitudinal Study. Frontiers in Physiology, 2017, 8, 454.	2.8	8
42	Relationship of locomotive syndrome with health-related quality of life among patients with obstructive sleep apnea syndrome. Journal of Physical Therapy Science, 2017, 29, 1129-1133.	0.6	5
43	Psychological distress between chronic hemodialysis patients with and without low back pain. Environmental Health and Preventive Medicine, 2016, 21, 487-491.	3.4	3
44	Cardiorespiratory Fitness Suppresses Ageâ€Related Arterial Stiffening in Healthy Adults: A 2‥ear Longitudinal Observational Study. Journal of Clinical Hypertension, 2016, 18, 292-298.	2.0	31
45	Prevalence of attention deficit hyperactivity disorder and/or autism spectrum disorder and its relation to lifestyle in female college students. Environmental Health and Preventive Medicine, 2016, 21, 455-459.	3.4	6
46	Relationship of toe pinch force to other muscle strength parameters in men with type 2 diabetes. Environmental Health and Preventive Medicine, 2016, 21, 179-185.	3.4	14
47	A pilot study of the relationship between diet and mental health in female university students enrolled in a training course for registered dietitians. Environmental Health and Preventive Medicine, 2016, 21, 345-349.	3.4	4
48	Comparison of health-related quality of life between type 2 diabetic patients with and without locomotive syndrome. Environmental Health and Preventive Medicine, 2016, 21, 356-360.	3.4	12
49	Correlation between suicide and meteorological parameters. Medicina (Lithuania), 2015, 51, 363-367.	2.0	11
50	Detection of chlorine and bromine in free liquid from the sphenoid sinus as an indicator of seawater drowning. Legal Medicine, 2015, 17, 299-303.	1.3	11
51	Higher vitamin D intake during pregnancy is associated with reduced risk of dental caries in young Japanese children. Annals of Epidemiology, 2015, 25, 620-625.	1.9	45
52	Evaluation of antireflux surgery using multichannel intraluminal impedance-pH measurement in neurologically impaired patients. Pediatric Surgery International, 2015, 31, 911-916.	1.4	3
53	Circulating leptin levels are associated with physical activity or physical fitness in Japanese. Environmental Health and Preventive Medicine, 2014, 19, 362-366.	3.4	15
54	Serum vaspin levels are associated with physical activity or physical fitness in Japanese: a pilot study. Environmental Health and Preventive Medicine, 2014, 19, 200-206.	3.4	4

#	Article	IF	CITATIONS
55	Comparison of serum ferritin and oxidative stress biomarkers between Japanese workers with and without metabolic syndrome. Obesity Research and Clinical Practice, 2014, 8, e271-e282.	1.8	21
56	Circulating adiponectin levels are associated with peak oxygen uptake in Japanese. Environmental Health and Preventive Medicine, 2014, 19, 279-285.	3.4	8
57	Lifestyle modification is associated with improving estimated glomerular filtration rate (eGFR) and proteinuria in Japanese with proteinuria. Acta Medica Okayama, 2014, 68, 43-6.	0.2	0
58	Protective effect of D-psicose against testicular atrophy induced by di(2-ethylhexyl) phthalate. International Journal of Environmental Studies, 2013, 70, 560-565.	1.6	1
59	Serum Interleukin-18 Levels Are Associated with Physical Activity in Japanese Men. PLoS ONE, 2013, 8, e81497.	2.5	6
60	Association between dietary intake of micronutrients and cardiorespiratory fitness in Japanese men. Journal of Nutritional Science, 2012, 1, e12.	1.9	12
61	Evaluation of anthropometric parameters and physical fitness in elderly Japanese. Environmental Health and Preventive Medicine, 2012, 17, 62-68.	3.4	4
62	The relation between ambulance transports stratified by heat stroke and air temperature in all 47 prefectures of Japan in August, 2009: ecological study. Environmental Health and Preventive Medicine, 2012, 17, 77-80.	3.4	20
63	Simple and Sensitive Headspace Gas Chromatographic Determination of Urinary Methanol by Means of Syringe Equilibration. Bunseki Kagaku, 2011, 60, 865-868.	0.2	0
64	Comparison of the changes in temperatures among rural, urban and metropolitan areas around the Inland Sea in Japan. Environmental Monitoring and Assessment, 2011, 181, 525-530.	2.7	8
65	Relation between cigarette smoking and ventilatory threshold in the Japanese. Environmental Health and Preventive Medicine, 2011, 16, 185-190.	3.4	8
66	The relation between estimated glomerular filtration rate and proteinuria in Okayama Prefecture, Japan. Environmental Health and Preventive Medicine, 2011, 16, 191-195.	3.4	5
67	Decreasing abdominal circumference is associated with improving estimated glomerular filtration rate (eGFR) with lifestyle modification in Japanese men: a pilot study. Acta Medica Okayama, 2011, 65, 363-7.	0.2	5
68	Relation between the Estimated Glomerular Filtration Rate and Pulse Wave Velocity in Japanese. Internal Medicine, 2010, 49, 1315-1320.	0.7	10
69	Changes in Metabolic Syndrome and Its Components with Lifestyle Modification in Japanese Men. Internal Medicine, 2010, 49, 261-265.	0.7	7
70	Evaluation of ventilatory threshold and its relation to exercise habits among Japanese. Environmental Health and Preventive Medicine, 2010, 15, 374-380.	3.4	7
71	Relationship between estimated glomerular filtration rate (eGFR) and metabolic syndrome in Japanese. Acta Medica Okayama, 2010, 64, 203-8.	0.2	9
72	Decreasing systolic blood pressure is associated with improving estimated glomerular filtration rate (eGFR) with lifestyle modification in Japanese healthy women. Acta Medica Okayama, 2010, 64, 339-43.	0.2	5

#	Article	lF	CITATIONS
73	Influence of cigarette smoking on estimated glomerular filtration rate (eGFR) in Japanese male workers. Acta Medica Okayama, 2010, 64, 385-90.	0.2	7