

Yoshihiro Tatsumi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2806974/publications.pdf>

Version: 2024-02-01

42
papers

957
citations

516710

16
h-index

477307

29
g-index

44
all docs

44
docs citations

44
times ranked

1611
citing authors

#	ARTICLE	IF	CITATIONS
1	CXCL1-Mediated Interaction of Cancer Cells with Tumor-Associated Macrophages and Cancer-Associated Fibroblasts Promotes Tumor Progression in Human Bladder Cancer. <i>Neoplasia</i> , 2016, 18, 636-646.	5.3	161
2	Collagen type IV alpha 1 (COL4A1) and collagen type XIII alpha 1 (COL13A1) produced in cancer cells promote tumor budding at the invasion front in human urothelial carcinoma of the bladder. <i>Oncotarget</i> , 2017, 8, 36099-36114.	1.8	76
3	Regulatory T Cells and Tumor-Associated Macrophages in the Tumor Microenvironment in Non-Muscle Invasive Bladder Cancer Treated with Intravesical Bacille Calmette-Guérin: A Long-Term Follow-Up Study of a Japanese Cohort. <i>International Journal of Molecular Sciences</i> , 2017, 18, 2186.	4.1	71
4	Clinical impact of postoperative loss in psoas major muscle and nutrition index after radical cystectomy for patients with urothelial carcinoma of the bladder. <i>BMC Cancer</i> , 2017, 17, 237.	2.6	68
5	Neutrophil-to-lymphocyte ratio as a detection marker of tumor recurrence in patients with muscle-invasive bladder cancer after radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016, 34, 257.e11-257.e17.	1.6	44
6	Exploration of risk factors predicting outcomes for primary T1 high-grade bladder cancer and validation of the Spanish Urological Club for Oncological Treatment scoring model: Long-term follow-up experience at a single institute. <i>International Journal of Urology</i> , 2015, 22, 541-547.	1.0	43
7	Diagnostic and prognostic role of urinary collagens in primary human bladder cancer. <i>Cancer Science</i> , 2017, 108, 2221-2228.	3.9	36
8	Preoperative predictive factors focused on inflammation-, nutrition-, and muscle-status in patients with upper urinary tract urothelial carcinoma undergoing nephroureterectomy. <i>International Journal of Clinical Oncology</i> , 2019, 24, 533-545.	2.2	33
9	Correlation of Immune Cells and Cytokines in the Tumor Microenvironment with Elevated Neutrophil-To-Lymphocyte Ratio in Blood: An Analysis of Muscle-Invasive Bladder Cancer. <i>Cancer Investigation</i> , 2018, 36, 395-405.	1.3	28
10	Prostate diseases and microbiome in the prostate, gut, and urine. <i>Prostate International</i> , 2022, 10, 96-107.	2.3	25
11	Expression of ferrochelatase has a strong correlation in protoporphyrin IX accumulation with photodynamic detection of bladder cancer. <i>Photodiagnosis and Photodynamic Therapy</i> , 2016, 13, 225-232.	2.6	23
12	Protoporphyrin IX induced by 5-aminolevulinic acid in bladder cancer cells in voided urine can be extracorporeally quantified using a spectrophotometer. <i>Photodiagnosis and Photodynamic Therapy</i> , 2015, 12, 282-288.	2.6	22
13	Tadalafil, a phosphodiesterase type 5 inhibitor, improves bladder blood supply and restores the initial phase of lower urinary tract dysfunction in diabetic rats. <i>Neurourology and Urodynamics</i> , 2018, 37, 666-672.	1.5	21
14	Gamma-Klotho exhibits multiple roles in tumor growth of human bladder cancer. <i>Oncotarget</i> , 2018, 9, 19508-19524.	1.8	21
15	Clinical significance of β - and β ² -Klotho in urothelial carcinoma of the bladder. <i>Oncology Reports</i> , 2016, 36, 2117-2125.	2.6	20
16	Changes in oncological outcomes after radical nephroureterectomy in patients with upper urinary tract urothelial carcinoma treated in the last two decades: a retrospective analysis based on a multicenter collaborative study. <i>Japanese Journal of Clinical Oncology</i> , 2016, 46, 1148-1155.	1.3	16
17	Outcomes of subsequent non-muscle-invasive bladder cancer treated with intravesical Bacillus Calmette-Guérin after radical nephroureterectomy for upper urinary tract urothelial carcinoma. <i>BJU International</i> , 2018, 121, 764-773.	2.5	16
18	The best objective response of target lesions and the incidence of treatment-related hypertension are associated with the survival of patients with metastatic renal cell carcinoma treated with sunitinib: a Japanese retrospective study. <i>BMC Research Notes</i> , 2016, 9, 79.	1.4	15

#	ARTICLE	IF	CITATIONS
19	Assessment of lower urinary symptom flare with overactive bladder symptom score and International Prostate Symptom Score in patients treated with iodine-125 implant brachytherapy: long-term follow-up experience at a single institute. <i>BMC Urology</i> , 2017, 17, 62.	1.4	15
20	Clinical Impact of Tumor-Infiltrating Lymphocytes and PD-L1-Positive Cells as Prognostic and Predictive Biomarkers in Urological Malignancies and Retroperitoneal Sarcoma. <i>Cancers</i> , 2020, 12, 3153.	3.7	15
21	Transperineal template-guided saturation biopsy aimed at sampling one core for each milliliter of prostate volume: 103 cases requiring repeat prostate biopsy. <i>BMC Urology</i> , 2017, 17, 28.	1.4	14
22	Biological Behavior and Long-Term Outcomes of Carcinoma In Situ in Upper Urinary Tract Managed by Radical Nephroureterectomy. <i>Journal of Urology</i> , 2018, 199, 933-939.	0.4	14
23	Topical and systemic immunoreaction triggered by intravesical chemotherapy in an N-butyl-N-(4-hydroxybutyl) nitrosamine induced bladder cancer mouse model. <i>PLoS ONE</i> , 2017, 12, e0175494.	2.5	13
24	Disabled Homolog 2 (DAB2) Protein in Tumor Microenvironment Correlates with Aggressive Phenotype in Human Urothelial Carcinoma of the Bladder. <i>Diagnostics</i> , 2020, 10, 54.	2.6	13
25	Changes in lower urinary tract symptoms and quality of life after salvage radiotherapy for biochemical recurrence of prostate cancer. <i>Radiotherapy and Oncology</i> , 2015, 115, 321-326.	0.6	12
26	Extended resection including adjacent organs and Ki-67 labeling index are prognostic factors in patients with retroperitoneal soft tissue sarcomas. <i>World Journal of Surgical Oncology</i> , 2016, 14, 43.	1.9	12
27	Expression of peptide transporter 1 has a positive correlation in protoporphyrin IX accumulation induced by 5-aminolevulinic acid with photodynamic detection of non-small cell lung cancer and metastatic brain tumor specimens originating from non-small cell lung cancer. <i>Photodiagnosis and Photodynamic Therapy</i> , 2019, 25, 309-316.	2.6	12
28	Clinical utility of bioelectrical impedance analysis in patients with locoregional muscle invasive or metastatic urothelial carcinoma: a subanalysis of changes in body composition during neoadjuvant systemic chemotherapy. <i>Supportive Care in Cancer</i> , 2018, 26, 1077-1086.	2.2	10
29	Evaluation of pro- and anti-tumor effects induced by three colony-stimulating factors, G-CSF, GM-CSF and M-CSF, in bladder cancer cells: Is G-CSF a friend of bladder cancer cells?. <i>International Journal of Oncology</i> , 2019, 54, 2237-2249.	3.3	10
30	Tumor Location Based Segmentation in Upper-Tract Urothelial Carcinoma Impacts on the Urothelial Recurrence-Free Survival: A Multi-Institutional Database Study. <i>Current Urology</i> , 2020, 14, 183-190.	0.6	9
31	Potential biomarkers for the therapeutic efficacy of sorafenib, sunitinib and everolimus. <i>Oncology Reports</i> , 2017, 37, 227-234.	2.6	8
32	Atypical small acinar proliferation and two or more cores of high-grade intraepithelial neoplasia on a previous prostate biopsy are significant predictors of cancer during a transperineal template-guided saturation biopsy aimed at sampling one core for each 1 mL of prostate volume. <i>Research and Reports in Urology</i> , 2017, Volume 9, 187-193.	1.0	8
33	Quality of life in patients who underwent 125I brachytherapy, 125I brachytherapy combined with three-dimensional conformal radiation therapy, or intensity-modulated radiation therapy, for prostate cancer. <i>Journal of Radiation Research</i> , 2019, 60, 270-280.	1.6	8
34	Inhibition of Heparanase Expression Results in Suppression of Invasion, Migration and Adhesion Abilities of Bladder Cancer Cells. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3789.	4.1	8
35	Klotho is correlated with resistance to docetaxel in castration-resistant prostate cancer. <i>Oncology Letters</i> , 2020, 19, 2306-2316.	1.8	6
36	Spectrophotometric photodynamic diagnosis of prostate cancer cells excreted in voided urine using 5-aminolevulinic acid. <i>Lasers in Medical Science</i> , 2018, 33, 1557-1563.	2.1	5

#	ARTICLE	IF	CITATIONS
37	<p>Comparison of cancer detection rates by transrectal prostate biopsy for prostate cancer using two different nomograms based on patient's age and prostate volume</p>. Research and Reports in Urology, 2019, Volume 11, 61-68.	1.0	5
38	Colony-stimulating factors detected in tumor cells and voided urine are potential prognostic markers for patients with muscle-invasive bladder cancer undergoing radical cystectomy. Research and Reports in Urology, 2018, Volume 10, 103-111.	1.0	4
39	Intravesical treatment of chemotherapeutic agents sensitizes bacillus Calmette-Guerin by the modulation of the tumor immune environment. Oncology Reports, 2019, 41, 1863-1874.	2.6	4
40	Orthotopic mouse model of renal pelvic cancer using a human bladder cancer cell line. International Journal of Urology, 2016, 23, 961-962.	1.0	3
41	Inhibitory Effect of Orally Administered 5-Aminolevulinic Acid on Prostate Carcinogenesis in the FVB-Transgenic Adenocarcinoma of a Mouse Prostate (FVB-TRAMP) Model. Asian Pacific Journal of Cancer Prevention, 2020, 21, 3743-3749.	1.2	2
42	Simple suturing of the bladder neck muscle layer at the vesicourethral anastomosis site to the dorsal vein complex during anterior reconstruction led to a better postoperative urinary continence after robot-assisted laparoscopic prostatectomy. Scandinavian Journal of Urology, 2020, 54, 470-474.	1.0	1