

Paula Clancy

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2814085/publications.pdf>

Version: 2024-02-01

39
papers

1,627
citations

279798

23
h-index

302126

39
g-index

39
all docs

39
docs citations

39
times ranked

2205
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Differential associations of ferritin and 25-hydroxyvitamin D with fasting glucose and diabetes risk in community dwelling older men. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3172. | 4.0 | 1 |
| 2 | High serum thrombospondin-1 concentration is associated with slower abdominal aortic aneurysm growth and deficiency of thrombospondin-1 promotes angiotensin II induced aortic aneurysm in mice. <i>Clinical Science</i> , 2017, 131, 1261-1281. | 4.3 | 26 |
| 3 | Involvement of Angiotensin II Type 1 and 2 Receptors in Gelatinase Regulation in Human Carotid Atheroma <i>in vitro</i> . <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 773-791. | 2.0 | 5 |
| 4 | Plasma ferritin concentrations are not associated with abdominal aortic aneurysm diagnosis, size or growth. <i>Atherosclerosis</i> , 2016, 251, 19-24. | 0.8 | 8 |
| 5 | The Impact of Prior Flavivirus Infections on the Development of Type 2 Diabetes Among the Indigenous Australians. <i>American Journal of Tropical Medicine and Hygiene</i> , 2016, 95, 265-268. | 1.4 | 2 |
| 6 | A Peptide Antagonist of Thrombospondin-1 Promotes Abdominal Aortic Aneurysm Progression in the Angiotensin II-Infused Apolipoprotein-E-Deficient Mouse. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 389-398. | 2.4 | 51 |
| 7 | Urocortin 2 is associated with abdominal aortic aneurysm and mediates anti-proliferative effects on vascular smooth muscle cells via corticotrophin releasing factor receptor 2. <i>Clinical Science</i> , 2014, 126, 517-527. | 4.3 | 27 |
| 8 | Serum Endostatin Concentrations Are Higher in Men with Symptoms of Intermittent Claudication. <i>Disease Markers</i> , 2014, 2014, 1-5. | 1.3 | 9 |
| 9 | Angiogenesis inhibition and depression in older men. <i>Journal of Psychiatry and Neuroscience</i> , 2014, 39, 200-205. | 2.4 | 11 |
| 10 | Tenascin-C is increased in atherothrombotic stroke patients and has an anti-inflammatory effect in the human carotid artery. <i>BioFactors</i> , 2014, 40, 448-457. | 5.4 | 9 |
| 11 | Angiotensin receptor 1 blockade reduces secretion of inflammation associated cytokines from cultured human carotid atheroma and vascular cells in association with reduced extracellular signal regulated kinase expression and activation. <i>Atherosclerosis</i> , 2014, 236, 108-115. | 0.8 | 37 |
| 12 | Plasma Angiotensin-1 Is Lower After Ischemic Stroke and Associated With Major Disability But Not Stroke Incidence. <i>Stroke</i> , 2014, 45, 1064-1068. | 2.0 | 22 |
| 13 | Role of the angiotensin converting enzyme 1/angiotensin II/angiotensin receptor 1 axis in interstitial collagenase expression in human carotid atheroma. <i>Atherosclerosis</i> , 2013, 229, 331-337. | 0.8 | 18 |
| 14 | Increased serum angiotensin-2 is associated with abdominal aortic aneurysm prevalence and cardiovascular mortality in older men. <i>International Journal of Cardiology</i> , 2013, 167, 1159-1163. | 1.7 | 18 |
| 15 | Relation Between Serum Thrombospondin-2 and Cardiovascular Mortality in Older Men Screened for Abdominal Aortic Aneurysm. <i>American Journal of Cardiology</i> , 2013, 111, 1800-1804. | 1.6 | 25 |
| 16 | Increased plasma levels of NGAL, a marker of neutrophil activation, in patients with abdominal aortic aneurysm. <i>Atherosclerosis</i> , 2012, 220, 552-556. | 0.8 | 52 |
| 17 | Proteomic analysis of intra-arterial thrombus secretions reveals a negative association of clusterin and thrombospondin-1 with abdominal aortic aneurysm. <i>Atherosclerosis</i> , 2011, 219, 432-439. | 0.8 | 42 |
| 18 | Circulating Concentrations of Stem-Cell-Mobilizing Cytokines Are Associated With Levels of Osteoprogenitor Cells and Aortic Calcification Severity. <i>Circulation Journal</i> , 2011, 75, 1227-1234. | 1.6 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Evaluation of the diagnostic and prognostic value of plasma D-dimer for abdominal aortic aneurysm. <i>European Heart Journal</i> , 2011, 32, 354-364. | 2.2 | 81 |
| 20 | The role of tenascin C in cardiovascular disease. <i>Cardiovascular Research</i> , 2011, 92, 19-28. | 3.8 | 68 |
| 21 | Association of PPAR α allelic variation, osteoprotegerin and abdominal aortic aneurysm. <i>Clinical Endocrinology</i> , 2010, 72, 128-132. | 2.4 | 34 |
| 22 | Association of statin prescription with small abdominal aortic aneurysm progression. <i>American Heart Journal</i> , 2010, 159, 307-313. | 2.7 | 152 |
| 23 | The Novel Association of the Chemokine CCL22 with Abdominal Aortic Aneurysm. <i>American Journal of Pathology</i> , 2010, 176, 2098-2106. | 3.8 | 39 |
| 24 | A Single-Nucleotide Polymorphism in the Gene Encoding Osteoprotegerin Is Associated With Diastolic Blood Pressure in Older Men. <i>American Journal of Hypertension</i> , 2009, 22, 1167-1170. | 2.0 | 6 |
| 25 | On a mouse monoclonal antibody that neutralizes all four dengue virus serotypes. <i>Journal of General Virology</i> , 2009, 90, 799-809. | 2.9 | 73 |
| 26 | Role of homocysteine in aortic calcification and osteogenic cell differentiation. <i>Atherosclerosis</i> , 2009, 202, 557-566. | 0.8 | 47 |
| 27 | Modulation of endothelial cell thrombomodulin by PPAR ligands α Variation according to environment. <i>Thrombosis Research</i> , 2008, 121, 827-834. | 1.7 | 14 |
| 28 | Relationship between CT anthropometric measurements, adipokines and abdominal aortic calcification. <i>Atherosclerosis</i> , 2008, 197, 428-434. | 0.8 | 45 |
| 29 | Reduced expansion rate of abdominal aortic aneurysms in patients with diabetes may be related to aberrant monocyte-matrix interactions. <i>European Heart Journal</i> , 2008, 29, 665-672. | 2.2 | 160 |
| 30 | Obesity, Adipokines, and Abdominal Aortic Aneurysm. <i>Circulation</i> , 2007, 116, 2275-2279. | 1.6 | 135 |
| 31 | Association Between Osteopontin and Human Abdominal Aortic Aneurysm. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007, 27, 655-660. | 2.4 | 114 |
| 32 | Effects of Peroxisome Proliferator-Activated Receptor Ligands in Modulating Tissue Factor and Tissue Factor Pathway Inhibitor in Acutely Symptomatic Carotid Atheromas. <i>Stroke</i> , 2007, 38, 1501-1508. | 2.0 | 27 |
| 33 | Association of obesity and metabolic syndrome with the severity and outcome of intermittent claudication. <i>Journal of Vascular Surgery</i> , 2007, 45, 40-46. | 1.1 | 80 |
| 34 | The domains carrying the opposing activities in adenylyltransferase are separated by a central regulatory domain. <i>FEBS Journal</i> , 2007, 274, 2865-2877. | 4.7 | 11 |
| 35 | Serum Osteoprotegerin as a Biomarker for Vascular Disease. <i>American Journal of Cardiology</i> , 2007, 100, 561. | 1.6 | 5 |
| 36 | Assessment of a Serum Assay for Quantification of Abdominal Aortic Calcification. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006, 26, 2574-2576. | 2.4 | 51 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Expression, purification, crystallization, and preliminary X-ray analysis of the N-terminal domain of Escherichia coli adenylyl transferase. Protein Expression and Purification, 2004, 34, 142-146. | 1.3 | 8 |
| 38 | The structures of the PII proteins from the cyanobacteria Synechococcus PCC 7942 and Synechocystis PCC 6803. Acta Crystallographica Section D: Biological Crystallography, 2003, 59, 2183-2190. | 2.5 | 47 |
| 39 | The Escherichia coli signal transducers PII (GlnB) and GlnK form heterotrimers in vivo: Fine tuning the nitrogen signal cascade. Proceedings of the National Academy of Sciences of the United States of America, 2000, 97, 3942-3947. | 7.1 | 51 |