

Jun-Sing Wang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2819953/jun-sing-wang-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

64

papers

710

citations

14

h-index

24

g-index

74

ext. papers

1,018

ext. citations

5.6

avg, IF

3.63

L-index

#	Paper	IF	Citations
64	Associations of Adherence to the DASH Diet and the Mediterranean Diet With All-Cause Mortality in Subjects With Various Glucose Regulation States.. <i>Frontiers in Nutrition</i> , 2022 , 9, 828792	6.2	1
63	The Impact of COVID-19 Surges in 2019-2021 on Patient-Reported Outcome Measures After Spine Surgery at an Academic Tertiary Referral Center in Taiwan: A Retrospective Observational Cohort Study.. <i>Frontiers in Surgery</i> , 2022 , 9, 853441	2.3	
62	Factors Associated with Postoperative Lipiduria and Hypoxemia in Patients Undergoing Surgery for Orthopedic Fractures.. <i>Frontiers in Surgery</i> , 2022 , 9, 814229	2.3	
61	The power of genetic diversity in genome-wide association studies of lipids. <i>Nature</i> , 2021 ,	50.4	24
60	Therapeutic Potential of Tpl2 (Tumor Progression Locus 2) Inhibition on Diabetic Vasculopathy Through the Blockage of the Inflammasome Complex. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021 , 41, e46-e62	9.4	
59	Distinct associations of self-monitoring of blood glucose with glycemic control and hypoglycemia between groups of recently diagnosed and long-term follow-up type 2 diabetes: The Taiwan Diabetes Registry. <i>International Journal of Clinical Practice</i> , 2021 , 75, e14410	2.9	1
58	Association of diurnal calorie trajectory with all-cause mortality: Findings from the National Health and Nutrition Examination Survey. <i>Clinical Nutrition</i> , 2021 , 40, 1920-1925	5.9	0
57	Associations of low-carbohydrate and low-fat intakes with all-cause mortality in subjects with prediabetes with and without insulin resistance. <i>Clinical Nutrition</i> , 2021 , 40, 3601-3607	5.9	3
56	Subjects with coronary artery disease and reduced ejection fraction have longer (GT) repeats in the heme-oxygenase 1 gene promoter. <i>Heart and Vessels</i> , 2021 , 36, 615-620	2.1	5
55	HbA1c trajectory and cardiovascular outcomes: an analysis of data from the Action to Control Cardiovascular Risk in Diabetes (ACCORD) study. <i>Therapeutic Advances in Chronic Disease</i> , 2021 , 12, 20406223211026391	4.9	0
54	The dawn phenomenon in type 2 diabetes: its association with glucose excursions and changes after oral glucose-lowering drugs. <i>Therapeutic Advances in Chronic Disease</i> , 2021 , 12, 204062232110336749	4.9	
53	Postchallenge glucose increment was associated with hemoglobin glycation index in subjects with no history of diabetes. <i>Journal of Investigative Medicine</i> , 2021 , 69, 1044-1049	2.9	2
52	Subjects with microvascular angina have longer GT repeats polymorphism in the haem oxygenase-1 gene promoter. <i>Biomarkers</i> , 2020 , 25, 144-148	2.6	1
51	Hypoglycaemic episodes increase the risk of ventricular arrhythmia and sudden cardiac arrest in patients with type 2 diabetes-A nationwide cohort study. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36, e3226	7.5	5
50	Systolic blood pressure trajectory and cardiovascular outcomes: An analysis using data in the Systolic Blood Pressure Intervention Trial. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13450	2.9	8
49	Associations of fear of hypoglycemia with second-line use of insulin secretagogues or insulin and subsequent glycemic control in patients with type 2 diabetes: An analysis using data from the DISCOVER study. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13485	2.9	2
48	Effect of low-protein intake on all-cause mortality in subjects with an estimated glomerular filtration rate higher than 60 mL/min/1.73 m ² with or without albuminuria. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13505	2.9	0

47	Association Between Serum Adipsin Levels and Insulin Resistance in Subjects With Various Degrees of Glucose Intolerance. <i>Journal of the Endocrine Society</i> , 2019 , 3, 403-410	0.4	14
46	Multi-ancestry genome-wide gene-smoking interaction study of 387,272 individuals identifies new loci associated with serum lipids. <i>Nature Genetics</i> , 2019 , 51, 636-648	36.3	59
45	Hospitalization in patients with type 2 diabetes mellitus in Taiwan: A nationwide population-based observational study. <i>Journal of the Formosan Medical Association</i> , 2019 , 118 Suppl 2, S90-S95	3.2	6
44	Effects of hyperuricemia on incident renal replacement therapy and all-cause mortality among patients with chronic kidney disease stages 3-5: a retrospective cohort study. <i>Sao Paulo Medical Journal</i> , 2019 , 137, 523-529	1.6	3
43	Effects of statins on all-cause mortality at different low-density-lipoprotein cholesterol levels in Asian patients with type 2 diabetes. <i>Current Medical Research and Opinion</i> , 2018 , 34, 1885-1892	2.5	5
42	Difference between observed and predicted glycated hemoglobin at baseline and treatment response to vildagliptin-based dual oral therapy in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018 , 138, 119-127	7.4	7
41	Levels of serum high mobility group box 1 were independently associated with cardiovascular risk in patients undergoing coronary angiography. <i>Clinica Chimica Acta</i> , 2018 , 483, 130-134	6.2	4
40	Testing for HbA1c, in addition to the oral glucose tolerance test, in screening for abnormal glucose regulation helps to reveal patients with early β cell function impairment. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 56, 1345-1352	5.9	1
39	Trajectories of fasting plasma glucose variability and mortality in type 2 diabetes. <i>Diabetes and Metabolism</i> , 2018 , 44, 121-128	5.4	24
38	Two-hour post-challenge hyperglycemia, but not fasting plasma glucose, associated with severity of coronary artery disease in patients with angina. <i>PLoS ONE</i> , 2018 , 13, e0202280	3.7	3
37	The association between brain-derived neurotrophic factor and central pulse pressure after an oral glucose tolerance test. <i>Clinica Chimica Acta</i> , 2018 , 476, 1-8	6.2	3
36	Regaining body weight after weight reduction further increases pulse wave velocity in obese men with metabolic syndrome. <i>Medicine (United States)</i> , 2018 , 97, e12730	1.8	2
35	Negative association between serum aryl hydrocarbon receptor concentrations and β cell function in patients with no history of diabetes undergoing coronary angiography. <i>Journal of Diabetes</i> , 2018 , 10, 958-964	3.8	2
34	Glycemic excursions are positively associated with HbA1c reduction from baseline after treatment with acarbose in patients with type 2 diabetes on metformin monotherapy. <i>Journal of Diabetes</i> , 2017 , 9, 248-255	3.8	3
33	The synergistic effect of vascular cell adhesion molecule-1 and coronary artery disease on brain-derived neurotrophic factor. <i>Clinica Chimica Acta</i> , 2017 , 466, 194-200	6.2	9
32	Protein-Truncating Variants at the Cholesteryl Ester Transfer Protein Gene and Risk for Coronary Heart Disease. <i>Circulation Research</i> , 2017 , 121, 81-88	15.7	48
31	The synergistic effect of inflammation and metabolic syndrome on intraocular pressure: A cross-sectional study. <i>Medicine (United States)</i> , 2017 , 96, e7851	1.8	6
30	Associations of fibroblast growth factor 21 with cardiovascular risk and β cell function in patients who had no history of diabetes. <i>Clinica Chimica Acta</i> , 2017 , 472, 80-85	6.2	3

29	Differential expression of circulating vascular cell adhesion molecule-1 in subjects with coronary artery disease and cardiac syndrome X without known diabetes mellitus. <i>Biomarkers</i> , 2017 , 22, 798-804	2.6	5
28	Post-challenge insulin concentration is useful for differentiating between coronary artery disease and cardiac syndrome X in subjects without known diabetes mellitus. <i>Diabetology and Metabolic Syndrome</i> , 2017 , 9, 10	5.6	5
27	Hemoglobin glycation index as a useful predictor of therapeutic responses to dipeptidyl peptidase-4 inhibitors in patients with type 2 diabetes. <i>PLoS ONE</i> , 2017 , 12, e0171753	3.7	10
26	Relationship between body weight and the increment in serum brain-derived neurotrophic factor after oral glucose challenge in men with obesity and metabolic syndrome: A prospective study. <i>Medicine (United States)</i> , 2016 , 95, e5260	1.8	15
25	Statin treatment is associated with a negative correlation between NT-proBNP and insulin resistance in patients without history of heart failure. <i>Clinica Chimica Acta</i> , 2016 , 459, 84-88	6.2	1
24	Glycemic excursions are positively associated with changes in duration of asymptomatic hypoglycemia after treatment intensification in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2016 , 113, 108-15	7.4	13
23	Association of MMP-9 to TIMP-1 ratio with long-term body weight and waist circumference after dietary weight reduction in men with metabolic syndrome. <i>Clinica Chimica Acta</i> , 2016 , 452, 182-4	6.2	
22	Epigenetic regulation of HGK/MAP4K4 in T cells of type 2 diabetes patients. <i>Oncotarget</i> , 2016 , 7, 10976-89	3.9	14
21	Physical activity and albuminuria were associated with painful diabetic polyneuropathy in type 2 diabetes in an ethnic Chinese population. <i>Clinica Chimica Acta</i> , 2016 , 462, 55-59	6.2	3
20	A new formula for estimation of low-density lipoprotein cholesterol in an ethnic Chinese population. <i>Clinical Chemistry and Laboratory Medicine</i> , 2015 , 53, 1871-9	5.9	1
19	Hydroxychloroquine reduces risk of incident diabetes mellitus in lupus patients in a dose-dependent manner: a population-based cohort study. <i>Rheumatology</i> , 2015 , 54, 1244-9	3.9	77
18	Coronary severity score and C-reactive protein predict major adverse cardiovascular events in patients with stable coronary artery disease (from the Taichung CAD study). <i>Clinica Chimica Acta</i> , 2015 , 445, 93-100	6.2	17
17	Comparing HbA1c, fasting and 2-h plasma glucose for screening for abnormal glucose regulation in patients undergoing coronary angiography. <i>Clinical Chemistry and Laboratory Medicine</i> , 2015 , 53, 1441-9	5.9	10
16	Factors associated with fibroblast growth factor 19 increment after oral glucose loading in patients who were previously admitted for coronary angiography. <i>Clinica Chimica Acta</i> , 2015 , 450, 237-42	6.2	3
15	Circulating adipokines and insulin resistance in subjects with combined cardiac and metabolic syndrome X. <i>Diabetology and Metabolic Syndrome</i> , 2015 , 7, 83	5.6	7
14	Weight loss reduces serum monocyte chemoattractant protein-1 concentrations in association with improvements in renal injury in obese men with metabolic syndrome. <i>Clinical Chemistry and Laboratory Medicine</i> , 2015 , 53, 623-9	5.9	6
13	Value of Chromosome 9p21 Polymorphism for Prediction of Cardiovascular Mortality in Han Chinese Without Coronary Lesions: An Observational Study. <i>Medicine (United States)</i> , 2015 , 94, e1538	1.8	3
12	Post-meal β -cell function predicts the efficacy of glycemic control in patients with type 2 diabetes inadequately controlled by metformin monotherapy after addition of glibenclamide or acarbose. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 68	5.6	6

11	Insulinoma in a young female patient with systemic lupus erythematosus: a case report. <i>Endocrine Practice</i> , 2014 , 20, e256-9	3.2	1
10	Correlation between reduction of superior interventricular groove epicardial fat thickness and improvement of insulin resistance after weight loss in obese men. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 115	5.6	6
9	Acarbose plus metformin fixed-dose combination outperforms acarbose monotherapy for type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2013 , 102, 16-24	7.4	20
8	Leptin to adiponectin ratio as a useful predictor for cardiac syndrome X. <i>Biomarkers</i> , 2013 , 18, 44-50	2.6	17
7	Trans-ethnic fine mapping identifies a novel independent locus at the 3' end of CDKAL1 and novel variants of several susceptibility loci for type 2 diabetes in a Han Chinese population. <i>Diabetologia</i> , 2013 , 56, 2619-28	10.3	25
6	Performance of HbA1c and fasting plasma glucose in screening for diabetes in patients undergoing coronary angiography. <i>Diabetes Care</i> , 2013 , 36, 1138-40	14.6	14
5	Genome-wide association study in a Chinese population with diabetic retinopathy. <i>Human Molecular Genetics</i> , 2013 , 22, 3165-73	5.6	77
4	Dynamic and dual effects of glycated hemoglobin on estimated glomerular filtration rate in type 2 diabetic outpatients. <i>American Journal of Nephrology</i> , 2013 , 38, 19-26	4.6	12
3	The beneficial effect of α-glucosidase inhibitor on glucose variability compared with sulfonylurea in Taiwanese type 2 diabetic patients inadequately controlled with metformin: preliminary data. <i>Journal of Diabetes and Its Complications</i> , 2011 , 25, 332-8	3.2	26
2	Effects of acarbose versus glibenclamide on glycemic excursion and oxidative stress in type 2 diabetic patients inadequately controlled by metformin: a 24-week, randomized, open-label, parallel-group comparison. <i>Clinical Therapeutics</i> , 2011 , 33, 1932-42	3.5	52
1	Effects of patient volume on quality of outpatient diabetes care. <i>Diabetes Research and Clinical Practice</i> , 2009 , 84, e27-9	7.4	9