

Bryan B Brimhall

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2821148/publications.pdf>

Version: 2024-02-01

10
papers

212
citations

1937457

4
h-index

1719901

7
g-index

10
all docs

10
docs citations

10
times ranked

403
citing authors

#	ARTICLE	IF	CITATIONS
1	Increased Incidence of Pseudoaneurysm Bleeding With Lumen-Apposing Metal Stents Compared to Double-Pigtail Plastic Stents in Patients With Peripancreatic Fluid Collections. <i>Clinical Gastroenterology and Hepatology</i> , 2018, 16, 1521-1528.	2.4	115
2	Medical Marijuana for Digestive Disorders: High Time to Prescribe?. <i>American Journal of Gastroenterology</i> , 2015, 110, 208-214.	0.2	47
3	Enteral Stents for Malignant Gastric Outlet Obstruction. <i>Gastrointestinal Endoscopy Clinics of North America</i> , 2011, 21, 389-403.	0.6	38
4	Current Endoscopic Approaches for the Treatment of Barrett Esophagus. <i>Journal of Clinical Gastroenterology</i> , 2017, 51, 2-11.	1.1	8
5	A low-volume polyethylene glycol solution was associated with an increased suboptimal bowel preparation rate but had similar recommendations for an early repeat colonoscopy, procedure times, and adenoma detection rates. <i>PLoS ONE</i> , 2017, 12, e0176265.	1.1	2
6	Esophageal Stents for the Treatment of Malignant Dysphagia in Patients with Esophageal Cancer. <i>Hospital Practice (1995)</i> , 2010, 38, 94-102.	0.5	1
7	Medicare Under Age 65 and Medicaid Patients Have Poorer Bowel Preparations: Implications for Recommendations for an Early Repeat Colonoscopy. <i>PLoS ONE</i> , 2016, 11, e0155208.	1.1	1
8	Mo1081 Effect of a Low-Volume Compared to a High-Volume Polyethylene Glycol Bowel Preparation Solution on Prep Quality, Procedure Times, and Adenoma Detection Rates. <i>Gastroenterology</i> , 2014, 146, S-552.	0.6	0
9	123 Medicare Patients Under the Age of 65 Years and Medicaid Patients Are More Likely to Have a Suboptimal Bowel Preparation for Colonoscopy. <i>Gastroenterology</i> , 2014, 146, S-34-S-35.	0.6	0
10	Simple, Low-Cost Educational Interventions Can Reduce Radiation Exposure. <i>Clinical Gastroenterology and Hepatology</i> , 2018, 16, 488-490.	2.4	0