

# Ge Yu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2821666/publications.pdf>

Version: 2024-02-01

28  
papers

652  
citations

623734

14  
h-index

580821

25  
g-index

29  
all docs

29  
docs citations

29  
times ranked

1124  
citing authors

#	ARTICLE	IF	CITATIONS
1	Mental health services in response to the COVID-19 pandemic in high-income countries: protocol for a rapid review. <i>BMJ Open</i> , 2022, 12, e062078.	1.9	4
2	A predictive algorithm using clinical and laboratory parameters may assist in ruling out and in diagnosing MDS. <i>Blood Advances</i> , 2021, 5, 3066-3075.	5.2	12
3	The anemia-independent impact of myelodysplastic syndromes on health-related quality of life. <i>Annals of Hematology</i> , 2021, 100, 2921-2932.	1.8	7
4	Impact of treatment with iron chelation therapy in patients with lower-risk myelodysplastic syndromes participating in the European MDS registry. <i>Haematologica</i> , 2020, 105, 640-651.	3.5	32
5	Novel dynamic outcome indicators and clinical endpoints in myelodysplastic syndrome; the European LeukemiaNet MDS Registry and MDS-RIGHT project perspective. <i>Haematologica</i> , 2020, 105, 2516-2523.	3.5	12
6	The association between patient-professional partnerships and self-management of chronic back pain: A mixed methods study. <i>European Journal of Pain</i> , 2018, 22, 1229-1244.	2.8	11
7	Health-related quality of life in lower-risk MDS patients compared with age- and sex-matched reference populations: a European LeukemiaNet study. <i>Leukemia</i> , 2018, 32, 1380-1392.	7.2	66
8	MDS Diagnosis: Many Patients May Not Require Bone Marrow Examination. <i>Blood</i> , 2018, 132, 4357-4357.	1.4	1
9	Excess Mortality in Low-Risk MDS Can be Explained By MDS and AML Related Causes of Death. <i>Blood</i> , 2018, 132, 4385-4385.	1.4	1
10	UK Utility Elicitation in Patients with Follicular Lymphoma. <i>Value in Health</i> , 2017, 20, A449.	0.3	2
11	The effects of patient-professional partnerships on the self-management and health outcomes for patients with chronic back pain: A quasi-experimental study. <i>International Journal of Nursing Studies</i> , 2016, 59, 197-207.	5.6	12
12	Impact of Treatment with Iron Chelators in Lower-Risk MDS Patients Participating in the European LeukemiaNet MDS (EUMDS) Registry. <i>Blood</i> , 2016, 128, 3186-3186.	1.4	14
13	Cost effectiveness of treatment with percutaneous Kirschner wires versus volar locking plate for adult patients with a dorsally displaced fracture of the distal radius. <i>Bone and Joint Journal</i> , 2015, 97-B, 1082-1089.	4.4	61
14	A multilevel cross-lagged structural equation analysis for reciprocal relationship between social capital and health. <i>Social Science and Medicine</i> , 2015, 142, 1-8.	3.8	33
15	UK DRAFFT: a randomised controlled trial of percutaneous fixation with Kirschner wires versus volar locking-plate fixation in the treatment of adult patients with a dorsally displaced fracture of the distal radius. <i>Health Technology Assessment</i> , 2015, 19, 1-124.	2.8	72
16	Measures of exposure to the Well London Phase-1 intervention and their association with health well-being and social outcomes. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 597-605.	3.7	19
17	Well London Phase-1: results among adults of a cluster-randomised trial of a community engagement approach to improving health behaviours and mental well-being in deprived inner-city neighbourhoods. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 606-614.	3.7	43
18	Cost-effectiveness of silicone and alginate impressions for complete dentures. <i>Journal of Dentistry</i> , 2014, 42, 902-907.	4.1	23

#	ARTICLE	IF	CITATIONS
19	Predictors of the timing of initiation of antenatal care in an ethnically diverse urban cohort in the UK. BMC Pregnancy and Childbirth, 2013, 13, 103.	2.4	53
20	Physical Activity in Deprived Communities in London: Examining Individual and Neighbourhood-Level Factors. PLoS ONE, 2013, 8, e69472.	2.5	13
21	Types of Social Capital and Mental Disorder in Deprived Urban Areas: A Multilevel Study of 40 Disadvantaged London Neighbourhoods. PLoS ONE, 2013, 8, e80127.	2.5	25
22	Well London: Results of a Cluster-Randomised Trial of a Community Development Approach to Improving Health Behaviours and Mental Wellbeing in Deprived Inner-City Neighbourhoods. Journal of Epidemiology and Community Health, 2012, 66, A3.1-A3.	3.7	2
23	Who stays, who drops out? Biosocial predictors of longer-term adherence in participants attending an exercise referral scheme in the UK. BMC Public Health, 2012, 12, 347.	2.9	36
24	The Well London program - a cluster randomized trial of community engagement for improving health behaviors and mental wellbeing: baseline survey results. Trials, 2012, 13, 105.	1.6	17
25	Think of your art-eries: Arts participation, behavioural cardiovascular risk factors and mental well-being in deprived communities in London. Public Health, 2012, 126, S57-S64.	2.9	35
26	A multilevel analysis of the association between social networks and support on leisure time physical activity: Evidence from 40 disadvantaged areas in London. Health and Place, 2011, 17, 1023-1029.	3.3	38
27	Prevalence of Low Physical Activity and its Relation to Social Environment in Deprived Areas in the London Borough of Redbridge. Social Indicators Research, 2011, 104, 311-322.	2.7	7
28	Determinants of physical activity in deprived communities in London: Examining the effects of individual and neighbourhood characteristics. Journal of Epidemiology and Community Health, 2011, 65, A37-A37.	3.7	1