

Kurt Hjlund

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2821740/kurt-hojlund-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

124
papers

8,034
citations

44
h-index

89
g-index

134
ext. papers

9,279
ext. citations

6.5
avg, IF

5.55
L-index

#	Paper	IF	Citations
124	Factors Mediating Exercise-induced Organ Crosstalk.. <i>Acta Physiologica</i> , 2022 , e13766	5.6	2
123	The effect of empagliflozin on growth differentiation factor 15 in patients with heart failure: a randomized controlled trial (Empire HF Biomarker).. <i>Cardiovascular Diabetology</i> , 2022 , 21, 34	8.7	0
122	Impact of Lean Body Mass and Insulin Sensitivity on the IGF-1-Bone Mass Axis in Adolescence: the EPICOM Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e772-e781	5.6	2
121	GAD65 autoantibodies and glucose tolerance in offspring born to women with and without type 1 diabetes (The EPICOM study). <i>Endocrinology, Diabetes and Metabolism</i> , 2021 , 5, e00310	2.7	1
120	Personalized phosphoproteomics identifies functional signaling. <i>Nature Biotechnology</i> , 2021 ,	44.5	8
119	Life-course Exposure to Perfluoroalkyl Substances in Relation to Markers of Glucose Homeostasis in Early Adulthood. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, 2495-2504	5.6	5
118	Gamma-glutamyltransferase, arterial remodeling and prehypertension in a healthy population at low cardiometabolic risk. <i>Journal of Human Hypertension</i> , 2021 , 35, 334-342	2.6	
117	Acute effects of delayed-release hydrolyzed pine nut oil on glucose tolerance, incretins, ghrelin and appetite in healthy humans. <i>Clinical Nutrition</i> , 2021 , 40, 2169-2179	5.9	2
116	Serum sclerostin and glucose homeostasis: No association in healthy men. Cross-sectional and prospective data from the EGIR-RISC study. <i>Bone</i> , 2021 , 143, 115681	4.7	0
115	Oral and intravenous pharmacokinetics of metformin with and without oral codeine intake in healthy subjects: A cross-over study. <i>Clinical and Translational Science</i> , 2021 , 14, 2408-2419	4.9	2
114	Effect of Exercise Training on Fat Loss-Energetic Perspectives and the Role of Improved Adipose Tissue Function and Body Fat Distribution. <i>Frontiers in Physiology</i> , 2021 , 12, 737709	4.6	5
113	Apolipoprotein D and transthyretin are reduced in female adolescent offspring of women with type 1 diabetes: The EPICOM study.. <i>Diabetic Medicine</i> , 2021 , e14776	3.5	1
112	A novel gene in early childhood diabetes: EDEM2 silencing decreases SLC2A2 and PXD1 expression, leading to impaired insulin secretion. <i>Molecular Genetics and Genomics</i> , 2020 , 295, 1253-1262	3.1	1
111	Circulating Follistatin and Activin A and Their Regulation by Insulin in Obesity and Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	10
110	Disruption of fasting and post-load glucose homeostasis are largely independent and sustained by distinct and early major beta-cell function defects: a cross-sectional and longitudinal analysis of the Relationship between Insulin Sensitivity and Cardiovascular risk (RISC) study cohort. <i>Metabolism: Clinical and Experimental</i> , 2020 , 105, 154185	12.7	5
109	Effect of long-term testosterone therapy on molecular regulators of skeletal muscle mass and fibre-type distribution in aging men with subnormal testosterone. <i>Metabolism: Clinical and Experimental</i> , 2020 , 112, 154347	12.7	2
108	The Mitochondrial Proteomic Signatures of Human Skeletal Muscle Linked to Insulin Resistance. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	3

107	Exercise Induction of Key Transcriptional Regulators of Metabolic Adaptation in Muscle Is Preserved in Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 4909-4920	5.6	6
106	Fatty Liver Among Adolescent Offspring of Women With Type 1 Diabetes (the EPICOM Study). <i>Diabetes Care</i> , 2019 , 42, 1560-1568	14.6	4
105	Obesity-Associated Hypermetabolism and Accelerated Senescence of Bone Marrow Stromal Stem Cells Suggest a Potential Mechanism for Bone Fragility. <i>Cell Reports</i> , 2019 , 27, 2050-2062.e6	10.6	41
104	In human nephrectomy specimens, the kidney level of tubular transport proteins does not correlate with their abundance in urinary extracellular vesicles. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 317, F560-F571	4.3	12
103	Acute Exercise Increases Plasma Levels of Muscle-Derived Microvesicles Carrying Fatty Acid Transport Proteins. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 4804-4814	5.6	7
102	Differential methylation of the type 2 diabetes susceptibility locus KCNQ1 is associated with insulin sensitivity and is predicted by CpG site specific genetic variation. <i>Diabetes Research and Clinical Practice</i> , 2019 , 148, 189-199	7.4	8
101	Lower mortality and cardiovascular event rates in patients with Latent Autoimmune Diabetes In Adults (LADA) as compared with type 2 diabetes and insulin deficient diabetes: A cohort study of 4368 patients. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 107-113	7.4	12
100	Adrenal activity and metabolic risk during randomized escitalopram or placebo treatment in PCOS. <i>Endocrine Connections</i> , 2018 , 7, 479-489	3.5	5
99	The relationship between bone turnover and insulin sensitivity and secretion: Cross-sectional and prospective data from the RISC cohort study. <i>Bone</i> , 2018 , 108, 98-105	4.7	7
98	Skeletal muscle O-GlcNAc transferase is important for muscle energy homeostasis and whole-body insulin sensitivity. <i>Molecular Metabolism</i> , 2018 , 11, 160-177	8.8	44
97	Easy, Fast, and Reproducible Quantification of Cholesterol and Other Lipids in Human Plasma by Combined High Resolution MSX and FTMS Analysis. <i>Journal of the American Society for Mass Spectrometry</i> , 2018 , 29, 34-41	3.5	15
96	Differential effects of age and sex on insulin sensitivity and body composition in adolescent offspring of women with type 1 diabetes: results from the EPICOM study. <i>Diabetologia</i> , 2018 , 61, 210-215	10.3	11
95	Renal function markers and insulin sensitivity after 3 years in a healthy cohort, the EGIR-RISC study. <i>BMC Nephrology</i> , 2018 , 19, 124	2.7	
94	Intact regulation of muscle expression and circulating levels of myokines in response to exercise in patients with type 2 diabetes. <i>Physiological Reports</i> , 2018 , 6, e13723	2.6	18
93	First Genome-Wide Association Study of Latent Autoimmune Diabetes in Adults Reveals Novel Insights Linking Immune and Metabolic Diabetes. <i>Diabetes Care</i> , 2018 , 41, 2396-2403	14.6	57
92	Proteomic study of skeletal muscle in obesity and type 2 diabetes: progress and potential. <i>Expert Review of Proteomics</i> , 2018 , 15, 817-828	4.2	3
91	Transcriptional and Epigenetic Changes Influencing Skeletal Muscle Metabolism in Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 4465-4477	5.6	35
90	The heterozygous N291S mutation in the lipoprotein lipase gene impairs whole-body insulin sensitivity and affects a distinct set of plasma metabolites in humans. <i>Journal of Clinical Lipidology</i> , 2017 , 11, 515-523.e6	4.9	

89	Autonomic nervous system activation mediates the increase in whole-body glucose uptake in response to electroacupuncture. <i>FASEB Journal</i> , 2017 , 31, 3288-3297	0.9	19
88	Abnormal levels of adipokines in adolescent offspring of women with type 1 diabetes - Results from the EPICOM study. <i>Metabolism: Clinical and Experimental</i> , 2017 , 72, 47-56	12.7	6
87	Characterization of the CLASP2 Protein Interaction Network Identifies SOGA1 as a Microtubule-Associated Protein. <i>Molecular and Cellular Proteomics</i> , 2017 , 16, 1718-1735	7.6	24
86	The Human Skeletal Muscle Proteome Project: a reappraisal of the current literature. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2017 , 8, 5-18	10.3	51
85	Intact initiation of autophagy and mitochondrial fission by acute exercise in skeletal muscle of patients with Type 2 diabetes. <i>Clinical Science</i> , 2017 , 131, 37-47	6.5	22
84	Metabolic risk profiles in diabetes stratified according to age at onset, islet autoimmunity and fasting C-peptide. <i>Diabetes Research and Clinical Practice</i> , 2017 , 134, 62-71	7.4	9
83	Effects of insulin and exercise training on FGF21, its receptors and target genes in obesity and type 2 diabetes. <i>Diabetologia</i> , 2017 , 60, 2042-2051	10.3	38
82	Mitochondrial phosphoproteomics of mammalian tissues. <i>Mitochondrion</i> , 2017 , 33, 45-57	4.9	22
81	Microvesicles Correlated with Components of Metabolic Syndrome in Men with Type 2 Diabetes Mellitus and Lowered Testosterone Levels But Were Unaltered by Testosterone Therapy. <i>Journal of Diabetes Research</i> , 2017 , 2017, 4257875	3.9	7
80	Beta-cell function is associated with carotid intima-media thickness independently of insulin resistance in healthy individuals. <i>Journal of Hypertension</i> , 2016 , 34, 685-91	1.9	8
79	Intact Regulation of the AMPK Signaling Network in Response to Exercise and Insulin in Skeletal Muscle of Male Patients With Type 2 Diabetes: Illumination of AMPK Activation in Recovery From Exercise. <i>Diabetes</i> , 2016 , 65, 1219-30	0.9	47
78	Follow-up duration influences the relative importance of OGTT and optimal timing of glucose measurements for predicting future type 2 diabetes. <i>European Journal of Endocrinology</i> , 2016 , 174, 591-600	6.5	8
77	Modification and Validation of the Triglyceride-to-HDL Cholesterol Ratio as a Surrogate of Insulin Sensitivity in White Juveniles and Adults without Diabetes Mellitus: The Single Point Insulin Sensitivity Estimator (SPISE). <i>Clinical Chemistry</i> , 2016 , 62, 1211-9	5.5	32
76	Effect of testosterone on insulin sensitivity, oxidative metabolism and body composition in aging men with type 2 diabetes on metformin monotherapy. <i>Diabetes, Obesity and Metabolism</i> , 2016 , 18, 980-9	6.7	37
75	Impaired Glucose Tolerance in Healthy Men Treated with St. John's Wort. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2016 , 118, 219-24	3.1	12
74	Intake of St John's wort improves the glucose tolerance in healthy subjects who ingest metformin compared with metformin alone. <i>British Journal of Clinical Pharmacology</i> , 2015 , 79, 298-306	3.8	24
73	Response to Comment on Pilz et al. Insulin sensitivity and albuminuria: the RISC study. <i>Diabetes Care</i> 2014;37:1597-1603. <i>Diabetes Care</i> , 2015 , 38, e31	14.6	
72	Dysregulation of muscle glycogen synthase in recovery from exercise in type 2 diabetes. <i>Diabetologia</i> , 2015 , 58, 1569-78	10.3	20

71	Increased metabolic risk in adolescent offspring of mothers with type 1 diabetes: the EPICOM study. <i>Diabetologia</i> , 2015 , 58, 1454-63	10.3	30
70	Endogenous glucose production increases in response to metformin treatment in the glycogen-depleted state in humans: a randomised trial. <i>Diabetologia</i> , 2015 , 58, 2494-502	10.3	22
69	Glucose tolerance is associated with differential expression of microRNAs in skeletal muscle: results from studies of twins with and without type 2 diabetes. <i>Diabetologia</i> , 2015 , 58, 363-73	10.3	43
68	Human muscle fiber type-specific insulin signaling: impact of obesity and type 2 diabetes. <i>Diabetes</i> , 2015 , 64, 485-97	0.9	105
67	Steady-state pharmacokinetics of metformin is independent of the OCT1 genotype in healthy volunteers. <i>European Journal of Clinical Pharmacology</i> , 2015 , 71, 691-697	2.8	36
66	Markers of autophagy are adapted to hyperglycaemia in skeletal muscle in type 2 diabetes. <i>Diabetologia</i> , 2015 , 58, 2087-95	10.3	45
65	Differential effects of strength training and testosterone treatment on soluble CD36 in aging men: Possible relation to changes in body composition. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015 , 75, 659-66	2	6
64	A PGC-1 β and muscle fibre type-related decrease in markers of mitochondrial oxidative metabolism in skeletal muscle of humans with inherited insulin resistance. <i>Diabetologia</i> , 2014 , 57, 1006-15	10.3	25
63	Insulin increases phosphorylation of mitochondrial proteins in human skeletal muscle in vivo. <i>Journal of Proteome Research</i> , 2014 , 13, 2359-69	5.6	20
62	Increased interaction with insulin receptor substrate 1, a novel abnormality in insulin resistance and type 2 diabetes. <i>Diabetes</i> , 2014 , 63, 1933-47	0.9	39
61	Effect of testosterone on markers of mitochondrial oxidative phosphorylation and lipid metabolism in muscle of aging men with subnormal bioavailable testosterone. <i>European Journal of Endocrinology</i> , 2014 , 171, 77-88	6.5	23
60	Insulin sensitivity and albuminuria: the RISC study. <i>Diabetes Care</i> , 2014 , 37, 1597-603	14.6	36
59	Metabolism and insulin signaling in common metabolic disorders and inherited insulin resistance. <i>Danish Medical Journal</i> , 2014 , 61, B4890	3.8	62
58	Studies of association of AGPAT6 variants with type 2 diabetes and related metabolic phenotypes in 12,068 Danes. <i>BMC Medical Genetics</i> , 2013 , 14, 113	2.1	2
57	Tissue specific phosphorylation of mitochondrial proteins isolated from rat liver, heart muscle, and skeletal muscle. <i>Journal of Proteome Research</i> , 2013 , 12, 4327-39	5.6	17
56	Rac1 signaling is required for insulin-stimulated glucose uptake and is dysregulated in insulin-resistant murine and human skeletal muscle. <i>Diabetes</i> , 2013 , 62, 1865-75	0.9	128
55	Euglycemic clamp insulin sensitivity and longitudinal systolic blood pressure: role of sex. <i>Hypertension</i> , 2013 , 62, 404-9	8.5	10
54	Influence of apolipoproteins on the association between lipids and insulin sensitivity: a cross-sectional analysis of the RISC Study. <i>Diabetes Care</i> , 2013 , 36, 4125-31	14.6	12

53	Hypoglycemia-associated electroencephalogram and electrocardiogram changes appear simultaneously. <i>Journal of Diabetes Science and Technology</i> , 2013 , 7, 93-9	4.1	10
52	Testosterone therapy increased muscle mass and lipid oxidation in aging men. <i>Age</i> , 2012 , 34, 145-56		51
51	Characterisation of adiponectin multimers and the IGF axis in humans with a heterozygote mutation in the tyrosine kinase domain of the insulin receptor gene. <i>European Journal of Endocrinology</i> , 2012 , 166, 511-9	6.5	5
50	Novel tyrosine phosphorylation sites in rat skeletal muscle revealed by phosphopeptide enrichment and HPLC-ESI-MS/MS. <i>Journal of Proteomics</i> , 2012 , 75, 4017-26	3.9	6
49	Moderate alcohol consumption is associated with improved insulin sensitivity, reduced basal insulin secretion rate and lower fasting glucagon concentration in healthy women. <i>Diabetologia</i> , 2012 , 55, 3228-37	10.3	55
48	A PGC1- β -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. <i>Nature</i> , 2012 , 481, 463-8	50.4	2762
47	Acute hyperinsulinemia is followed by increased serum concentrations of fibroblast growth factor 23 in type 2 diabetes patients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2012 , 72, 108-13	2	14
46	Genome-wide analysis of DNA methylation differences in muscle and fat from monozygotic twins discordant for type 2 diabetes. <i>PLoS ONE</i> , 2012 , 7, e51302	3.7	148
45	Boström et al. reply. <i>Nature</i> , 2012 , 488, E10-E11	50.4	13
44	Plasma FGF21 displays a circadian rhythm during a 72-h fast in healthy female volunteers. <i>Clinical Endocrinology</i> , 2011 , 75, 514-9	3.4	57
43	Phosphoproteome analysis of functional mitochondria isolated from resting human muscle reveals extensive phosphorylation of inner membrane protein complexes and enzymes. <i>Molecular and Cellular Proteomics</i> , 2011 , 10, M110.000299	7.6	120
42	The SNARE protein SNAP23 and the SNARE-interacting protein Munc18c in human skeletal muscle are implicated in insulin resistance/type 2 diabetes. <i>Diabetes</i> , 2010 , 59, 1870-8	0.9	31
41	Increased reactive oxygen species production and lower abundance of complex I subunits and carnitine palmitoyltransferase 1B protein despite normal mitochondrial respiration in insulin-resistant human skeletal muscle. <i>Diabetes</i> , 2010 , 59, 2444-52	0.9	131
40	Proteomics analysis of human skeletal muscle reveals novel abnormalities in obesity and type 2 diabetes. <i>Diabetes</i> , 2010 , 59, 33-42	0.9	185
39	Increased subsarcolemmal lipids in type 2 diabetes: effect of training on localization of lipids, mitochondria, and glycogen in sedentary human skeletal muscle. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2010 , 298, E706-13	6	116
38	Circulating levels of insulin-like growth factor-II/mannose-6-phosphate receptor in obesity and type 2 diabetes. <i>Growth Hormone and IGF Research</i> , 2010 , 20, 185-91	2	24
37	Automated detection of hypoglycemia-induced EEG changes recorded by subcutaneous electrodes in subjects with type 1 diabetes--the brain as a biosensor. <i>Diabetes Research and Clinical Practice</i> , 2010 , 88, 22-8	7.4	46
36	Dysregulation of glycogen synthase COOH- and NH ₂ -terminal phosphorylation by insulin in obesity and type 2 diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 4547-56	5.6	52

35	Acute hyperinsulinemia decreases plasma osteoprotegerin with diminished effect in type 2 diabetes and obesity. <i>European Journal of Endocrinology</i> , 2009 , 161, 95-101	6.5	28
34	Fasting insulin has a stronger association with an adverse cardiometabolic risk profile than insulin resistance: the RISC study. <i>European Journal of Endocrinology</i> , 2009 , 161, 223-30	6.5	16
33	Fatty liver is associated with insulin resistance, risk of coronary heart disease, and early atherosclerosis in a large European population. <i>Hepatology</i> , 2009 , 49, 1537-44	11.2	248
32	Lipid droplets interact with mitochondria using SNAP23. <i>Cell Biology International</i> , 2009 , 33, 934-40	4.5	82
31	Circulating soluble CD36 is a novel marker of liver injury in subjects with altered glucose tolerance. <i>Journal of Nutritional Biochemistry</i> , 2009 , 20, 477-84	6.3	23
30	Proteome profile of functional mitochondria from human skeletal muscle using one-dimensional gel electrophoresis and HPLC-ESI-MS/MS. <i>Journal of Proteomics</i> , 2009 , 72, 1046-60	3.9	59
29	In vivo phosphoproteome of human skeletal muscle revealed by phosphopeptide enrichment and HPLC-ESI-MS/MS. <i>Journal of Proteome Research</i> , 2009 , 8, 4954-65	5.6	71
28	The assembly of lipid droplets and its relation to cellular insulin sensitivity. <i>Biochemical Society Transactions</i> , 2009 , 37, 981-5	5.1	21
27	Lipid droplets and their role in the development of insulin resistance and diabetic dyslipidemia. <i>Clinical Lipidology</i> , 2009 , 4, 611-622		4
26	Mitochondrial dysfunction in type 2 diabetes and obesity. <i>Endocrinology and Metabolism Clinics of North America</i> , 2008 , 37, 713-31, x	5.5	96
25	Characterization of the human skeletal muscle proteome by one-dimensional gel electrophoresis and HPLC-ESI-MS/MS. <i>Molecular and Cellular Proteomics</i> , 2008 , 7, 257-67	7.6	88
24	Impaired insulin-stimulated phosphorylation of Akt and AS160 in skeletal muscle of women with polycystic ovary syndrome is reversed by pioglitazone treatment. <i>Diabetes</i> , 2008 , 57, 357-66	0.9	103
23	Soluble CD36 and risk markers of insulin resistance and atherosclerosis are elevated in polycystic ovary syndrome and significantly reduced during pioglitazone treatment. <i>Diabetes Care</i> , 2008 , 31, 328-34	14.6	78
22	Impaired insulin activation and dephosphorylation of glycogen synthase in skeletal muscle of women with polycystic ovary syndrome is reversed by pioglitazone treatment. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 3618-26	5.6	29
21	Total and high molecular weight (HMW) adiponectin levels and measures of glucose and lipid metabolism following pioglitazone treatment in a randomized placebo-controlled study in polycystic ovary syndrome. <i>Clinical Endocrinology</i> , 2008 , 68, 165-74	3.4	29
20	Association of fasting glucagon and proinsulin concentrations with insulin resistance. <i>Diabetologia</i> , 2007 , 50, 2342-7	10.3	63
19	Reduced expression of nuclear-encoded genes involved in mitochondrial oxidative metabolism in skeletal muscle of insulin-resistant women with polycystic ovary syndrome. <i>Diabetes</i> , 2007 , 56, 2349-55	0.9	135
18	Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes. <i>Diabetes</i> , 2007 , 56, 1592-9	0.9	397

17	Identification of the oxidized low-density lipoprotein scavenger receptor CD36 in plasma: a novel marker of insulin resistance. <i>Circulation</i> , 2006 , 114, 1169-76	16.7	117
16	Energy expenditure, body composition and insulin response to glucose in male twins discordant for the Trp64Arg polymorphism of the beta3-adrenergic receptor gene. <i>Diabetes, Obesity and Metabolism</i> , 2006 , 8, 322-30	6.7	15
15	Impaired glycogen synthase activity and mitochondrial dysfunction in skeletal muscle: markers or mediators of insulin resistance in type 2 diabetes?. <i>Current Diabetes Reviews</i> , 2006 , 2, 375-95	2.7	45
14	Free rather than total circulating insulin-like growth factor-I determines the feedback on growth hormone release in normal subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 366-71	5.6	59
13	Reduced insulin-mediated citrate synthase activity in cultured skeletal muscle cells from patients with type 2 diabetes: evidence for an intrinsic oxidative enzyme defect. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2005 , 1741, 206-14	6.9	66
12	Fasting unmasks a strong inverse association between ghrelin and cortisol in serum: studies in obese and normal-weight subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 741-6	5.6	116
11	Skeletal muscle lipid accumulation in type 2 diabetes may involve the liver X receptor pathway. <i>Diabetes</i> , 2005 , 54, 1108-15	0.9	81
10	A novel syndrome of autosomal-dominant hyperinsulinemic hypoglycemia linked to a mutation in the human insulin receptor gene. <i>Diabetes</i> , 2004 , 53, 1592-8	0.9	93
9	The primary defect in glycogen synthase activity is not based on increased glycogen synthase kinase-3alpha activity in diabetic myotubes. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 319, 1235-40	3.4	11
8	AMPK activity and isoform protein expression are similar in muscle of obese subjects with and without type 2 diabetes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2004 , 286, E239-44	6	68
7	Increased phosphorylation of skeletal muscle glycogen synthase at NH2-terminal sites during physiological hyperinsulinemia in type 2 diabetes. <i>Diabetes</i> , 2003 , 52, 1393-402	0.9	109
6	Does overnight normalization of plasma glucose by insulin infusion affect assessment of glucose metabolism in Type 2 diabetes?. <i>Diabetic Medicine</i> , 2003 , 20, 816-22	3.5	3
5	Proteome analysis reveals phosphorylation of ATP synthase beta -subunit in human skeletal muscle and proteins with potential roles in type 2 diabetes. <i>Journal of Biological Chemistry</i> , 2003 , 278, 10436-42	5.4	176
4	The diabetic phenotype is conserved in myotubes established from diabetic subjects: evidence for primary defects in glucose transport and glycogen synthase activity. <i>Diabetes</i> , 2002 , 51, 921-7	0.9	131
3	Development and clinical evaluation of a novel immunoassay for the binary complex of IGF-I and IGF-binding protein-1 in human serum. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 260-6	5.6	55
2	Reference intervals for glucose, beta-cell polypeptides, and counterregulatory factors during prolonged fasting. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2001 , 280, E50-8	6	60
1	Tissue Biopsies in Diabetes Research	265-288	1