

# Janny H Dekker

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2827/publications.pdf>

Version: 2024-02-01

39  
papers

481  
citations

686830

13  
h-index

713013

21  
g-index

40  
all docs

40  
docs citations

40  
times ranked

408  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of pelvic floor muscle training compared with watchful waiting in older women with symptomatic mild pelvic organ prolapse: randomised controlled trial in primary care. <i>BMJ</i> , The, 2014, 349, g7378-g7378.	3.0	70
2	Severity, not type, is the main predictor of decreased quality of life in elderly women with urinary incontinence: a population-based study as part of a randomized controlled trial in primary care. <i>Health and Quality of Life Outcomes</i> , 2012, 10, 153.	1.0	39
3	Systematic screening for urinary incontinence in older women: Who could benefit from it?. <i>Scandinavian Journal of Primary Health Care</i> , 2012, 30, 21-28.	0.6	36
4	Effectiveness and cost-effectiveness of pessary treatment compared with pelvic floor muscle training in older women with pelvic organ prolapse: 2-year follow-up of a randomized controlled trial in primary care. <i>Menopause</i> , 2016, 23, 1307-1318.	0.8	34
5	Predictors of unsuccessful pessary fitting in women with prolapse: a cross-sectional study in general practice. <i>International Urogynecology Journal</i> , 2017, 28, 307-313.	0.7	31
6	Pelvic floor muscle training versus watchful waiting or pessary treatment for pelvic organ prolapse (POPPS): Design and participant baseline characteristics of two parallel pragmatic randomized controlled trials in primary care. <i>Maturitas</i> , 2014, 77, 168-173.	1.0	27
7	Sexual function in older women with pelvic floor symptoms: a cross-sectional study in general practice. <i>British Journal of General Practice</i> , 2014, 64, e144-e150.	0.7	22
8	Minimal important change in the pelvic floor distress inventory-20 among women opting for conservative prolapse treatment. <i>American Journal of Obstetrics and Gynecology</i> , 2017, 216, 397.e1-397.e7.	0.7	21
9	The impact of a mobile application-based treatment for urinary incontinence in adult women: Design of a mixed-methods randomized controlled trial in a primary care setting. <i>Neurourology and Urodynamics</i> , 2018, 37, 2167-2176.	0.8	21
10	Levonorgestrel-releasing intrauterine system versus endometrial ablation for heavy menstrual bleeding. <i>American Journal of Obstetrics and Gynecology</i> , 2021, 224, 187.e1-187.e10.	0.7	20
11	App-Based Treatment in Primary Care for Urinary Incontinence: A Pragmatic, Randomized Controlled Trial. <i>Annals of Family Medicine</i> , 2021, 19, 102-109.	0.9	19
12	The effect of systematic screening of older women for urinary incontinence on treatment uptake: The URINO trial. <i>Maturitas</i> , 2013, 74, 334-340.	1.0	17
13	Incidence and treatment of heavy menstrual bleeding in general practice. <i>Family Practice</i> , 2017, 34, 673-678.	0.8	17
14	Active encouragement of older women with urinary incontinence in primary care to undergo diagnosis and treatment: A matched-pair cluster randomized controlled trial. <i>Maturitas</i> , 2015, 80, 212-219.	1.0	11
15	Physiotherapy for Children with Functional Constipation: A Pragmatic Randomized Controlled Trial in Primary Care. <i>Journal of Pediatrics</i> , 2020, 216, 25-31.e2.	0.9	11
16	One year effectiveness of an app-based treatment for urinary incontinence in comparison to care as usual in Dutch general practice: A pragmatic randomised controlled trial over 12 months. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2022, 129, 1474-1480.	1.1	11
17	Predictors of Success for Pelvic Floor Muscle Training in Pelvic Organ Prolapse. <i>Physical Therapy</i> , 2019, 99, 109-117.	1.1	10
18	Factors affecting patient recruitment to trials: qualitative research in general practice. <i>BJGP Open</i> , 2020, 4, bjgpopen20X101056.	0.9	10

#	ARTICLE	IF	CITATIONS
19	Free uroflowmetry for voiding dysfunction measurement in women with pelvic organ prolapse and urinary incontinence in primary care. <i>International Journal of Urology</i> , 2015, 22, 801-802.	0.5	8
20	Pelvic Physiotherapy in Children With Functional Constipation: Promising But More Research Needed. <i>Gastroenterology</i> , 2017, 152, 2080-2081.	0.6	8
21	Cost-effectiveness of an app-based treatment for urinary incontinence in comparison with care as usual in Dutch general practice: a pragmatic randomised controlled trial over 12 months. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2022, 129, 1538-1545.	1.1	7
22	Pelvic floor muscle therapy or alpha-blocking agents for treatment of men with lower urinary tract symptoms: An exploratory randomized controlled trial. <i>International Journal of Urology</i> , 2017, 24, 473-474.	0.5	5
23	Physiotherapy plus conventional treatment versus conventional treatment only in the treatment of functional constipation in children: design of a randomized controlled trial and cost-effectiveness study in primary care. <i>BMC Pediatrics</i> , 2018, 18, 249.	0.7	5
24	Barriers and Facilitators Associated With App-Based Treatment for Female Urinary Incontinence: Mixed Methods Evaluation. <i>JMIR MHealth and UHealth</i> , 2021, 9, e25878.	1.8	5
25	Is the hymen a suitable cut-off point for clinically relevant pelvic organ prolapse?. <i>Maturitas</i> , 2017, 99, 86-91.	1.0	3
26	Pelvic organ prolapse: prevention by training?. <i>Lancet, The</i> , 2017, 389, 336-337.	6.3	3
27	Cost-effectiveness of physiotherapy in childhood functional constipation: a randomized controlled trial in primary care. <i>Family Practice</i> , 2022, 39, 662-668.	0.8	3
28	More research is needed before we regard POP-Q stage 1 prolapse as normal. <i>International Urogynecology Journal</i> , 2014, 25, 1291-1291.	0.7	2
29	Cost-effectiveness of a nurse-led intervention to optimise implementation of guideline-concordant continence care: Study protocol of the COCON study. <i>BMC Nursing</i> , 2017, 16, 10.	0.9	2
30	Urethritis of cystitis: soa of urineweginfectie?. <i>Huisarts En Wetenschap</i> , 2015, 58, 25-25.	0.0	1
31	Authors' reply re: Cost-effectiveness of a proactive approach of urinary incontinence in women. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016, 123, 2229-2230.	1.1	1
32	Effect of lower VWF and FXI levels on levonorgestrel IUS and endometrial ablation treatment success in heavy menstrual bleeding: An exploratory study. <i>Haemophilia</i> , 2021, 27, e756-e759.	1.0	1
33	Prolaps. <i>Bijblijven (Amsterdam, Netherlands)</i> , 2015, 31, 756-769.	0.0	0
34	De bekkenbodembelicht. <i>Bijblijven (Amsterdam, Netherlands)</i> , 2015, 31, 707-719.	0.0	0
35	Mictieklachten bij mannen "meer dan een vergrote prostaat". <i>Bijblijven (Amsterdam, Netherlands)</i> , 2015, 31, 730-737.	0.0	0
36	Authors' reply to the comment by Hanis et al. on "Predictors of unsuccessful pessary fitting in women with prolapse: a cross-sectional study in general practice," by Panman et al.. <i>International Urogynecology Journal</i> , 2017, 28, 1441-1441.	0.7	0

#	ARTICLE	IF	CITATIONS
37	Testis niet in het scrotum. <i>Bijblijven</i> (Amsterdam, Netherlands), 2017, 33, 603-609.	0.0	0
38	Quality of life in children with functional constipation: Are child self-reports and parent proxy-reports interchangeable?. <i>Journal of Pediatrics</i> , 2020, 217, 216.	0.9	0
39	High rates of anorectal chlamydia in women: Cross-sectional study in general practice. <i>BJGP Open</i> , 2022, , BJGPO.2021.0223.	0.9	0