

Bing Hua

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2836456/publications.pdf>

Version: 2024-02-01

14
papers

185
citations

1162367

8
h-index

1281420

11
g-index

17
all docs

17
docs citations

17
times ranked

174
citing authors

#	ARTICLE	IF	CITATIONS
1	High triglyceride-glucose index is associated with adverse cardiovascular outcomes in patients with acute myocardial infarction. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 2351-2362.	1.1	46
2	Predictive effect of triglyceride-glucose index on clinical events in patients with type 2 diabetes mellitus and acute myocardial infarction: results from an observational cohort study in China. <i>Cardiovascular Diabetology</i> , 2021, 20, 43.	2.7	40
3	High Triglyceride-Glucose Index is Associated with Poor Cardiovascular Outcomes in Nondiabetic Patients with ACS with LDL-C below 1.8 mmol/L. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 268-281.	0.9	32
4	Pregnancy-Associated Plasma Protein A Induces Inflammatory Cytokine Expression by Activating IGF-1/PI3K/Akt Pathways. <i>Mediators of Inflammation</i> , 2019, 2019, 1-12.	1.4	10
5	Liraglutide inhibited AGEs induced coronary smooth muscle cell phenotypic transition through inhibiting the NF- κ B signal pathway. <i>Peptides</i> , 2019, 112, 125-132.	1.2	10
6	Real-world use of ACEI/ARB in diabetic hypertensive patients before the initial diagnosis of obstructive coronary artery disease: patient characteristics and long-term follow-up outcome. <i>Journal of Translational Medicine</i> , 2020, 18, 150.	1.8	10
7	Dynamic fluctuations of advanced glycation end products and its C-terminal truncated receptor level in patients with acute ST-segment elevation myocardial infarction and undergoing diabetes or not. <i>Medicine (United States)</i> , 2018, 97, e11278.	0.4	9
8	AGEs impair Kv channel-mediated vasodilation of coronary arteries by activating the NF- κ B signaling pathway in ZDF rats. <i>Biomedicine and Pharmacotherapy</i> , 2019, 120, 109527.	2.5	9
9	Pioglitazone downregulates Twist-1 expression in the kidney and protects renal function of Zucker diabetic fatty rats. <i>Biomedicine and Pharmacotherapy</i> , 2019, 118, 109346.	2.5	9
10	Protective role of activating PPAR γ 3 in advanced glycation end products-induced impairment of coronary artery vasodilation via inhibiting p38 phosphorylation and reactive oxygen species production. <i>Biomedicine and Pharmacotherapy</i> , 2022, 147, 112641.	2.5	8
11	TCTAP A-051 Nicorandil Protects Cardiac Microvascular Endothelial Cells from Advanced Glycation End Products Induced Cytotoxicity by Promoting Autophagy. <i>Journal of the American College of Cardiology</i> , 2017, 69, S27.	1.2	1
12	Role of peroxisome proliferators-activated receptor-gamma in advanced glycation end product-mediated functional loss of voltage-gated potassium channel in rat coronary arteries. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 337.	0.7	1
13	GLYCATED ALBUMIN IS A PREDICTOR OF MAJOR ADVERSE CARDIO-CEREBRAL EVENTS IN NONDIABETIC PATIENTS WITH ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION UNDERWENT PERCUTANEOUS CORONARY INTERVENTION. <i>Journal of the American College of Cardiology</i> , 2019, 73, 86.	1.2	0
14	ADVANCED GLYCATION END PRODUCTS IMPAIRED VOLTAGE-GATED K+ CHANNEL ON RAT CORONARY SMOOTH MUSCLE CELLS BY ACTIVATING NF-KB SIGNALING PATHWAY. <i>Journal of the American College of Cardiology</i> , 2019, 73, 2059.	1.2	0