

Michael P Stone

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2836562/michael-p-stone-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

96
papers

2,717
citations

29
h-index

47
g-index

104
ext. papers

2,883
ext. citations

6.4
avg, IF

4.42
L-index

#	Paper	IF	Citations
96	Structure of a Stable Interstrand DNA Cross-Link Involving a β -Glycosyl Linkage Between an -dA Amino Group and an Abasic Site. <i>Biochemistry</i> , 2021 , 60, 41-52	3.2	3
95	DNA Sequence Modulates the Efficiency of NEIL1-Catalyzed Excision of the Aflatoxin B-Induced Formamidopyrimidine Guanine Adduct. <i>Chemical Research in Toxicology</i> , 2021 , 34, 901-911	4	0
94	Recognition of DNA adducts by edited and unedited forms of DNA glycosylase NEIL1. <i>DNA Repair</i> , 2020 , 85, 102741	4.3	7
93	Characterization of rare NEIL1 variants found in East Asian populations. <i>DNA Repair</i> , 2019 , 79, 32-39	4.3	4
92	Processing of N-substituted formamidopyrimidine DNA adducts by DNA glycosylases NEIL1 and NEIL3. <i>DNA Repair</i> , 2019 , 73, 49-54	4.3	10
91	Configurational and Conformational Equilibria of N-(2-Deoxy-d-erythro-pentofuranosyl)-2,6-diamino-3,4-dihydro-4-oxo-5-N-methylformamidopyrimidine (MeFapy-dG) Lesion in DNA. <i>Chemical Research in Toxicology</i> , 2018 , 31, 924-935	4	2
90	NEIL1 protects against aflatoxin-induced hepatocellular carcinoma in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, 4207-4212	11.5	26
89	DNA polymerase β limits chromosomal damage and promotes cell survival following aflatoxin exposure. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 13774-13779	11.5	17
88	Mechanism of Error-Free Bypass of the Environmental Carcinogen N-(2-Deoxyguanosin-8-yl)-3-aminobenzanthrone Adduct by Human DNA Polymerase β . <i>ChemBioChem</i> , 2016 , 17, 2033-2037	3.8	5
87	Base-Displaced Intercalated Conformation of the 2-Amino-3-methylimidazo[4,5-f]quinoline N(2)-dG DNA Adduct Positioned at the Nonreiterated G(1) in the NarI Restriction Site. <i>Chemical Research in Toxicology</i> , 2015 , 28, 1455-68	4	5
86	Structural Basis for Error-Free Bypass of the 5-N-Methylformamidopyrimidine-dG Lesion by Human DNA Polymerase β and <i>Sulfolobus solfataricus</i> P2 Polymerase IV. <i>Journal of the American Chemical Society</i> , 2015 , 137, 7011-4	16.4	12
85	Base-Displaced Intercalated Structure of the N-(2-Deoxyguanosin-8-yl)-3-aminobenzanthrone DNA Adduct. <i>Chemical Research in Toxicology</i> , 2015 , 28, 2253-66	4	8
84	DNA Sequence Modulates Geometrical Isomerism of the trans-8,9-Dihydro-8-(2,6-diamino-4-oxo-3,4-dihydropyrimid-5-yl-formamido)-9-hydroxy Aflatoxin B1 Adduct. <i>Chemical Research in Toxicology</i> , 2015 , 28, 225-37	4	8
83	Molecular basis of aflatoxin-induced mutagenesis-role of the aflatoxin B1-formamidopyrimidine adduct. <i>Carcinogenesis</i> , 2014 , 35, 1461-8	4.6	37
82	Major groove orientation of the (2S)-N(6)-(2-hydroxy-3-buten-1-yl)-2-deoxyadenosine DNA adduct induced by 1,2-epoxy-3-butene. <i>Chemical Research in Toxicology</i> , 2014 , 27, 1675-86	4	5
81	Structures of exocyclic R,R- and S,S-N(6),N(6)-(2,3-dihydroxybutan-1,4-diyl)-2-deoxyadenosine adducts induced by 1,2,3,4-diepoxybutane. <i>Chemical Research in Toxicology</i> , 2014 , 27, 805-17	4	10
80	Error-prone replication bypass of the primary aflatoxin B1 DNA adduct, AFB1-N7-Gua. <i>Journal of Biological Chemistry</i> , 2014 , 289, 18497-506	5.4	35

79	Molecular mechanisms underlying aflatoxin-induced mutagenesis. <i>FASEB Journal</i> , 2013 , 27, lb78	0.9	
78	Structure of (5S)-8,5Vcyclo-2Vdeoxyguanosine in DNA. <i>Journal of the American Chemical Society</i> , 2011 , 133, 20357-68	16.4	38
77	Chemistry and Biology of Aflatoxin-DNA Adducts. <i>ACS Symposium Series</i> , 2011 , 147-166	0.4	3
76	Chemistry and structural biology of DNA damage and biological consequences. <i>Chemistry and Biodiversity</i> , 2011 , 8, 1571-615	2.5	36
75	Hydroxy-1,N2-propano-2Vdeoxyguanosine DNA adduct conjugates the N-terminal amine of the KWKK peptide via a carbinolamine linkage. <i>Chemical Research in Toxicology</i> , 2011 , 24, 1123-33	4	5
74	Bypass of aflatoxin B1 adducts by the <i>Sulfolobus solfataricus</i> DNA polymerase IV. <i>Journal of the American Chemical Society</i> , 2011 , 133, 12556-68	16.4	29
73	Formation of a N2-dG:N2-dG carbinolamine DNA cross-link by the trans-4-hydroxynonenal-derived (6S,8R,11S) 1,N2-dG adduct. <i>Journal of the American Chemical Society</i> , 2011 , 133, 16101-10	16.4	10
72	Structural Consequences of Epimerization of Thymine Glycol Lesions in Duplex DNA: Implications for DNA Repair. <i>ACS Symposium Series</i> , 2010 , 11-28	0.4	
71	Selective Incision of the alpha-N-Methyl-Formamidopyrimidine Anomer by <i>Escherichia coli</i> Endonuclease IV. <i>Journal of Nucleic Acids</i> , 2010 , 2010,	2.3	23
70	Binding of the human nucleotide excision repair proteins XPA and XPC/HR23B to the 5R-thymine glycol lesion and structure of the cis-(5R,6S) thymine glycol epimer in the 5VGTgG-3Vsequence: destabilization of two base pairs at the lesion site. <i>Nucleic Acids Research</i> , 2010 , 38, 428-40	20.1	60
69	DNA cross-link induced by trans-4-hydroxynonenal. <i>Environmental and Molecular Mutagenesis</i> , 2010 , 51, 625-34	3.2	35
68	Inherent stereospecificity in the reaction of aflatoxin B(1) 8,9-epoxide with deoxyguanosine and efficiency of DNA catalysis. <i>Chemical Research in Toxicology</i> , 2009 , 22, 913-7	4	29
67	Conformational interconversion of the trans-4-hydroxynonenal-derived (6S,8R,11S) 1,N(2)-deoxyguanosine adduct when mismatched with deoxyadenosine in DNA. <i>Chemical Research in Toxicology</i> , 2009 , 22, 187-200	4	12
66	The cis-(5R,6S)-thymine glycol lesion occupies the wobble position when mismatched with deoxyguanosine in DNA. <i>Biochemistry</i> , 2009 , 48, 9722-33	3.2	19
65	Stereospecific formation of the (R)-gamma-hydroxytrimethylene interstrand N2-dG:N2-dG cross-link arising from the gamma-OH-1,N2-propano-2Vdeoxyguanosine adduct in the 5VcPcG-3V DNA sequence. <i>Journal of the American Chemical Society</i> , 2009 , 131, 8416-24	16.4	13
64	Chemistry and biology of DNA containing 1,N(2)-deoxyguanosine adducts of the alpha,beta-unsaturated aldehydes acrolein, crotonaldehyde, and 4-hydroxynonenal. <i>Chemical Research in Toxicology</i> , 2009 , 22, 759-78	4	141
63	Structural perturbations induced by the alpha-anomer of the aflatoxin B(1) formamidopyrimidine adduct in duplex and single-strand DNA. <i>Journal of the American Chemical Society</i> , 2009 , 131, 16096-107	16.4	20
62	Differential base stacking interactions induced by trimethylene interstrand DNA cross-links in the 5VcPcG-3Vand 5VgPc-3Vsequence contexts. <i>Chemical Research in Toxicology</i> , 2009 , 22, 1810-6	4	9

61	Structure of the 1,N2-ethenodeoxyguanosine adduct opposite cytosine in duplex DNA: Hoogsteen base pairing at pH 5.2. <i>Chemical Research in Toxicology</i> , 2008 , 21, 1795-805	4	14
60	The stereochemistry of trans-4-hydroxynonenal-derived exocyclic 1,N2-2Vdeoxyguanosine adducts modulates formation of interstrand cross-links in the 5VCpG-3Vsequence. <i>Biochemistry</i> , 2008 , 47, 11457-72	3.2	17
59	Insertion of dNTPs opposite the 1,N2-propanodeoxyguanosine adduct by <i>Sulfolobus solfataricus</i> P2 DNA polymerase IV. <i>Biochemistry</i> , 2008 , 47, 7322-34	3.2	21
58	Interconversion of the cis-5R,6S- and trans-5R,6R-thymine glycol lesions in duplex DNA. <i>Journal of the American Chemical Society</i> , 2008 , 130, 11701-10	16.4	27
57	Rearrangement of the (6S,8R,11S) and (6R,8S,11R) exocyclic 1,N2-deoxyguanosine adducts of trans-4-hydroxynonenal to N2-deoxyguanosine cyclic hemiacetal adducts when placed complementary to cytosine in duplex DNA. <i>Journal of the American Chemical Society</i> , 2008 , 130, 10898-906	16.4	16
56	Interstrand DNA cross-links induced by alpha,beta-unsaturated aldehydes derived from lipid peroxidation and environmental sources. <i>Accounts of Chemical Research</i> , 2008 , 41, 793-804	24.3	142
55	Site-specific synthesis and characterization of oligonucleotides containing an N6-(2-deoxy-D-erythro-pentofuranosyl)-2,6-diamino-3,4-dihydro-4-oxo-5-N-methylformamidopyrimidine lesion, the ring-opened product from N7-methylation of deoxyguanosine. <i>Chemical Research in Toxicology</i> , 2008 , 21, 2324-33	4	26
54	Stereochemistry modulates the stability of reduced interstrand cross-links arising from R- and S-alpha-CH3-gamma-OH-1,N2-propano-2Vdeoxyguanosine in the 5VCpG-3VDNA sequence. <i>Biochemistry</i> , 2007 , 46, 2608-21	3.2	16
53	Structure of the 1,N2-etheno-2Vdeoxyguanosine adduct in duplex DNA at pH 8.6. <i>Chemical Research in Toxicology</i> , 2007 , 20, 1601-11	4	15
52	Bulge migration of the malondialdehyde OPdG DNA adduct when placed opposite a two-base deletion in the (CpG)3 frameshift hotspot of the <i>Salmonella typhimurium</i> hisD3052 gene. <i>Chemical Research in Toxicology</i> , 2007 , 20, 1200-10	4	8
51	Base-displaced intercalated structure of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the <i>NarI</i> restriction enzyme, a hotspot for -2 bp deletions. <i>Journal of the American Chemical Society</i> , 2006 , 128, 10085-95	16.4	57
50	Stereospecific formation of interstrand carbinolamine DNA cross-links by crotonaldehyde- and acetaldehyde-derived alpha-CH3-gamma-OH-1,N2-propano-2Vdeoxyguanosine adducts in the 5VCpG-3Vsequence. <i>Chemical Research in Toxicology</i> , 2006 , 19, 195-208	4	49
49	Orientation of the crotonaldehyde-derived N2-[3-Oxo-1(S)-methyl-propyl]-dG DNA adduct hinders interstrand cross-link formation in the 5VCpG-3Vsequence. <i>Chemical Research in Toxicology</i> , 2006 , 19, 1019-29	4	19
48	Unraveling the aflatoxin-FAPY conundrum: structural basis for differential replicative processing of isomeric forms of the formamidopyrimidine-type DNA adduct of aflatoxin B1. <i>Journal of the American Chemical Society</i> , 2006 , 128, 15188-99	16.4	49
47	Spectroscopic characterization of interstrand carbinolamine cross-links formed in the 5VCpG-3V sequence by the acrolein-derived gamma-OH-1,N2-propano-2Vdeoxyguanosine DNA adduct. <i>Journal of the American Chemical Society</i> , 2005 , 127, 17686-96	16.4	36
46	In vitro bypass of malondialdehyde-deoxyguanosine adducts: differential base selection during extension by the Klenow fragment of DNA polymerase I is the critical determinant of replication outcome. <i>Biochemistry</i> , 2004 , 43, 11828-35	3.2	21
45	Stereospecific structural perturbations arising from adenine N(6) butadiene triol adducts in duplex DNA. <i>Chemical Research in Toxicology</i> , 2004 , 17, 1007-19	4	15
44	Structure of a site specific major groove (2S,3S)-N6-(2,3,4-trihydroxybutyl)-2Vdeoxyadenosyl DNA adduct of butadiene diol epoxide. <i>Chemical Research in Toxicology</i> , 2004 , 17, 717-30	4	15

43	Wobble dC.dA pairing 5Vto the cationic guanine N7 8,9-dihydro-8-(N7-guanyl)-9-hydroxyaflatoxin B1 adduct: implications for nontargeted AFB1 mutagenesis. <i>Biochemistry</i> , 2003 , 42, 7023-34	3.2	12
42	DNA interchain cross-links formed by acrolein and crotonaldehyde. <i>Journal of the American Chemical Society</i> , 2003 , 125, 50-61	16.4	167
41	NMR determination of the conformation of a trimethylene interstrand cross-link in an oligodeoxynucleotide duplex containing a 5Vd(GpC) motif. <i>Journal of the American Chemical Society</i> , 2003 , 125, 62-72	16.4	36
40	Thermal stabilization of the DNA duplex by adducts of aflatoxin B1. <i>Biopolymers</i> , 2002 , 65, 190-201	2.2	13
39	Structure of the 1,N(2)-propanodeoxyguanosine adduct in a three-base DNA hairpin loop derived from a palindrome in the Salmonella typhimurium hisD3052 gene. <i>Chemical Research in Toxicology</i> , 2002 , 15, 140-52	4	12
38	Detection of an interchain carbinolamine cross-link formed in a CpG sequence by the acrolein DNA adduct gamma-OH-1,N(2)-propano-2Vdeoxyguanosine. <i>Journal of the American Chemical Society</i> , 2002 , 124, 9324-5	16.4	44
37	Structure of an oligodeoxynucleotide containing a 1,N(2)-propanodeoxyguanosine adduct positioned in a palindrome derived from the Salmonella typhimurium hisD3052 gene: Hoogsteen pairing at pH 5.2. <i>Chemical Research in Toxicology</i> , 2002 , 15, 127-39	4	32
36	Structural refinement of the 8,9-dihydro-8-(N7-guanyl)-9-hydroxy-aflatoxin B(1) adduct in a 5VCp(AFB)G-3Vsequence. <i>Chemical Research in Toxicology</i> , 2002 , 15, 638-47	4	20
35	Mispairing of the 8,9-dihydro-8-(N7-guanyl)-9-hydroxy-aflatoxin B1 adduct with deoxyadenosine results in extrusion of the mismatched dA toward the major groove. <i>Biochemistry</i> , 2002 , 41, 5462-72	3.2	11
34	Structural studies of an oligodeoxynucleotide containing a trimethylene interstrand cross-link in a 5V(CpG) motif: model of a malondialdehyde cross-link. <i>Journal of the American Chemical Society</i> , 2001 , 123, 1730-9	16.4	52
33	The exocyclic 1,N2-deoxyguanosine pyrimidopurinone M1G is a chemically stable DNA adduct when placed opposite a two-base deletion in the (CpG)3 frameshift hotspot of the Salmonella typhimurium hisD3052 gene. <i>Biochemistry</i> , 2001 , 40, 15638-49	3.2	15
32	The nonmutagenic (R)- and (S)-beta-(N(6)-adenyl)styrene oxide adducts are oriented in the major groove and show little perturbation to DNA structure. <i>Biochemistry</i> , 2001 , 40, 9780-91	3.2	13
31	Intercalation of the (1R,2S,3R,4S)-N6-[1-(1,2,3,4-tetrahydro-2,3,4-trihydroxybenz[a]anthracenyl)]-2Vdeoxyadenosyl adduct in the N-ras codon 61 sequence: DNA sequence effects. <i>Biochemistry</i> , 2001 , 40, 6743-55	3.2	16
30	Structure of the malondialdehyde deoxyguanosine adduct M1G when placed opposite a two-base deletion in the (CpG)3 frameshift hotspot of the Salmonella typhimurium hisD3052 gene. <i>Advances in Experimental Medicine and Biology</i> , 2001 , 500, 513-6	3.6	5
29	Synthesis of oligonucleotides containing the alkali-labile pyrimidopurinone adduct, M(1)G. <i>Chemical Research in Toxicology</i> , 2000 , 13, 90-5	4	24
28	Replication of a site-specific trans-8,9-dihydro-8-(N7-guanyl)-9-hydroxyaflatoxin B(1) adduct by the exonuclease deficient klenow fragment of DNA polymerase I. <i>Chemical Research in Toxicology</i> , 2000 , 13, 1158-64	4	11
27	Solution structure of an oligodeoxynucleotide containing the malondialdehyde deoxyguanosine adduct N2-(3-oxo-1-propenyl)-dG (ring-opened M1G) positioned in a (CpG)3 frameshift hotspot of the Salmonella typhimurium hisD3052 gene. <i>Biochemistry</i> , 1999 , 38, 13491-501	3.2	28
26	Intercalation of the (1S,2R,3S,4R)-N6-[1-(1,2,3,4-tetrahydro-2,3,4-trihydroxybenz[a]anthracenyl)]-2Vdeoxyadenosyl adduct in an oligodeoxynucleotide containing the human N-ras codon 61 sequence. <i>Biochemistry</i> , 1999 , 38, 16045-57	3.2	20

25	Site-specific synthesis of aflatoxin B(1) adducts within an oligodeoxyribonucleotide containing the human p53 codon 249 sequence. <i>Chemical Research in Toxicology</i> , 1999 , 12, 707-14	4	5
24	DNA abasic lesions in a different light: solution structure of an endogenous topoisomerase II poison. <i>Biochemistry</i> , 1999 , 38, 15500-7	3.2	36
23	Intercalation of the (-)-(1R,2S,3R, 4S)-N6-[1-benz[a]anthracenyl]-2' deoxyadenosyl adduct in an oligodeoxynucleotide containing the human N-ras codon 61 sequence. <i>Biochemistry</i> , 1999 , 38, 2969-81	3.2	17
22	Influence of the R(61,2)- and S(61,2)-alpha-(N6-adenyl)styrene oxide adducts on the A.C mismatched base pair in an oligodeoxynucleotide containing the human N-ras codon 61. <i>Biochemistry</i> , 1999 , 38, 8635-46	3.2	7
21	Refined structure of the doubly intercalated d(TATAFBGCATA) ₂ aflatoxin B1 adduct. <i>Chemical Research in Toxicology</i> , 1998 , 11, 873-81	4	19
20	Multiple conformations of an intercalated (-)-(7S,8R,9S, 10R)-N6-[10-(7,8,9,10-tetrahydrobenzo[a]pyrenyl)]-2' deoxyadenosyl adduct in the N-ras codon 61 sequence. <i>Biochemistry</i> , 1998 , 37, 16516-28	3.2	39
19	An intercalated and thermally stable FAPY adduct of aflatoxin B1 in a DNA duplex: structural refinement from 1H NMR. <i>Biochemistry</i> , 1998 , 37, 4374-87	3.2	56
18	Site-specific targeting of aflatoxin adduction directed by triple helix formation in the major groove of oligodeoxyribonucleotides. <i>Nucleic Acids Research</i> , 1998 , 26, 1070-5	20.1	9
17	Major groove (S)-alpha-(N6-adenyl)styrene oxide adducts in an oligodeoxynucleotide containing the human N-ras codon 61 sequence: conformations of the S(61,2) and S(61,3) sequence isomers from 1H NMR. <i>Biochemistry</i> , 1996 , 35, 7316-29	3.2	35
16	Adduction of the human N-ras codon 61 sequence with (-)-(7S,8R,9R,10S)-7,8-dihydroxy-9,10-epoxy-7,8,9,10-tetrahydrobenzo[a] pyrene: structural refinement of the intercalated SRSR(61,2) (-)-(7S,8R,9S,10R)-N6-[10-(7,8,9,10-tetrahydrobenzo[a]pyrenyl)]-2' deoxyadenosyl adduct from 1H NMR. <i>Biochemistry</i> , 1996 , 35, 6913-24	3.2	71
15	Sequence and Stereospecific Consequences of Major Groove [(N6-Adenyl)] Styrene Oxide Adducts in an Oligodeoxynucleotide Containing the Human N-ras Codon 61 Sequence. <i>Magnetic Resonance in Chemistry</i> , 1996 , 34, S105-S114	2.1	5
14	Structure of a duplex oligodeoxynucleotide containing propanodeoxyguanosine opposite a two-base deletion in the (CpG) ₃ frame shift hotspot of Salmonella typhimurium hisD3052 determined by 1H NMR and restrained molecular dynamics. <i>Biochemistry</i> , 1995 , 34, 50-64	3.2	29
13	Major groove (R)-alpha-(N6-adenyl)styrene oxide adducts in an oligodeoxynucleotide containing the human N-ras codon 61 sequence: conformations of the R(61,2) and R(61,3) sequence isomers from 1H NMR. <i>Biochemistry</i> , 1995 , 34, 14021-36	3.2	31
12	1H NMR characterization of a duplex oligodeoxynucleotide containing propanodeoxyguanosine opposite a two-base deletion in the (CpG) ₃ frameshift hotspot of Salmonella typhimurium hisD3052. <i>Chemical Research in Toxicology</i> , 1994 , 7, 319-28	4	22
11	1H NMR of an oligodeoxynucleotide containing a propanodeoxyguanosine adduct positioned in a (CG) ₃ frameshift hotspot of Salmonella typhimurium hisD3052: Hoogsteen base-pairing at pH 5.8. <i>Chemical Research in Toxicology</i> , 1993 , 6, 825-36	4	54
10	DNA conformation mediates aflatoxin B1-DNA binding and the formation of guanine N7 adducts by aflatoxin B1 8,9-exo-epoxide. <i>Chemical Research in Toxicology</i> , 1993 , 6, 64-8	4	69
9	A postoligomerization synthesis of oligodeoxynucleotides containing polycyclic aromatic hydrocarbon adducts at the N6 position of deoxyadenosine.. <i>Journal of the American Chemical Society</i> , 1992 , 114, 5480-5481	16.4	64
8	Alteration of the aflatoxin cyclopentenone ring to a delta-lactone reduces intercalation with DNA and decreases formation of guanine N7 adducts by aflatoxin epoxides. <i>Chemical Research in Toxicology</i> , 1990 , 3, 254-61	4	38

7	Intercalation of aflatoxin B1 in two oligodeoxynucleotide adducts: comparative ¹ H NMR analysis of d(ATCAFBGAT).d(ATCGAT) and d(ATAFBGCAT) ² . <i>Biochemistry</i> , 1990 , 29, 10438-48	3.2	90
6	Aflatoxin-DNA Binding and the Characterization of Aflatoxin B1-Oligodeoxynucleotide Adducts by ¹ H NMR Spectroscopy. <i>Jerusalem Symposia on Quantum Chemistry and Biochemistry</i> , 1990 , 451-480		4
5	THIONO COMPOUNDS. 10. STRUCTURES AND REACTIONS OF INTERMEDIATES FROM THE OXIDATION OF PHOSPHOROTHIOATES ^{1,2} . <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 1989 , 44, 39-52	1	2
4	Preparation of the 8,9-epoxide of the mycotoxin aflatoxin B1: the ultimate carcinogenic species. <i>Journal of the American Chemical Society</i> , 1988 , 110, 7929-7931	16.4	222
3	THIONO COMPOUNDS. 9. USE OF SPECTRA TO STUDY INTERMEDIATES IN THE OXIDATION OF THIONO PHOSPHORUS, COMPOUNDS. <i>Phosphorous and Sulfur and the Related Elements</i> , 1988 , 35, 159-172		3
2	Carcinogen-nucleic acid interactions: equilibrium binding studies of aflatoxins B1 and B2 with DNA and the oligodeoxynucleotide d(ATGCAT) ² . <i>Journal of Biomolecular Structure and Dynamics</i> , 1988 , 5, 1023-41	2.6	18
1	Interstrand DNA Cross-Linking 1,N ² -Deoxyguanosine Adducts Derived from β -Unsaturated Aldehydes: StructureFunction Relationships ²⁰¹⁻²¹⁶		2