

Mohsen Kerkeni

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/284473/publications.pdf>

Version: 2024-02-01

19
papers

542
citations

687363

13
h-index

888059

17
g-index

19
all docs

19
docs citations

19
times ranked

738
citing authors

#	ARTICLE	IF	CITATIONS
1	Pentosidine as a biomarker for microvascular complications in type 2 diabetic patients. <i>Diabetes and Vascular Disease Research</i> , 2013, 10, 239-245.	2.0	64
2	Comparative Study on in Vitro Effects of Homocysteine Thiolactone and Homocysteine on HUVEC Cells: Evidence for a Stronger Proapoptotic and Proinflammatory Homocysteine Thiolactone. <i>Molecular and Cellular Biochemistry</i> , 2006, 291, 119-126.	3.1	58
3	Hyperhomocysteinemia, paraoxonase activity and risk of coronary artery disease. <i>Clinical Biochemistry</i> , 2006, 39, 821-825.	1.9	58
4	Hyperhomocysteinemia, Endothelial Nitric Oxide Synthase Polymorphism, and Risk of Coronary Artery Disease. <i>Clinical Chemistry</i> , 2006, 52, 53-58.	3.2	50
5	Hyperhomocysteinemia, methylenetetrahydrofolate reductase polymorphism and risk of coronary artery disease. <i>Annals of Clinical Biochemistry</i> , 2006, 43, 200-206.	1.6	47
6	Elevated serum levels of AGEs, sRAGE, and pentosidine in Tunisian patients with severity of diabetic retinopathy. <i>Microvascular Research</i> , 2012, 84, 378-383.	2.5	41
7	Plasma Levels of Pentosidine, Carboxymethyl-Lysine, Soluble Receptor for Advanced Glycation End Products, and Metabolic Syndrome: The Metformin Effect. <i>Disease Markers</i> , 2016, 2016, 1-8.	1.3	39
8	Increased serum concentrations of pentosidine are related to presence and severity of coronary artery disease. <i>Thrombosis Research</i> , 2014, 134, 633-638.	1.7	37
9	Increased serum homocitrulline concentrations are associated with the severity of coronary artery disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2015, 53, 103-10.	2.3	31
10	RAGE receptor: May be a potential inflammatory mediator for SARS-COV-2 infection?. <i>Medical Hypotheses</i> , 2020, 144, 109950.	1.5	27
11	Hyperhomocysteinemia and parameters of antioxidative defence in Tunisian patients with coronary heart disease. <i>Annals of Clinical Biochemistry</i> , 2008, 45, 193-198.	1.6	19
12	Methylglyoxal: A Relevant Marker of Disease Activity in Patients with Rheumatoid Arthritis. <i>Disease Markers</i> , 2018, 2018, 1-6.	1.3	17
13	Increased serum concentrations of N ^ε -carboxymethyllysine are related to the presence and the severity of rheumatoid arthritis. <i>Annals of Clinical Biochemistry</i> , 2018, 55, 430-436.	1.6	15
14	Endothelial nitric oxide synthetase, methylenetetrahydrofolate reductase polymorphisms, and cardiovascular complications in Tunisian patients with nondiabetic renal disease. <i>Clinical Biochemistry</i> , 2009, 42, 958-964.	1.9	14
15	Hyperhomocysteinemia, paraoxonase concentration and cardiovascular complications in Tunisian patients with nondiabetic renal disease. <i>Clinical Biochemistry</i> , 2009, 42, 777-782.	1.9	11
16	Skin Autofluorescence Measurement in Subclinical Atheromatous Disease: Results from the ILERVAS Project. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 879-889.	2.0	9
17	Apolipoprotein A-I, apolipoprotein B, high-sensitivity C-reactive protein and severity of coronary artery disease in tunisian population. <i>Cardiovascular Pathology</i> , 2012, 21, 455-460.	1.6	5
18	Advanced Glycation End Products Levels and Carotid Intima Media Thickness in Patients With Rheumatoid Arthritis. <i>Journal of Biology and Life Science</i> , 2019, 10, 58.	0.2	0

#	ARTICLE	IF	CITATIONS
19	Weak Association between Skin Autofluorescence Levels and Prediabetes with an ILERVAS Cross-Sectional Study. <i>Nutrients</i> , 2022, 14, 1102.	4.1	0