

Robert Z Qi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/285448/publications.pdf>

Version: 2024-02-01

21
papers

980
citations

758635

12
h-index

794141

19
g-index

21
all docs

21
docs citations

21
times ranked

1307
citing authors

#	ARTICLE	IF	CITATIONS
1	Protocol to measure inter-centrosome distance in adherent cells using epifluorescence microscopy. STAR Protocols, 2022, 3, 101227.	0.5	0
2	Proteolysis of β -tubulin small complex proteins is mediated by the ubiquitin-proteasome system. FEBS Letters, 2021, 595, 1987-1996.	1.3	4
3	Nek2-mediated GAS2L1 phosphorylation and centrosome-linker disassembly induce centrosome disjunction. Journal of Cell Biology, 2020, 219, .	2.3	12
4	Cytolinker Gas2L1 regulates axon morphology through microtubule-modulated actin stabilization. EMBO Reports, 2019, 20, e47732.	2.0	45
5	The catalytic subunit of DNA polymerase δ is a nucleocytoplasmic shuttling protein. Experimental Cell Research, 2019, 375, 36-40.	1.2	5
6	Epigenetic Alterations of Wnt Signaling Pathways in Nasopharyngeal Carcinoma. Current Pharmacology Reports, 2018, 4, 337-345.	1.5	0
7	Microtubule plus-end tracking of end-binding protein 1 (EB1) is regulated by CDK5 regulatory subunit-associated protein 2. Journal of Biological Chemistry, 2017, 292, 7675-7687.	1.6	7
8	GAS2L1 Is a Centriole-Associated Protein Required for Centrosome Dynamics and Disjunction. Developmental Cell, 2017, 40, 81-94.	3.1	31
9	The catalytic subunit of DNA polymerase δ inhibits β TuRC activity and regulates Golgi-derived microtubules. Nature Communications, 2017, 8, 554.	5.8	12
10	Noncontact Viscoelastic Imaging of Living Cells Using a Long-Needle Atomic Force Microscope with Dual-Frequency Modulation. Physical Review Applied, 2017, 8, .	1.5	21
11	Molecular Pathway of Microtubule Organization at the Golgi Apparatus. Developmental Cell, 2016, 39, 44-60.	3.1	114
12	Transmembrane and Coiled-Coil Domain Family 1 Is a Novel Protein of the Endoplasmic Reticulum. PLoS ONE, 2014, 9, e85206.	1.1	15
13	Assaying Microtubule Nucleation by the β -Tubulin Ring Complex. Methods in Enzymology, 2014, 540, 119-130.	0.4	5
14	A novel myomegalin isoform functions in Golgi microtubule organization and ER-Golgi transport. Journal of Cell Science, 2014, 127, 4904-17.	1.2	46
15	NME7 is a functional component of the β -tubulin ring complex. Molecular Biology of the Cell, 2014, 25, 2017-2025.	0.9	64
16	Dynamic Recruitment of CDK5RAP2 to Centrosomes Requires Its Association with Dynein. PLoS ONE, 2013, 8, e68523.	1.1	25
17	The neuronal p35 activator of Cdk5 is a novel F-actin binding and bundling protein. Cellular and Molecular Life Sciences, 2011, 68, 1633-1643.	2.4	16
18	Identification of p10 as a neurotoxic product generated from the proteolytic cleavage of the neuronal Cdk5 activator. Journal of Cellular Biochemistry, 2010, 111, 1359-1366.	1.2	6

#	ARTICLE	IF	CITATIONS
19	CDK5RAP2 stimulates microtubule nucleation by the $\hat{\beta}$ -tubulin ring complex. <i>Journal of Cell Biology</i> , 2010, 191, 1089-1095.	2.3	248
20	CDK5RAP2 Is a Pericentriolar Protein That Functions in Centrosomal Attachment of the $\hat{\beta}$ -Tubulin Ring Complex. <i>Molecular Biology of the Cell</i> , 2008, 19, 115-125.	0.9	242
21	The Protein SET Binds the Neuronal Cdk5 Activator p35 and Modulates Cdk5/p35 Activity. <i>Journal of Biological Chemistry</i> , 2002, 277, 7324-7332.	1.6	62