

Hideyuku Inui

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2864914/hideyuku-inui-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

65

papers

998

citations

20

h-index

29

g-index

68

ext. papers

1,157

ext. citations

4.8

avg, IF

4.17

L-index

#	Paper	IF	Citations
65	Differential uptake for dioxin-like compounds by zucchini subspecies. <i>Chemosphere</i> , 2008 , 73, 1602-7	8.4	71
64	Herbicide resistance in transgenic plants with mammalian P450 monooxygenase genes. <i>Pest Management Science</i> , 2005 , 61, 286-91	4.6	60
63	Metabolism of Herbicides and Other Chemicals in Human Cytochrome P450 Species and in Transgenic Potato Plants Co-Expressing Human CYP1A1, CYP2B6 and CYP2C19. <i>Journal of Pesticide Sciences</i> , 2001 , 26, 28-40	2.7	53
62	Herbicide Metabolism and Cross-Tolerance in Transgenic Potato Plants Co-Expressing Human CYP1A1, CYP2B6, and CYP2C19. <i>Pesticide Biochemistry and Physiology</i> , 2000 , 66, 116-129	4.9	41
61	A major latex-like protein is a key factor in crop contamination by persistent organic pollutants. <i>Plant Physiology</i> , 2013 , 161, 2128-35	6.6	39
60	Transgenic rice containing human CYP2B6 detoxifies various classes of herbicides. <i>Journal of Agricultural and Food Chemistry</i> , 2005 , 53, 3461-7	5.7	38
59	Herbicide Metabolism and Cross-Tolerance in Transgenic Potato Plants Expressing Human CYP1A1. <i>Pesticide Biochemistry and Physiology</i> , 1999 , 64, 33-46	4.9	38
58	Enhanced herbicide cross-tolerance in transgenic rice plants co-expressing human CYP1A1, CYP2B6, and CYP2C19. <i>Plant Science</i> , 2005 , 168, 773-781	5.3	36
57	Structural basis of species differences between human and experimental animal CYP1A1s in metabolism of 3,3',4,4'-tetrachlorobiphenyl. <i>Journal of Biochemistry</i> , 2011 , 149, 487-94	3.1	34
56	Enhancement of metabolizing herbicides in young tubers of transgenic potato plants with the rat CYP1A1 gene. <i>Theoretical and Applied Genetics</i> , 2002 , 105, 515-520	6	29
55	Efficient biodegradation of petroleum -alkanes and polycyclic aromatic hydrocarbons by polyextremophilic <i>Saia</i> with multidegradative capacity.. <i>RSC Advances</i> , 2020 , 10, 14060-14070	3.7	28
54	Mammalian cytochrome P450-dependent metabolism of polychlorinated dibenzo-p-dioxins and coplanar polychlorinated biphenyls. <i>International Journal of Molecular Sciences</i> , 2014 , 15, 14044-57	6.3	28
53	Herbicide Metabolism and Tolerance in the Transgenic Rice Plants Expressing Human CYP2C9 and CYP2C19. <i>Pesticide Biochemistry and Physiology</i> , 2001 , 71, 156-169	4.9	28
52	Metabolism of the Herbicide Chlortoluron in Transgenic Tobacco Plants Expressing the Fused Enzyme between Rat Cytochrome P4501A1 and Yeast NADPH-Cytochrome P450 Oxidoreductase. <i>Pesticide Biochemistry and Physiology</i> , 1996 , 54, 190-198	4.9	26
51	Molecular Mechanisms of Herbicide Resistance with Special Emphasis on Cytochrome P450 Monooxygenases.. <i>Plant Biotechnology</i> , 1998 , 15, 173-176	1.3	25
50	Congener specificity in the accumulation of dioxins and dioxin-like compounds in zucchini plants grown hydroponically. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011 , 75, 705-10	2.1	24
49	Expression of human cytochromes P450 1A1 and P450 1A2 as fused enzymes with yeast NADPH-cytochrome P450 oxidoreductase in transgenic tobacco plants. <i>Bioscience, Biotechnology and Biochemistry</i> , 2000 , 64, 2025-33	2.1	24

48	Aryl hydrocarbon receptor (AhR)-mediated reporter gene expression systems in transgenic tobacco plants. <i>Planta</i> , 2007 , 227, 37-45	4.7	22
47	Metabolism of agrochemicals and related environmental chemicals based on cytochrome P450s in mammals and plants. <i>Pest Management Science</i> , 2015 , 71, 824-8	4.6	20
46	Structure-selective accumulation of polychlorinated biphenyls in <i>Cucurbita pepo</i> . <i>Journal of Pesticide Sciences</i> , 2011 , 36, 363-369	2.7	20
45	Inducible cross-tolerance to herbicides in transgenic potato plants with the rat CYP1A1 gene. <i>Theoretical and Applied Genetics</i> , 2002 , 104, 308-314	6	19
44	Hormonal regulation and effects of four environmental pollutants on vitellogenin gene transcription in the giant water bug, <i>Lethocerus deyrollei</i> (Hemiptera: Belostomatidae). <i>Journal of Insect Conservation</i> , 2011 , 15, 421-431	2.1	18
43	Distribution of perfluoroalkyl compounds in Osaka Bay and coastal waters of Western Japan. <i>Chemosphere</i> , 2017 , 170, 260-265	8.4	16
42	Phytomonitoring and phytoremediation of agrochemicals and related compounds based on recombinant cytochrome P450s and aryl hydrocarbon receptors (AhRs). <i>Journal of Agricultural and Food Chemistry</i> , 2011 , 59, 2870-5	5.7	16
41	Recombinant aryl hydrocarbon receptors for bioassay of aryl hydrocarbon receptor ligands in transgenic tobacco plants. <i>Plant Biotechnology Journal</i> , 2009 , 7, 119-28	11.6	16
40	A scFv antibody-based immunoaffinity chromatography column for clean-up of bisphenol A-contaminated water samples. <i>Journal of Agricultural and Food Chemistry</i> , 2009 , 57, 353-8	5.7	16
39	Molecular characterization of specifically active recombinant fused enzymes consisting of CYP3A4, NADPH-cytochrome P450 oxidoreductase, and cytochrome b5. <i>Biochemistry</i> , 2007 , 46, 10213-21	3.2	16
38	Overexpression of <i>Arabidopsis thaliana</i> LOV KELCH REPEAT PROTEIN 2 promotes tuberization in potato (<i>Solanum tuberosum</i> cv. May Queen). <i>FEBS Letters</i> , 2010 , 584, 2393-6	3.8	13
37	Uptake mechanisms of polychlorinated biphenyls in <i>Cucurbita pepo</i> via xylem sap containing major latex-like proteins. <i>Environmental and Experimental Botany</i> , 2019 , 162, 399-405	5.9	12
36	Defluorination of perfluoroalkyl acids is followed by production of monofluorinated fatty acids. <i>Science of the Total Environment</i> , 2018 , 636, 355-359	10.2	12
35	Zinc finger protein genes from <i>Cucurbita pepo</i> are promising tools for conferring non-Cucurbitaceae plants with ability to accumulate persistent organic pollutants. <i>Chemosphere</i> , 2015 , 123, 48-54	8.4	12
34	Factors regulating the differential uptake of persistent organic pollutants in cucurbits and non-cucurbits. <i>Journal of Plant Physiology</i> , 2020 , 245, 153094	3.6	12
33	From the Cover: Structural Determinants of the Position of 2,3,4,4',5-Pentachlorobiphenyl (CB118) Hydroxylation by Mammalian Cytochrome P450 Monooxygenases. <i>Toxicological Sciences</i> , 2016 , 152, 340-344	4.4	11
32	Molecular mechanisms of herbicide-inducible gene expression of tobacco CYP71AH11 metabolizing the herbicide chlorotoluron. <i>Pesticide Biochemistry and Physiology</i> , 2014 , 108, 49-57	4.9	11
31	Molecular insights into the role of a distal F240A mutation that alters CYP1A1 activity towards persistent organic pollutants. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2017 , 1861, 2852-2860	4	11

30	Different uptake pathways between hydrophilic and hydrophobic compounds in lateral roots of Cucurbita pepo. <i>Journal of Pesticide Sciences</i> , 2015 , 40, 99-105	2.7	10
29	Enzyme-linked immunosorbent assay with monoclonal and single-chain variable fragment antibodies selective to coplanar polychlorinated biphenyls. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 1605-12	5.7	8
28	Enhanced expression of CYP2C9 and tolerance to sulfonylurea herbicides in transgenic rice plants. <i>Plant Biotechnology</i> , 2005 , 22, 89-96	1.3	8
27	Assays of dioxins and dioxin-like compounds in actually contaminated soils using transgenic tobacco plants carrying a recombinant mouse aryl hydrocarbon receptor-mediated β glucuronidase reporter gene expression system. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2012 , 47, 599-607	2.2	7
26	Assays of PCB congeners and organochlorine insecticides with the transgenic Arabidopsis and tobacco plants carrying recombinant guinea pig AhR and GUS reporter genes. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2012 , 47, 599-607	2.2	7
25	Effects of biosurfactants on assays of PCB congeners in transgenic arabidopsis plants carrying a recombinant guinea pig AhR-mediated GUS reporter gene expression system. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2012 , 47, 599-607	2.2	6
24	A selectable marker using cytochrome P450 monooxygenases for Arabidopsis transformation. <i>Plant Biotechnology</i> , 2005 , 22, 281-286	1.3	6
23	Metabolic enhancement of 2,3,4,4',5-pentachlorobiphenyl (CB118) using cytochrome P450 monooxygenase isolated from soil bacterium under the presence of perfluorocarboxylic acids (PFCAs) and the structural basis of its metabolism. <i>Chemosphere</i> , 2018 , 210, 376-383	8.4	5
22	Review: Biological functions of major latex-like proteins in plants. <i>Plant Science</i> , 2021 , 306, 110856	5.3	5
21	Capillary zone electrophoresis determination of fluoride in seawater using transient isotachopheresis. <i>Analytical and Bioanalytical Chemistry</i> , 2018 , 410, 1825-1831	4.4	4
20	Recombinant human AhR-mediated GUS reporter gene assays for PCB congeners in transgenic tobacco plants in comparison with recombinant mouse and guinea pig AhRs. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2010 , 45, 741-9	2.2	4
19	Bioassay of estrogenic compounds in transgenic Arabidopsis plants carrying a recombinant human estrogen receptor gene and a GFP reporter gene. <i>Transgenic Research</i> , 2009 , 18, 899-909	3.3	4
18	High temperatures promote the uptake of hydrophobic pollutants by altered gene expression levels of major latex-like proteins. <i>Journal of Pesticide Sciences</i> , 2020 , 45, 75-80	2.7	4
17	Suppression of the genes responsible for transporting hydrophobic pollutants leads to the production of safer crops. <i>Science of the Total Environment</i> , 2020 , 741, 140439	10.2	4
16	Pesticide treatment reduces hydrophobic pollutant contamination in Cucurbita pepo through competitive binding to major latex-like proteins. <i>Environmental Pollution</i> , 2020 , 266, 115179	9.3	4
15	Cytochrome P450 Monooxygenases Metabolizing Herbicides. <i>Biotechnology and Biotechnological Equipment</i> , 1998 , 12, 17-22	1.6	3
14	Effect of amending soil with organic acids on perylene uptake into Cucurbita pepo. <i>Journal of Pesticide Sciences</i> , 2014 , 39, 162-164	2.7	3
13	Designed recombinant transcription factor with antibody-variable regions. <i>Analytical Chemistry</i> , 2009 , 81, 10162-6	7.8	2

12	Transport enhancement of hydrophobic pollutants by the expression of zucchini major latex-like protein genes in tobacco plants. <i>Journal of Plant Physiology</i> , 2021 , 263, 153464	3.6	2
11	Assays of dioxins and dioxin-like compounds in actually contaminated soils using transgenic tobacco plants carrying a recombinant mouse aryl hydrocarbon receptor-mediated β -glucuronidase reporter gene expression system. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2012 , 47, 233-9	2.2	1
10	Molecular analysis of specificity of anti-nonylphenol polyethoxylate single-chain antibody fragments by grafting and designed point mutations. <i>Molecular Immunology</i> , 2009 , 46, 3125-30	4.3	1
9	Herbicide Resistant Transgenic Plants Expressing Cytochrome P450 Monooxygenases Metabolizing Xenobiotics. <i>ACS Symposium Series</i> , 2000 , 116-126	0.4	1
8	Effects of Arabidopsis Ku80 deletion on the integration of the left border of T-DNA into plant chromosomal DNA via <i>Agrobacterium tumefaciens</i> . <i>Genes and Genetic Systems</i> , 2020 , 95, 173-182	1.4	1
7	MLP-PG1, a major latex-like protein identified in <i>Cucurbita pepo</i> , confers resistance through the induction of pathogenesis-related genes. <i>Planta</i> , 2021 , 255, 10	4.7	1
6	How does the Cucurbitaceae family take up organic pollutants (POPs, PAHs, and PPCPs)? <i>Reviews in Environmental Science and Biotechnology</i> , 2021 , 20, 751-779	13.9	1
5	Bioassay for Persistent Organic Pollutants in Transgenic Plants with Ah Receptor and GUS Reporter Genes 431-438		
4	Simple monitoring of endocrine-disrupting chemicals using transgenic Arabidopsis plants expressing medaka estrogen receptor. <i>Chemosphere</i> , 2022 , 286, 131633	8.4	0
3	Engineering of Transgenic Plants Expressing Drug-Metabolizing Enzymes for Reduction of Pesticide Residues. <i>Journal of Pesticide Sciences</i> , 2001 , 26, 318-326	2.7	
2	Herbicide Metabolism and Resistance of Transgenic Potato Plants Expressing Rat Cytochrome P4501A1. <i>Breeding Science</i> , 1998 , 48, 135-143		
1	Hydroxylation and dechlorination of 3,3',4,4'-tetrachlorobiphenyl (CB77) by rat and human CYP1A1s and critical roles of amino acids composing their substrate-binding cavity. <i>Science of the Total Environment</i> , 2022 , 155848	10.2	