

Cheryl E Rockwell

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2867798/cheryl-e-rockwell-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

41
papers

1,235
citations

20
h-index

35
g-index

45
ext. papers

1,415
ext. citations

4.5
avg, IF

4.36
L-index

#	Paper	IF	Citations
41	Trivalent arsenic impairs the effector response of human CD4 and CD8 T cells to influenza A virus ex vivo.. <i>Food and Chemical Toxicology</i> , 2022 , 113122	4.7	
40	Phenotypic Changes in T Cell and Macrophage Subtypes in Perivascular Adipose Tissues Precede High-Fat Diet-Induced Hypertension. <i>Frontiers in Physiology</i> , 2021 , 12, 616055	4.6	3
39	The role of Nrf2 in autoimmunity and infectious disease: Therapeutic possibilities. <i>Advances in Pharmacology</i> , 2021 , 91, 61-110	5.7	1
38	Nrf2 ^{hi} , Regulatory, Activated, and Memory Immune Cells Co-exist in PVATs That Are Comparable in Density to Non-PVAT Fats in Health. <i>Frontiers in Physiology</i> , 2020 , 11, 58	4.6	12
37	Nrf2-dependent and -independent effects of tBHQ in activated murine B cells. <i>Food and Chemical Toxicology</i> , 2020 , 145, 111595	4.7	4
36	Chronic low-level cadmium exposure in rats affects cytokine production by activated T cells. <i>Toxicology Research</i> , 2019 , 8, 227-237	2.6	11
35	Dichotomous Role of Plasmin in Regulation of Macrophage Function after Acetaminophen Overdose. <i>American Journal of Pathology</i> , 2019 , 189, 1986-2001	5.8	2
34	Differential Sensitivity of Kupffer Cells and Hepatic Monocyte-Derived Macrophages to Bacterial Lipopolysaccharide 2019 , 1,		5
33	The Immune Response to Influenza is Suppressed by the Synthetic Food Additive and Nrf2 Activator, tert-butylhydroquinone (tBHQ). <i>FASEB Journal</i> , 2019 , 33, 505.3	0.9	1
32	Identification of an Unfavorable Immune Signature in Advanced Lung Tumors from Nrf2-Deficient Mice. <i>Antioxidants and Redox Signaling</i> , 2018 , 29, 1535-1552	8.4	20
31	Comparison of Hepatic NRF2 and Aryl Hydrocarbon Receptor Binding in 2,3,7,8-Tetrachlorodibenzo-p-dioxin-Treated Mice Demonstrates NRF2-Independent PKM2 Induction. <i>Molecular Pharmacology</i> , 2018 , 94, 876-884	4.3	9
30	Differential effects of the Nrf2 activators tBHQ and CDDO-Im on the early events of T cell activation. <i>Biochemical Pharmacology</i> , 2018 , 147, 67-76	6	20
29	The Nrf2 activator tBHQ inhibits the activation of primary murine natural killer cells. <i>Food and Chemical Toxicology</i> , 2018 , 121, 231-236	4.7	9
28	Nrf2-Dependent and -Independent Effects of t-Butylhydroquinone, CDDO-Im, and HO in Human Jurkat T Cells as Determined by CRISPR/Cas9 Gene Editing. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2017 , 361, 259-267	4.7	12
27	Persistent alterations in immune cell populations and function from a single dose of perfluorononanoic acid (PFNA) in C57Bl/6 mice. <i>Food and Chemical Toxicology</i> , 2017 , 100, 24-33	4.7	11
26	Pharmacological Inhibition of Myocardin-related Transcription Factor Pathway Blocks Lung Metastases of RhoC-Overexpressing Melanoma. <i>Molecular Cancer Therapeutics</i> , 2017 , 16, 193-204	6.1	24
25	Inhibition of early T cell cytokine production by arsenic trioxide occurs independently of Nrf2. <i>PLoS ONE</i> , 2017 , 12, e0185579	3.7	12

24	Fibrin deposition following bile duct injury limits fibrosis through an $\text{M}\alpha$ -dependent mechanism. <i>Blood</i> , 2016 , 127, 2751-62	2.2	23
23	The Nrf2 activator tBHQ inhibits T cell activation of primary human CD4 T cells. <i>Cytokine</i> , 2015 , 71, 289-95	4.6	38
22	Individual bile acids have differential effects on bile acid signaling in mice. <i>Toxicology and Applied Pharmacology</i> , 2015 , 283, 57-64	4.6	54
21	Determination of the Effects of Nrf2 upon the Early Events of Jurkat T Cell Activation by Use of CRISPR-CAS9 Mediated Mutation. <i>FASEB Journal</i> , 2015 , 29, 621.10	0.9	
20	Hepatic stellate cells orchestrate clearance of necrotic cells in a hypoxia-inducible factor-1 β -dependent manner by modulating macrophage phenotype in mice. <i>Journal of Immunology</i> , 2014 , 192, 3847-3857	5.3	33
19	Fas-induced apoptosis increases hepatocyte tissue factor procoagulant activity in vitro and in vivo. <i>Toxicological Sciences</i> , 2014 , 141, 453-64	4.4	12
18	IL-17A synergistically enhances bile acid-induced inflammation during obstructive cholestasis. <i>American Journal of Pathology</i> , 2013 , 183, 1498-1507	5.8	51
17	The Nrf2 activator, tBHQ, differentially affects early events following stimulation of Jurkat cells. <i>Toxicological Sciences</i> , 2013 , 136, 63-71	4.4	22
16	Hepatocyte tissue factor activates the coagulation cascade in mice. <i>Blood</i> , 2013 , 121, 1868-74	2.2	50
15	Acute Immunotoxic Effects of Perfluorononanoic Acid (PFNA) in C57BL/6 Mice. <i>Clinical & Experimental Pharmacology</i> , 2013 , Suppl 4,	1	5
14	Th2 skewing by activation of Nrf2 in CD4(+) T cells. <i>Journal of Immunology</i> , 2012 , 188, 1630-7	5.3	80
13	Effect of bile duct ligation on bile acid composition in mouse serum and liver. <i>Liver International</i> , 2012 , 32, 58-69	7.9	119
12	A critical role for the inducible proteasomal subunits LMP7 and MECL1 in cytokine production by activated murine splenocytes. <i>Pharmacology</i> , 2012 , 89, 117-26	2.3	10
11	Protease-activated receptor 1 and hematopoietic cell tissue factor are required for hepatic steatosis in mice fed a Western diet. <i>American Journal of Pathology</i> , 2011 , 179, 2278-89	5.8	30
10	ChIPing the cistrome of PXR in mouse liver. <i>Nucleic Acids Research</i> , 2010 , 38, 7943-63	20.1	54
9	Transcriptional regulation of renal cytoprotective genes by Nrf2 and its potential use as a therapeutic target to mitigate cisplatin-induced nephrotoxicity. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2010 , 335, 2-12	4.7	138
8	Differential effects of lactacystin on cytokine production in activated Jurkat cells and murine splenocytes. <i>Cytokine</i> , 2010 , 51, 12-7	4	6
7	Lipid A-mediated tolerance and cancer therapy. <i>Advances in Experimental Medicine and Biology</i> , 2010 , 667, 81-99	3.6	7

6	A COX-2 metabolite of the endogenous cannabinoid, 2-arachidonyl glycerol, mediates suppression of IL-2 secretion in activated Jurkat T cells. <i>Biochemical Pharmacology</i> , 2008 , 76, 353-61	6	34
5	A combination of proteasome inhibitors and antibiotics prevents lethality in a septic shock model. <i>Innate Immunity</i> , 2008 , 14, 319-29	2.7	21
4	Interleukin-2 suppression by 2-arachidonyl glycerol is mediated through peroxisome proliferator-activated receptor gamma independently of cannabinoid receptors 1 and 2. <i>Molecular Pharmacology</i> , 2006 , 70, 101-11	4.3	132
3	2-Arachidonoyl-glycerol suppresses interferon-gamma production in phorbol ester/ionomycin-activated mouse splenocytes independent of CB1 or CB2. <i>Journal of Leukocyte Biology</i> , 2005 , 77, 966-74	6.5	20
2	A cyclooxygenase metabolite of anandamide causes inhibition of interleukin-2 secretion in murine splenocytes. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004 , 311, 683-90	4.7	74
1	Evidence for cannabinoid receptor-dependent and -independent mechanisms of action in leukocytes. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003 , 306, 1077-85	4.7	65