

Agnieszka Hernik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/287214/publications.pdf>

Version: 2024-02-01

17
papers

515
citations

932766

10
h-index

676716

22
g-index

36
all docs

36
docs citations

36
times ranked

966
citing authors

#	ARTICLE	IF	CITATIONS
1	Biopesticides - towards increased consumer safety in the European Union. <i>Pest Management Science</i> , 2015, 71, 3-6.	1.7	124
2	Pregnancy serum concentrations of perfluorinated alkyl substances and offspring behaviour and motor development at age 5-9 years - a prospective study. <i>Environmental Health</i> , 2015, 14, 2.	1.7	65
3	Hazard quotient profiles used as a risk assessment tool for PFOS and PFOA serum levels in three distinctive European populations. <i>Environment International</i> , 2015, 74, 112-118.	4.8	61
4	No association between body mass index and sperm DNA integrity. <i>Human Reproduction</i> , 2015, 30, 1704-1713.	0.4	50
5	Perfluorinated chemicals in blood serum of inhabitants in central Poland in relation to gender and age. <i>Science of the Total Environment</i> , 2015, 532, 548-555.	3.9	44
6	Levels of polybrominated diphenyl ethers in house dust in Central Poland. <i>Indoor Air</i> , 2017, 27, 128-135.	2.0	21
7	PCDD/Fs and DL-PCBs intake from fish caught in Polish fishing grounds in the Baltic Sea - Characterizing the risk for consumers. <i>Environment International</i> , 2013, 56, 32-41.	4.8	19
8	Characterising the individual health risk in infants exposed to organochlorine pesticides via breast milk by applying appropriate margins of safety derived from estimated daily intakes. <i>Chemosphere</i> , 2014, 94, 158-163.	4.2	14
9	Polybrominated diphenyl ethers, polychlorinated biphenyls and organochlorine pesticides in human milk as markers of environmental exposure to these compounds. <i>Annals of Agricultural and Environmental Medicine</i> , 2011, 18, 113-8.	0.5	13
10	Polybrominated diphenyl ethers and polychlorinated biphenyls in cord blood from women in Poland. <i>Chemosphere</i> , 2013, 93, 526-531.	4.2	12
11	Is the fact of parenting couples cohabitation affecting the serum levels of persistent organohalogen pollutants?. <i>International Journal of Hygiene and Environmental Health</i> , 2015, 218, 392-400.	2.1	7
12	Alternative toxicological methods for establishing residue definitions applied for dietary risk assessment of pesticides in the European Union. <i>Food and Chemical Toxicology</i> , 2020, 137, 111120.	1.8	7
13	Consumer Risk Assessment Associated with Intake of Pesticide Residues in Food of Plant Origin from the Retail Market in Poland. <i>Human and Ecological Risk Assessment (HERA)</i> , 2015, 21, 2036-2061.	1.7	5
14	Relationship between paired cord blood and milk POPs levels as a tool for assessing perinatal exposure, a pilot study. <i>Human and Ecological Risk Assessment (HERA)</i> , 2016, 22, 1456-1468.	1.7	5
15	Risk assessment for pesticides' MRL non-compliances in Poland in the years 2011-2015. <i>Roczniki Panstwowego Zakladu Higieny</i> , 2015, 66, 309-17.	0.5	2
16	Validation of the analytical method for the simultaneous determination of selected polybrominated diphenyl ethers, polychlorinated biphenyls and organochlorine pesticides in human blood serum by gas chromatography with microelectron capture detector. <i>Roczniki Panstwowego Zakladu Higieny</i> , 2016, 67, 113-20.	0.5	1
17	Different risk assessment methodologies applied for infant's exposure for polybrominated diphenyl ethers: Implications for public health. <i>Human and Ecological Risk Assessment (HERA)</i> , 2021, 27, 1954-1964.	1.7	0