

Agnieszka Hernik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/287214/publications.pdf>

Version: 2024-02-01

17
papers

515
citations

933447

10
h-index

677142

22
g-index

36
all docs

36
docs citations

36
times ranked

966
citing authors

#	ARTICLE	IF	CITATIONS
1	Different risk assessment methodologies applied for infant's exposure for polybrominated diphenyl ethers: Implications for public health. Human and Ecological Risk Assessment (HERA), 2021, 27, 1954-1964.	3.4	0
2	Alternative toxicological methods for establishing residue definitions applied for dietary risk assessment of pesticides in the European Union. Food and Chemical Toxicology, 2020, 137, 111120.	3.6	7
3	Levels of polybrominated diphenyl ethers in house dust in Central Poland. Indoor Air, 2017, 27, 128-135.	4.3	21
4	Relationship between paired cord blood and milk POPs levels as a tool for assessing perinatal exposure, a pilot study. Human and Ecological Risk Assessment (HERA), 2016, 22, 1456-1468.	3.4	5
5	Validation of the analytical method for the simultaneous determination of selected polybrominated diphenyl ethers, polychlorinated biphenyls and organochlorine pesticides in human blood serum by gas chromatography with microelectron capture detector. Roczniki Panstwowego Zakladu Higieny, 2016, 67, 113-20.	0.7	1
6	No association between body mass index and sperm DNA integrity. Human Reproduction, 2015, 30, 1704-1713.	0.9	50
7	Pregnancy serum concentrations of perfluorinated alkyl substances and offspring behaviour and motor development at age 5-9 years - a prospective study. Environmental Health, 2015, 14, 2.	4.0	65
8	Is the fact of parenting couples cohabitation affecting the serum levels of persistent organohalogen pollutants?. International Journal of Hygiene and Environmental Health, 2015, 218, 392-400.	4.3	7
9	Perfluorinated chemicals in blood serum of inhabitants in central Poland in relation to gender and age. Science of the Total Environment, 2015, 532, 548-555.	8.0	44
10	Consumer Risk Assessment Associated with Intake of Pesticide Residues in Food of Plant Origin from the Retail Market in Poland. Human and Ecological Risk Assessment (HERA), 2015, 21, 2036-2061.	3.4	5
11	Biopesticides - towards increased consumer safety in the European Union. Pest Management Science, 2015, 71, 3-6.	3.4	124
12	Hazard quotient profiles used as a risk assessment tool for PFOS and PFOA serum levels in three distinctive European populations. Environment International, 2015, 74, 112-118.	10.0	61
13	Risk assessment for pesticides' MRL non-compliances in Poland in the years 2011-2015. Roczniki Panstwowego Zakladu Higieny, 2015, 66, 309-17.	0.7	2
14	Characterising the individual health risk in infants exposed to organochlorine pesticides via breast milk by applying appropriate margins of safety derived from estimated daily intakes. Chemosphere, 2014, 94, 158-163.	8.2	14
15	Polybrominated diphenyl ethers and polychlorinated biphenyls in cord blood from women in Poland. Chemosphere, 2013, 93, 526-531.	8.2	12
16	PCDD/Fs and DL-PCBs intake from fish caught in Polish fishing grounds in the Baltic Sea - Characterizing the risk for consumers. Environment International, 2013, 56, 32-41.	10.0	19
17	Polybrominated diphenyl ethers, polychlorinated biphenyls and organochlorine pesticides in human milk as markers of environmental exposure to these compounds. Annals of Agricultural and Environmental Medicine, 2011, 18, 113-8.	1.0	13