
## **Xixiong Kang**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2872157/publications.pdf Version: 2024-02-01



XIXIONC KANC

#	Article	IF	CITATIONS
1	The diagnostic value of cerebrospinal fluids procalcitonin and lactate for the differential diagnosis of post-neurosurgical bacterial meningitis and aseptic meningitis. Clinical Biochemistry, 2015, 48, 50-54.	1.9	57
2	Chemiluminescence immunoassay based on microfluidic chips for α-fetoprotein. Clinica Chimica Acta, 2014, 431, 113-117.	1.1	44
3	High expression of UBE2C is associated with the aggressive progression and poor outcome of malignant glioma. Oncology Letters, 2016, 11, 2300-2304.	1.8	42
4	Identification and functional characterization of IncRNAs acting as ceRNA involved in the malignant progression of glioblastoma multiforme. Oncology Reports, 2016, 36, 2911-2925.	2.6	34
5	Toll-like receptor 2 and -4 are involved in the pathogenesis of the Guillain-Barré syndrome. Molecular Medicine Reports, 2015, 12, 3207-3213.	2.4	20
6	Labelâ€free serum detection based on Raman spectroscopy for the diagnosis and classification of glioma. Journal of Raman Spectroscopy, 2020, 51, 1977-1985.	2.5	18
7	Rapid Detection of mecA and femA Genes by Loop-Mediated Isothermal Amplification in a Microfluidic System for Discrimination of Different Staphylococcal Species and Prediction of Methicillin Resistance. Frontiers in Microbiology, 2020, 11, 1487.	3.5	15
8	The Clinical Significance of Soluble Programmed Cell Death-Ligand 1 (sPD-L1) in Patients With Gliomas. Frontiers in Oncology, 2020, 10, 9.	2.8	14
9	Association of SLCO1B1 and ABCB1 Genetic Variants with Atorvastatin-induced Myopathy in Patients with Acute Ischemic Stroke. Current Pharmaceutical Design, 2019, 25, 1663-1670.	1.9	11
10	The combination of cerebrospinal fluid procalcitonin, lactate, interleukin-8 and interleukin-10 concentrations for the diagnosis of postneurosurgical bacterial meningitis: A prospective study. Annals of Clinical Biochemistry, 2019, 56, 133-140.	1.6	10
11	Soluble cytotoxic T-lymphocyte–associated antigen 4 (sCTLA-4) as a potential biomarker for diagnosis and evaluation of the prognosis in Glioma. BMC Immunology, 2021, 22, 33.	2.2	9
12	Higher numbers of circulating endothelial progenitor cells in stroke patients with intracranial arterial stenosis. BMC Neurology, 2013, 13, 161.	1.8	8
13	Association between acromegaly and a single nucleotide polymorphism (rs2854744) in the IGFBP3 gene. BMC Medical Genetics, 2018, 19, 182.	2.1	8
14	A pilot study of cdc6 as a biomarker for circulating tumor cells in patients with lung cancer. Journal of Clinical Laboratory Analysis, 2020, 34, e23245.	2.1	7
15	Risk Factors of Recurrent Ischemic Events after Acute Noncardiogenic Ischemic Stroke. Current Pharmaceutical Design, 2020, 25, 4827-4834.	1.9	7
16	EMT-Related Markers in Serum Exosomes are Potential Diagnostic Biomarkers for Invasive Pituitary Adenomas. Neuropsychiatric Disease and Treatment, 2021, Volume 17, 3769-3780.	2.2	6
17	DHX32 expression is an indicator of poor breast cancer prognosis. Oncology Letters, 2017, 13, 942-948.	1.8	5
18	A rapid procedure for bacterial identification and antimicrobial susceptibility testing directly from positive blood cultures. Analyst, The, 2021, 147, 147-154.	3.5	5

XIXIONG KANG

#	Article	IF	CITATIONS
19	Seizure Control of Current Shunt on Rats with Temporal Lobe Epilepsy and Neocortical Epilepsy. PLoS ONE, 2014, 9, e86477.	2.5	2
20	Soluble programmed death-1 (sPD-1) and programmed death ligand 1 (sPD-L1) as potential biomarkers for the diagnosis and prognosis of glioma patients. Journal of Medical Biochemistry, 2020, 39, 444-451.	1.7	2
21	Influence of the IGFBP3-202A/C Gene Polymorphism on Clinical Features and Surgery Outcome in Acromegalic Patients. Frontiers in Endocrinology, 2018, 9, 751.	3.5	1
22	Inhibitory Effect of Tetramerized Single-Chain Variable Fragment of Anti-Cyclic Citrullinated Peptide Antibodies on the Proliferation, Activation, and Secretion of Cytokines of Fibroblast-Like Synoviocytes in Rheumatoid Arthritis In Vitro Co-Culture System. Inflammation, 2020, 43, 2245-2255.	3.8	0