Elizabeth K Speliotes

List of Publications by Citations

Source: https://exaly.com/author-pdf/2875006/elizabeth-k-speliotes-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 80
 22,469
 44
 90

 papers
 citations
 h-index
 g-index

 90
 26,223
 15.4
 4.96

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
80	Genetic studies of body mass index yield new insights for obesity biology. <i>Nature</i> , 2015 , 518, 197-206	50.4	2687
79	Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. <i>Science</i> , 2007 , 316, 1331-6	33.3	2364
78	Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. <i>Nature Genetics</i> , 2010 , 42, 937-48	36.3	2267
77	Hundreds of variants clustered in genomic loci and biological pathways affect human height. <i>Nature</i> , 2010 , 467, 832-8	50.4	1514
76	Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. <i>Nature Genetics</i> , 2009 , 41, 25-34	36.3	1368
75	Defining the role of common variation in the genomic and biological architecture of adult human height. <i>Nature Genetics</i> , 2014 , 46, 1173-86	36.3	1339
74	Common variants near MC4R are associated with fat mass, weight and risk of obesity. <i>Nature Genetics</i> , 2008 , 40, 768-75	36.3	1048
73	New genetic loci link adipose and insulin biology to body fat distribution. <i>Nature</i> , 2015 , 518, 187-196	50.4	920
7 2	Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. <i>Nature Genetics</i> , 2010 , 42, 949-60	36.3	724
71	Loss-of-function mutations in APOC3, triglycerides, and coronary disease. <i>New England Journal of Medicine</i> , 2014 , 371, 22-31	59.2	721
70	Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. <i>PLoS Genetics</i> , 2011 , 7, e1001324	6	629
69	Biological interpretation of genome-wide association studies using predicted gene functions. <i>Nature Communications</i> , 2015 , 6, 5890	17.4	489
68	Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. <i>Nature Genetics</i> , 2013 , 45, 501-12	36.3	437
67	Genome-wide association study identifies loci influencing concentrations of liver enzymes in plasma. <i>Nature Genetics</i> , 2011 , 43, 1131-8	36.3	415
66	Rare and low-frequency coding variants alter human adult height. <i>Nature</i> , 2017 , 542, 186-190	50.4	412
65	Genome-wide association scan meta-analysis identifies three Loci influencing adiposity and fat distribution. <i>PLoS Genetics</i> , 2009 , 5, e1000508	6	393
64	Exome-wide association study of plasma lipids in >300,000 individuals. <i>Nature Genetics</i> , 2017 , 49, 1758-	·1 36. 6	310

63	FTO genotype is associated with phenotypic variability of body mass index. <i>Nature</i> , 2012 , 490, 267-72	50.4	304
62	Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. <i>PLoS Genetics</i> , 2013 , 9, e1003500	6	277
61	Fatty liver is associated with dyslipidemia and dysglycemia independent of visceral fat: the Framingham Heart Study. <i>Hepatology</i> , 2010 , 51, 1979-87	11.2	277
60	PNPLA3 variants specifically confer increased risk for histologic nonalcoholic fatty liver disease but not metabolic disease. <i>Hepatology</i> , 2010 , 52, 904-12	11.2	267
59	Genetic variation near IRS1 associates with reduced adiposity and an impaired metabolic profile. <i>Nature Genetics</i> , 2011 , 43, 753-60	36.3	237
58	Association Between Telomere Length and Risk of Cancer and Non-Neoplastic Diseases: A Mendelian Randomization Study. <i>JAMA Oncology</i> , 2017 , 3, 636-651	13.4	236
57	The survivin-like C. elegans BIR-1 protein acts with the Aurora-like kinase AIR-2 to affect chromosomes and the spindle midzone. <i>Molecular Cell</i> , 2000 , 6, 211-23	17.6	223
56	A meta-analysis identifies new loci associated with body mass index in individuals of African ancestry. <i>Nature Genetics</i> , 2013 , 45, 690-6	36.3	192
55	Population genetic differentiation of height and body mass index across Europe. <i>Nature Genetics</i> , 2015 , 47, 1357-62	36.3	186
54	Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. <i>Nature Genetics</i> , 2018 , 50, 26-41	36.3	186
53	Low-frequency and rare exome chip variants associate with fasting glucose and type 2 diabetes susceptibility. <i>Nature Communications</i> , 2015 , 6, 5897	17.4	147
52	17-Beta Hydroxysteroid Dehydrogenase 13IIs a Hepatic Retinol Dehydrogenase Associated With Histological Features of Nonalcoholic Fatty Liver Disease. <i>Hepatology</i> , 2019 , 69, 1504-1519	11.2	133
51	Genetic evidence for a normal-weight "metabolically obese" phenotype linking insulin resistance, hypertension, coronary artery disease, and type 2 diabetes. <i>Diabetes</i> , 2014 , 63, 4369-77	0.9	131
50	Hepatic steatosis and cardiovascular disease outcomes: An analysis of the Framingham Heart Study. Journal of Hepatology, 2015 , 63, 470-6	13.4	119
49	Sugar-sweetened beverage, diet soda, and fatty liver disease in the Framingham Heart Study cohorts. <i>Journal of Hepatology</i> , 2015 , 63, 462-9	13.4	112
48	Association between variants in or near PNPLA3, GCKR, and PPP1R3B with ultrasound-defined steatosis based on data from the third National Health and Nutrition Examination Survey. <i>Clinical Gastroenterology and Hepatology</i> , 2013 , 11, 1183-1190.e2	6.9	95
47	Liver fat is reproducibly measured using computed tomography in the Framingham Heart Study. Journal of Gastroenterology and Hepatology (Australia), 2008 , 23, 894-9	4	94
46	Genome-wide association of body fat distribution in African ancestry populations suggests new loci. <i>PLoS Genetics</i> , 2013 , 9, e1003681	6	92

45	Characterization of European ancestry nonalcoholic fatty liver disease-associated variants in individuals of African and Hispanic descent. <i>Hepatology</i> , 2013 , 58, 966-75	11.2	91
44	Common body mass index-associated variants confer risk of extreme obesity. <i>Human Molecular Genetics</i> , 2009 , 18, 3502-7	5.6	91
43	Increased expression of basic fibroblast growth factor (bFGF) following focal cerebral infarction in the rat. <i>Molecular Brain Research</i> , 1996 , 39, 31-42		77
42	Intramuscular fat and associations with metabolic risk factors in the Framingham Heart Study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 863-70	9.4	69
41	Genetic variation at NCAN locus is associated with inflammation and fibrosis in non-alcoholic fatty liver disease in morbid obesity. <i>Human Heredity</i> , 2013 , 75, 34-43	1.1	66
40	TM6SF2: catch-22 in the fight against nonalcoholic fatty liver disease and cardiovascular disease?. <i>Gastroenterology</i> , 2015 , 148, 679-84	13.3	58
39	Association of Nonalcoholic Fatty Liver Disease With Lower Brain Volume in Healthy Middle-aged Adults in the Framingham Study. <i>JAMA Neurology</i> , 2018 , 75, 97-104	17.2	54
38	Nonalcoholic fatty liver disease and vascular function: cross-sectional analysis in the Framingham heart study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015 , 35, 1284-91	9.4	53
37	Genome-wide association analyses identify 39 new susceptibility loci for diverticular disease. <i>Nature Genetics</i> , 2018 , 50, 1359-1365	36.3	49
36	Gene-based meta-analysis of genome-wide association studies implicates new loci involved in obesity. <i>Human Molecular Genetics</i> , 2015 , 24, 6849-60	5.6	44
35	Protein-coding variants implicate novel genes related to lipid homeostasis contributing to body-fat distribution. <i>Nature Genetics</i> , 2019 , 51, 452-469	36.3	44
34	Insights from Genome-Wide Association Analyses of Nonalcoholic Fatty Liver Disease. <i>Seminars in Liver Disease</i> , 2015 , 35, 375-91	7.3	37
33	Novel association of rs58542926 genotype with increased serum tyrosine levels and decreased apoB-100 particles in Finns. <i>Journal of Lipid Research</i> , 2017 , 58, 1471-1481	6.3	35
32	A Peripheral Blood DNA Methylation Signature of Hepatic Fat Reveals a Potential Causal Pathway for Nonalcoholic Fatty Liver Disease. <i>Diabetes</i> , 2019 , 68, 1073-1083	0.9	25
31	Body mass index trajectories in young adulthood predict non-alcoholic fatty liver disease in middle age: The CARDIA cohort study. <i>Liver International</i> , 2018 , 38, 706-714	7.9	24
30	rs641738C>T near MBOAT7 is associated with liver fat, ALT and fibrosis in NAFLD: A meta-analysis. <i>Journal of Hepatology</i> , 2021 , 74, 20-30	13.4	24
29	Adipose Tissue Depots and Their Cross-Sectional Associations With Circulating Biomarkers of Metabolic Regulation. <i>Journal of the American Heart Association</i> , 2016 , 5,	6	23
28	Genetic variants that associate with cirrhosis have pleiotropic effects on human traits. <i>Liver International</i> , 2020 , 40, 405-415	7.9	22

(2017-2019)

27	Insulin Resistance Exacerbates Genetic Predisposition to Nonalcoholic Fatty Liver Disease in Individuals Without Diabetes. <i>Hepatology Communications</i> , 2019 , 3, 894-907	6	21
26	Recent Advances in Human Genetics and Epigenetics of Adiposity: Pathway to Precision Medicine?. <i>Gastroenterology</i> , 2017 , 152, 1695-1706	13.3	20
25	Identification of seven novel loci associated with amino acid levels using single-variant and gene-based tests in 8545 Finnish men from the METSIM study. <i>Human Molecular Genetics</i> , 2018 , 27, 16	64 ⁵ 167	4 ²⁰
24	A Comprehensive Analysis of Common and Rare Variants to Identify Adiposity Loci in Hispanic Americans: The IRAS Family Study (IRASFS). <i>PLoS ONE</i> , 2015 , 10, e0134649	3.7	17
23	Comparison of the potency of competitive NMDA antagonists against the neurotoxicity of glutamate and NMDA. <i>Journal of Neurochemistry</i> , 1994 , 63, 879-85	6	16
22	Treatment of Dyslipidemia in Common Liver Diseases. <i>Clinical Gastroenterology and Hepatology</i> , 2018 , 16, 1189-1196	6.9	16
21	Independent markers of nonalcoholic fatty liver disease in a gentrifying population-based Chinese cohort. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3156	7.5	15
20	Genome-wide family-based linkage analysis of exome chip variants and cardiometabolic risk. <i>Genetic Epidemiology</i> , 2014 , 38, 345-52	2.6	14
19	Genome-wide association study of serum liver enzymes implicates diverse metabolic and liver pathology. <i>Nature Communications</i> , 2021 , 12, 816	17.4	14
18	Twenty-five-year trajectories of insulin resistance and pancreatic Eell response and diabetes risk in nonalcoholic fatty liver disease. <i>Liver International</i> , 2018 , 38, 2069-2081	7.9	10
17	Body Composition and Genetic Lipodystrophy Risk Score Associate With Nonalcoholic Fatty Liver Disease and Liver Fibrosis. <i>Hepatology Communications</i> , 2019 , 3, 1073-1084	6	9
16	Genome-Wide Study of Subcutaneous and Visceral Adipose Tissue Reveals Novel Sex-Specific Adiposity Loci in Mexican Americans. <i>Obesity</i> , 2018 , 26, 202-212	8	9
15	Discovery and fine-mapping of height loci via high-density imputation of GWASs in individuals of African ancestry. <i>American Journal of Human Genetics</i> , 2021 , 108, 564-582	11	7
14	Empirical characteristics of family-based linkage to a complex trait: the ADIPOQ region and adiponectin levels. <i>Human Genetics</i> , 2015 , 134, 203-13	6.3	6
13	Allele-specific variation at APOE increases nonalcoholic fatty liver disease and obesity but decreases risk of Alzheimer@ disease and myocardial infarction. <i>Human Molecular Genetics</i> , 2021 , 30, 1443-1456	5.6	5
12	Loci identified by a genome-wide association study of carotid artery stenosis in the eMERGE network. <i>Genetic Epidemiology</i> , 2021 , 45, 4-15	2.6	5
11	Functional Analysis of the Dengue Virus Genome Using an Insertional Mutagenesis Screen. <i>Journal of Virology</i> , 2018 , 92,	6.6	4
10	Genome-wide linkage and association analysis of cardiometabolic phenotypes in Hispanic Americans. <i>Journal of Human Genetics</i> , 2017 , 62, 175-184	4.3	4

9	rs641738C>T near MBOAT7 is positively associated with liver fat, ALT, and histological severity of NAFLD: a meta-analysis		3
8	A Noncoding Variant Near PPP1R3B Promotes Liver Glycogen Storage and MetS, but Protects Against Myocardial Infarction. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, 372-387	5.6	3
7	The genetic determinants of common human obesity. Current Cardiovascular Risk Reports, 2009, 3, 411	-41.79	2
6	Knockout of murine Lyplal1 confers sex-specific protection against diet-induced obesity		1
5	Treatment of Dyslipidemia in Common Liver Diseases. Clinical Liver Disease, 2019, 14, 161-162	2.2	O
4	Hepatic decompensation is accelerated in patients with cirrhosis and alpha-1 antitrypsin Pi*MZ genotype <i>JHEP Reports</i> , 2022 , 4, 100483	10.3	O
3	Thwart your destiny; effect of nonacoholic fatty liver disease genes on steatosis, liver injury and cirrhosis varies by body mass index. <i>Hepatology</i> , 2018 , 68, 372-374	11.2	
2	Commentary on Diagnostic Problems in Hepatology Cases. Seminars in Liver Disease, 2015 , 35, 432-3	7.3	

Genetic Pleiotropies of Obesity **2014**, 93-111