
## Heiko Sahling

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2884905/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Oil and gas seepage offshore Georgia (Black Sea) – Geochemical evidences for a paleogene-neogene hydrocarbon source rock. Marine and Petroleum Geology, 2021, 128, 104995.	3.3	8
2	Methane gas emissions of the Black Sea—mapping from the Crimean continental margin to the Kerch Peninsula slope. Geo-Marine Letters, 2020, 40, 467-480.	1.1	13
3	<strong>A new genus <em>Turneroconcha</em> (Bivalvia: Vesicomyidae: Pliocardiinae) for the giant hydrothermal vent clam â€~<em>Calyptogena</em>' <em>magnifica</em></strong> . Zootaxa, 2020, 4808, 79-100.	0.5	2
4	Anaerobic Degradation of Non-Methane Alkanes by " <i>Candidatus</i> Methanoliparia―in Hydrocarbon Seeps of the Gulf of Mexico. MBio, 2019, 10, .	4.1	63
5	Characteristics and hydrocarbon seepage at the Challenger Knoll in the Sigsbee Basin, Gulf of Mexico. Geo-Marine Letters, 2019, 39, 391-399.	1.1	4
6	Fueled by methane: deep-sea sponges from asphalt seeps gain their nutrition from methane-oxidizing symbionts. ISME Journal, 2019, 13, 1209-1225.	9.8	68
7	Amount and Fate of Gas and Oil Discharged at 3400 m Water Depth From a Natural Seep Site in the Southern Gulf of Mexico. Frontiers in Marine Science, 2019, 6, .	2.5	29
8	Morphology and activity of the Helgoland Mud Volcano in the Sorokin Trough, northern Black Sea. Marine and Petroleum Geology, 2019, 99, 227-236.	3.3	8
9	Mud Volcanism in a Canyon: Morphodynamic Evolution of the Active Venere Mud Volcano and Its Interplay With Squillace Canyon, Central Mediterranean. Geochemistry, Geophysics, Geosystems, 2018, 19, 356-378.	2.5	12
10	Resolving the status of the families Vesicomyidae and Kelliellidae (Bivalvia: Venerida), with notes on their ecology. Journal of Molluscan Studies, 2018, 84, 69-91.	1.2	6
11	Seafloor sealing, doming, and collapse associated with gas seeps and authigenic carbonate structures at Venere mud volcano, Central Mediterranean. Deep-Sea Research Part I: Oceanographic Research Papers, 2018, 137, 76-96.	1.4	31
12	Slow Volcanoes: The Intriguing Similarities Between Marine Asphalt and Basalt Lavas. Oceanography, 2018, 31, .	1.0	10
13	Short-chain alkanes fuel mussel and sponge Cycloclasticus symbionts from deep-sea gas and oil seeps. Nature Microbiology, 2017, 2, 17093.	13.3	80
14	Phylogeny and origins of chemosynthetic vesicomyid clams. Systematics and Biodiversity, 2017, 15, 346-360.	1.2	53
15	Massive asphalt deposits, oil seepage, and gas venting support abundant chemosynthetic communities at the Campeche Knolls, southern Gulf of Mexico. Biogeosciences, 2016, 13, 4491-4512.	3.3	40
16	Fluxes and fate of dissolved methane released at the seafloor at the landward limit of the gas hydrate stability zone offshore western Svalbard. Journal of Geophysical Research: Oceans, 2015, 120, 6185-6201.	2.6	57
17	Seasonal methane accumulation and release from a gas emission site in the central North Sea. Biogeosciences, 2015, 12, 5261-5276.	3.3	32
18	Gas emissions at the continental margin west of Svalbard: mapping, sampling, and quantification. Biogeosciences, 2014, 11, 6029-6046.	3.3	73

HEIKO SAHLING

#	Article	IF	CITATIONS
19	Distribution and temporal variation of megaâ€fauna at the <scp>R</scp> egab pockmark ( <scp>N</scp> orthern <scp>C</scp> ongo <scp>F</scp> an), based on a comparison of videomosaics and geographic information systems analyses. Marine Ecology, 2014, 35, 77-95.	1.1	34
20	Fluid flow regimes and growth of a giant pockmark. Geology, 2014, 42, 63-66.	4.4	50
21	Natural oil seepage at Kobuleti Ridge, eastern Black Sea. Marine and Petroleum Geology, 2014, 50, 68-82.	3.3	60
22	Seepage of methane at Jaco Scar, a slide caused by seamount subduction offshore Costa Rica. International Journal of Earth Sciences, 2014, 103, 1801-1815.	1.8	16
23	Methane fluxes and carbonate deposits at a cold seep area of the Central Nile Deep Sea Fan, Eastern Mediterranean Sea. Marine Geology, 2014, 347, 27-42.	2.1	65
24	Megafaunal distribution and assessment of total methane and sulfide consumption by mussel beds at Menez Gwen hydrothermal vent, based on geo-referenced photomosaics. Deep-Sea Research Part I: Oceanographic Research Papers, 2013, 75, 93-109.	1.4	33
25	LAPM: a tool for underwater large-area photo-mosaicking. Geoscientific Instrumentation, Methods and Data Systems, 2013, 2, 189-198.	1.6	9
26	Quantification of gas bubble emissions from submarine hydrocarbon seeps at the Makran continental margin (offshore Pakistan). Journal of Geophysical Research, 2012, 117, .	3.3	108
27	Molecular taxonomy reveals broad trans-oceanic distributions and high species diversity of deep-sea clams (Bivalvia: Vesicomyidae: Pliocardiinae) in chemosynthetic environments. Systematics and Biodiversity, 2012, 10, 403-415.	1.2	40
28	Geological control and magnitude of methane ebullition from a high-flux seep area in the Black Sea—the Kerch seep area. Marine Geology, 2012, 319-322, 57-74.	2.1	92
29	Interaction between hydrocarbon seepage, chemosynthetic communities, and bottom water redox at cold seeps of the Makran accretionary prism: insights from habitat-specific pore water sampling and modeling. Biogeosciences, 2012, 9, 2013-2031.	3.3	87
30	Patterns of carbonate authigenesis at the Kouilou pockmarks on the Congo deep-sea fan. Marine Geology, 2010, 268, 129-136.	2.1	100
31	Morpho-acoustic variability of cold seeps on the continental slope offshore Nicaragua: Result of fluid flow interaction with sedimentary processes. Marine Geology, 2010, 275, 53-65.	2.1	20
32	Automated gas bubble imaging at sea floor – a new method of in situ gas flux quantification. Ocean Science, 2010, 6, 549-562.	3.4	19
33	Abyssogena: a new genus of the family Vesicomyidae (Bivalvia) from deep-water vents and seeps. Journal of Molluscan Studies, 2010, 76, 107-132.	1.2	48
34	Origin, distribution, and alteration of asphalts at Chapopote Knoll, Southern Gulf of Mexico. Marine and Petroleum Geology, 2010, 27, 1093-1106.	3.3	50
35	Molecular and isotopic partitioning of low-molecular-weight hydrocarbons during migration and gas hydrate precipitation in deposits of a high-flux seepage site. Chemical Geology, 2010, 269, 350-363.	3.3	102
36	Vesicomyidae (Bivalvia): Current Taxonomy and Distribution. PLoS ONE, 2010, 5, e9957.	2.5	101

HEIKO SAHLING

#	Article	IF	CITATIONS
37	Mineralization of vestimentiferan tubes at methane seeps on the Congo deep-sea fan. Deep-Sea Research Part I: Oceanographic Research Papers, 2009, 56, 283-293.	1.4	43
38	Salt tectonics and mud volcanism in the Latakia and Cyprus Basins, eastern Mediterranean. Tectonophysics, 2009, 470, 173-182.	2.2	27
39	Vodyanitskii mud volcano, Sorokin trough, Black Sea: Geological characterization and quantification of gas bubble streams. Marine and Petroleum Geology, 2009, 26, 1799-1811.	3.3	93
40	Pockmarks in the Northern Congo Fan area, SW Africa: Complex seafloor features shaped by fluid flow. Marine Geology, 2008, 249, 206-225.	2.1	95
41	Hydrogeological system of erosional convergent margins and its influence on tectonics and interplate seismogenesis. Geochemistry, Geophysics, Geosystems, 2008, 9, .	2.5	159
42	Fluid seepage at the continental margin offshore Costa Rica and southern Nicaragua. Geochemistry, Geophysics, Geosystems, 2008, 9, .	2.5	123
43	Hydroacoustic methodology for detection, localization, and quantification of gas bubbles rising from the seafloor at gas seeps from the eastern Black Sea. Geochemistry, Geophysics, Geosystems, 2008, 9, .	2.5	101
44	Estimates of methane output from mud extrusions at the erosive convergent margin off Costa Rica. Marine Geology, 2006, 225, 129-144.	2.1	94
45	Acoustic investigation of cold seeps offshore Georgia, eastern Black Sea. Marine Geology, 2006, 231, 51-67.	2.1	84
46	Recent bivalve molluscs of the genus Calyptogena (Vesicomyidae). Journal of Molluscan Studies, 2006, 72, 359-395.	1.2	67
47	Presence of two phylogenetically distinct groups in the deep-sea mussel Acharax (Mollusca: Bivalvia:) Tj ETQq1	1 0.78431 1.9	4 rgBT /Over
48	The physicochemical habitat of <i>Sclerolinum</i> sp. at Hook Ridge hydrothermal vent, Bransfield Strait, Antarctica. Limnology and Oceanography, 2005, 50, 598-606.	3.1	45
49	Styles and Productivity of Mud Diapirism along the Middle American Margin. , 2005, , 49-76.		14
50	Mapping deep-water gas emissions with sidescan sonar. Eos, 2005, 86, 341.	0.1	28
51	Fluid venting activity on the Costa Rica margin: new results from authigenic carbonates. International Journal of Earth Sciences, 2004, 93, 596.	1.8	96
52	Depth-related structure and ecological significance of cold-seep communities—a case study from the Sea of Okhotsk. Deep-Sea Research Part I: Oceanographic Research Papers, 2003, 50, 1391-1409.	1.4	136
53	Characteristics of an active vent in the fore-arc basin of the Sunda Arc, Indonesia. Marine Geology, 2002, 184, 121-141.	2.1	22
54	Hot vents in an ice-cold ocean: Indications for phase separation at the southernmost area of hydrothermal activity, Bransfield Strait, Antarctica. Earth and Planetary Science Letters, 2001, 193, 381-394.	4.4	34

Heiko Sahling

#	Article	IF	CITATIONS
55	Discovery of new hydrothermal vent sites in Bransfield Strait, Antarctica. Earth and Planetary Science Letters, 2001, 193, 395-407.	4.4	86
56	Hydrothermal activity at Hook Ridge in the Central Bransfield Basin, Antarctica. Geo-Marine Letters, 1998, 18, 277-284.	1.1	31
57	Fluid venting in the eastern Aleutian Subduction Zone. Journal of Geophysical Research, 1998, 103, 2597-2614.	3.3	123
58	Quantifying fluid flow, solute mixing, and biogeochemical turnover at cold vents of the eastern Aleutian subduction zone. Geochimica Et Cosmochimica Acta, 1997, 61, 5209-5219.	3.9	143