

Yoshio Ohno

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/28932/publications.pdf>

Version: 2024-02-01

51
papers

1,163
citations

567281

15
h-index

395702

33
g-index

54
all docs

54
docs citations

54
times ranked

1832
citing authors

#	ARTICLE	IF	CITATIONS
1	Prognostic Value of Neutrophil-to-lymphocyte Ratio and Establishment of Novel Preoperative Risk Stratification Model in Bladder Cancer Patients Treated With Radical Cystectomy. <i>Urology</i> , 2012, 79, 1085-1091.	1.0	212
2	Pretreatment Neutrophil-to-Lymphocyte Ratio as an Independent Predictor of Recurrence in Patients With Nonmetastatic Renal Cell Carcinoma. <i>Journal of Urology</i> , 2010, 184, 873-878.	0.4	182
3	Followup of Neutrophil-to-Lymphocyte Ratio and Recurrence of Clear Cell Renal Cell Carcinoma. <i>Journal of Urology</i> , 2012, 187, 411-417.	0.4	114
4	Sarcopenia as a Novel Preoperative Prognostic Predictor for Survival in Patients with Bladder Cancer Undergoing Radical Cystectomy. <i>Annals of Surgical Oncology</i> , 2016, 23, 1048-1054.	1.5	72
5	Association of legumain expression pattern with prostate cancer invasiveness and aggressiveness. <i>World Journal of Urology</i> , 2013, 31, 359-364.	2.2	62
6	Role of systemic inflammatory response markers in urological malignancy. <i>International Journal of Urology</i> , 2019, 26, 31-47.	1.0	50
7	Clinical variables for predicting metastatic renal cell carcinoma patients who might not benefit from cytoreductive nephrectomy: neutrophil-to-lymphocyte ratio and performance status. <i>International Journal of Clinical Oncology</i> , 2014, 19, 139-145.	2.2	49
8	Impact of Tumor Size on Renal Function and Prediction of Renal Insufficiency After Radical Nephrectomy in Patients With Renal Cell Carcinoma. <i>Journal of Urology</i> , 2011, 186, 1242-1246.	0.4	39
9	Clinical significance of preoperative peripheral blood neutrophil count in patients with non-metastatic upper urinary tract carcinoma. <i>World Journal of Urology</i> , 2013, 31, 953-958.	2.2	33
10	C-reactive protein-albumin ratio as a prognostic factor in renal cell carcinoma – A data from multi-institutional study in Japan. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 812.e1-812.e8.	1.6	29
11	Clinical Implications of Preoperative Serum Total Cholesterol in Patients With Clear Cell Renal Cell Carcinoma. <i>Urology</i> , 2014, 83, 154-158.	1.0	26
12	Prediction of biochemical recurrence after robot-assisted radical prostatectomy: Analysis of 784 Japanese patients. <i>International Journal of Urology</i> , 2015, 22, 188-193.	1.0	21
13	Prediction of non-muscle invasive bladder cancer recurrence using machine learning of quantitative nuclear features. <i>Modern Pathology</i> , 2022, 35, 533-538.	5.5	21
14	Prognostic value of preoperative pyuria in patients with non-muscle-invasive bladder cancer. <i>International Journal of Urology</i> , 2015, 22, 645-649.	1.0	18
15	Factors predicting incisional surgical site infection in patients undergoing open radical cystectomy for bladder cancer. <i>International Journal of Clinical Oncology</i> , 2014, 19, 935-939.	2.2	15
16	Preoperative predictive factors and further risk stratification of biochemical recurrence in clinically localized high-risk prostate cancer. <i>International Journal of Clinical Oncology</i> , 2016, 21, 595-600.	2.2	15
17	Development of a Nomogram for Predicting Severe Neutropenia Associated With Docetaxel-Based Chemotherapy in Patients With Castration-Resistant Prostate Cancer. <i>Clinical Genitourinary Cancer</i> , 2017, 15, 176-181.	1.9	15
18	The Impact of Lateral Bladder Neck Preservation on Urinary Continence Recovery After Robot-Assisted Radical Prostatectomy. <i>Journal of Endourology</i> , 2018, 32, 40-45.	2.1	15

#	ARTICLE	IF	CITATIONS
19	Prediction of renal function after nephroureterectomy in patients with upper tract urothelial carcinoma. Japanese Journal of Clinical Oncology, 2015, 45, 1064-1068.	1.3	14
20	Prognostic Value of the LATITUDE and CHAARTED Risk Criteria for Predicting the Survival of Men with Bone Metastatic Hormone-Naïve Prostate Cancer Treated with Combined Androgen Blockade Therapy: Real-World Data from a Japanese Multi-Institutional Study. BioMed Research International, 2020, 2020, 1-7.	1.9	14
21	The CANLPH Score, an Integrative Model of Systemic Inflammation and Nutrition Status (SINS), Predicts Clinical Outcomes After Surgery in Renal Cell Carcinoma: Data From a Multicenter Cohort in Japan. Annals of Surgical Oncology, 2019, 26, 2994-3004.	1.5	13
22	Pneumatosis intestinalis and hepatic portal venous gas in a patient receiving sorafenib. International Journal of Urology, 2012, 19, 1041-1042.	1.0	11
23	Association between preoperative serum total cholesterol level and biochemical recurrence in prostate cancer patients who underwent radical prostatectomy. Molecular and Clinical Oncology, 2016, 4, 1073-1077.	1.0	11
24	Impact of a preoperatively estimated prostate volume using transrectal ultrasonography on surgical and oncological outcomes in a single surgeon's experience with robot-assisted radical prostatectomy. Surgical Endoscopy and Other Interventional Techniques, 2016, 30, 3702-3708.	2.4	10
25	Preoperative determinant of early postoperative renal function following radical cystectomy and intestinal urinary diversion. International Urology and Nephrology, 2017, 49, 233-238.	1.4	10
26	Characterization and gene expression analysis of novel matched primary and metastatic renal cell carcinoma cell lines. Oncology Reports, 2008, 20, 501-9.	2.6	10
27	Prognostic significance of tenascin-C expression in clear cell renal cell carcinoma. Oncology Reports, 2008, 20, 511-6.	2.6	10
28	Prognostic Value of Platelet Counts in Patients with Metastatic Prostate Cancer Treated with Endocrine Therapy. Urology Journal, 2020, 17, 42-49.	0.4	9
29	Prognostic implication of infiltrative growth pattern and establishment of novel risk stratification model for survival in patients with upper urinary tract urothelial carcinoma. International Journal of Clinical Oncology, 2014, 19, 373-378.	2.2	7
30	Clinical significance of preoperative renal function and gross hematuria for intravesical recurrence after radical nephroureterectomy for upper tract urothelial carcinoma. International Journal of Urology, 2017, 24, 111-116.	1.0	6
31	Clinical efficacy and prognostic factors of tumor progression in Japanese patients with advanced renal cell carcinoma treated with sorafenib. Japanese Journal of Clinical Oncology, 2015, 45, 274-280.	1.3	5
32	No clinical significance of the time interval between biopsy and robotic-assisted radical prostatectomy for patients with clinically localized prostate cancer on biochemical recurrence: a propensity score matching analysis. Japanese Journal of Clinical Oncology, 2017, 47, 1083-1089.	1.3	5
33	The significance of micro-lymphatic invasion and pathological Gleason score in prostate cancer patients with pathologically organ-confined disease and negative surgical margins after robot-assisted radical prostatectomy. International Journal of Clinical Oncology, 2020, 25, 377-383.	2.2	5
34	Adjuvant chemotherapy improves overall survival in patients with localized upper tract urothelial carcinoma harboring pathologic vascular invasion: a propensity score-matched analysis of multi-institutional cohort. World Journal of Urology, 2020, 38, 3183-3190.	2.2	5
35	Preoperative prognostic factors for biochemical recurrence after robot-assisted radical prostatectomy in Japan. International Journal of Clinical Oncology, 2014, 19, 702-707.	2.2	4
36	Ureteral intussusception associated with a fibroepithelial polyp: a case report. Clinical Imaging, 2015, 39, 901-903.	1.5	4

#	ARTICLE	IF	CITATIONS
37	Editorial Comment to High neutrophil-to-lymphocyte ratio predicts worse overall survival in patients with advanced/metastatic urothelial bladder cancer. <i>International Journal of Urology</i> , 2018, 25, 238-239.	1.0	3
38	Predicting factors for progression to castration resistance prostate cancer after biochemical recurrence in patients with clinically localized prostate cancer who underwent radical prostatectomy. <i>International Journal of Clinical Oncology</i> , 2020, 25, 1704-1710.	2.2	3
39	Distinct effect of body mass index by sex as a prognostic factor in localized renal cell carcinoma treated with nephrectomy – data from a multi-institutional study in Japan ~. <i>BMC Cancer</i> , 2021, 21, 201.	2.6	3
40	Re: Impact of ABO Blood Type on Outcomes in Patients with Primary Nonmuscle Invasive Bladder Cancer. <i>European Urology</i> , 2014, 66, 391.	1.9	2
41	Is scintigraphy necessary to detect migration of 125I seeds after brachytherapy for early prostate cancer?. <i>International Journal of Clinical Oncology</i> , 2016, 21, 397-401.	2.2	2
42	Prostate-specific antigen screening impacts on biochemical recurrence in patients with clinically localized prostate cancer. <i>International Journal of Urology</i> , 2018, 25, 561-567.	1.0	2
43	Salvage radiation therapy for prostate cancer patients after prostatectomy. <i>Japanese Journal of Clinical Oncology</i> , 2019, 49, 281-286.	1.3	2
44	Prostate-specific antigen nomogram to predict advanced prostate cancer using area under the receiver operating characteristic curve boosting. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2022, 40, 162.e9-162.e16.	1.6	2
45	Prediction of Multifocal Lesions in Patients With Upper Tract Urothelial Carcinoma. <i>Urology</i> , 2014, 84, 869-874.	1.0	1
46	A Case of Primary Small-Cell Carcinoma of the Bladder. <i>Case Reports in Oncology</i> , 2017, 9, 574-579.	0.7	1
47	Fatty acids bound to albumin induce prostaglandin E2 production in human renal proximal tubular epithelial cell line HK-2. <i>Biochemical and Biophysical Research Communications</i> , 2020, 530, 273-277.	2.1	1
48	Abstract of Poster Presentation. <i>Human Cell</i> , 2005, 18, 43-65.	2.7	0
49	Editorial Comment. <i>Journal of Urology</i> , 2014, 192, 1618-1618.	0.4	0
50	Re: Low Pretreatment Neutrophil-to-Lymphocyte Ratio Predicts for Good Outcomes in Patients Receiving Neoadjuvant Chemotherapy Before Radical Cystectomy for Muscle Invasive Bladder Cancer. <i>European Urology</i> , 2017, 72, 1026.	1.9	0
51	Prognostic significance of the presence of tertiary Gleason grade 5 in robot-assisted radical prostatectomy specimens in Japanese patients with clinically localized prostate cancer. <i>Japanese Journal of Clinical Oncology</i> , 2019, 49, 276-280.	1.3	0