Hatice Tugba Dogmus Lehtijrvi

List of Publications by Citations

Source:

https://exaly.com/author-pdf/2898278/hatice-tugba-dogmus-lehtijarvi-publications-by-citations.pdf **Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

8 18 359 21 h-index g-index citations papers 21 1.5 2.24 445 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
21	Widespread Phytophthora infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of Phytophthora diseases. <i>Forest Pathology</i> , 2016 , 46, 134-163	1.2	187
20	Global geographic distribution and host range of Dothistroma species: a comprehensive review. <i>Forest Pathology</i> , 2016 , 46, 408-442	1.2	61
19	A review of Pinaceae resistance mechanisms against needle and shoot pathogens with a focus on the Dothistroma P inus interaction. <i>Forest Pathology</i> , 2016 , 46, 453-471	1.2	17
18	Heterobasidion abietinum on Abies species in western Turkey. Forest Pathology, 2006, 36, 280-286	1.2	15
17	Heterobasidion on Abies nordmanniana in north-eastern Turkey. Forest Pathology, 2007 , 37, 387-390	1.2	11
16	The efficacy of selected biological and chemical control agents against Heterobasidion abietinum on Abies cilicica. <i>Forest Pathology</i> , 2011 , 41, 470-476	1.2	10
15	Invasive forest pathogens in Europe: Cross-country variation in public awareness but consistency in policy acceptability. <i>Ambio</i> , 2019 , 48, 1-12	6.5	9
14	Dothistroma spp. in Western Ukraine and Georgia. Forest Pathology, 2018, 48, e12409	1.2	8
13	Oomycota species associated with deciduous and coniferous seedlings in forest tree nurseries of Western Turkey. <i>Forest Pathology</i> , 2017 , 47, e12363	1.2	7
12	Cedrus libani: the most susceptible Turkish conifer species to local Heterobasidion isolates in spring inoculations. <i>Forest Pathology</i> , 2011 , 41, 1-6	1.2	6
11	Pathogenicity of Heterobasidion annosum (Fr.) Bref. sensu stricto on coniferous tree species in Turkey. <i>Forest Pathology</i> , 2016 , 46, 22-28	1.2	6
10	Ceratocystis platani is killing plane trees in Istanbul (Turkey). Forest Pathology, 2018, 48, e12375	1.2	6
9	Armillaria ostoyae in managed coniferous forests in Kastamonu in Turkey. <i>Forest Pathology</i> , 2017 , 47, e12364	1.2	4
8	Turkish Heterobasidion abietinum is pathogenic to inoculated Abies nordmanniana ssp. nordmanniana and ssp. bornm[leriana. <i>Forest Pathology</i> , 2009 , 39, 200-209	1.2	4
7	Phytophthora species detected in the rhizosphere of Alnus glutinosa stands in the Floodplain Forests of Western Turkey. <i>Forest Pathology</i> , 2018 , 48, e12470	1.2	4
6	First Report of Brown Felt Blight Caused by Herpotrichia juniperi on Cedrus libani in Turkey. <i>Plant Disease</i> , 2011 , 95, 222	1.5	2
5	European pear rust on Juniperus excelsa L. in south-western Turkey. <i>Forest Pathology</i> , 2009 , 39, 35-42	1.2	1

LIST OF PUBLICATIONS

4	Impacts of inoculation with Herpotrichia pinetorum, Gremmenia infestans and Gremmeniella abietina on Pinus nigra subsp. pallasiana and Cedrus libani seedlings in the field. <i>Forest Pathology</i> , 2016 , 46, 47-53	1.2	1
3	Population structure of Heterobasidion annosum (Fr.) Bref. sensu stricto in Pinus brutia Ten. in south-western Turkey. <i>Forest Pathology</i> , 2021 , 51, e12715	1.2	O
2	Seed quantity affects the fungal community composition detected using metabarcoding <i>Scientific Reports</i> , 2022 , 12, 3060	4.9	
1	Detection and Identification of the Causal Agents of Dothistroma Needle Blight. <i>Methods in Molecular Biology</i> , 2022 , 155-166	1.4	