

Jon S Harding

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2907504/publications.pdf>

Version: 2024-02-01

45
papers

1,441
citations

430874

18
h-index

330143

37
g-index

45
all docs

45
docs citations

45
times ranked

1659
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Stream biomonitoring using macroinvertebrates around the globe: a comparison of large-scale programs. <i>Environmental Monitoring and Assessment</i> , 2015, 187, 4132. | 2.7 | 209 |
| 2 | Habitat loss drives threshold response of benthic invertebrate communities to deposited sediment in agricultural streams. <i>Ecological Applications</i> , 2013, 23, 1036-1047. | 3.8 | 172 |
| 3 | Consequences of acid mine drainage for the structure and function of benthic stream communities: a review. <i>Freshwater Science</i> , 2012, 31, 108-120. | 1.8 | 158 |
| 4 | The Biological Assessment and Rehabilitation of the World's Rivers: An Overview. <i>Water (Switzerland)</i> , 2021, 13, 371. | 2.7 | 88 |
| 5 | Effects of contrasting land use on physicochemical conditions and benthic assemblages of streams in a Canterbury (South Island, New Zealand) river system. <i>New Zealand Journal of Marine and Freshwater Research</i> , 1995, 29, 479-492. | 2.0 | 81 |
| 6 | Riparian shading mitigates stream eutrophication in agricultural catchments. <i>Freshwater Science</i> , 2014, 33, 73-84. | 1.8 | 71 |
| 7 | Anthropogenic and natural sources of acidity and metals and their influence on the structure of stream food webs. <i>Environmental Pollution</i> , 2012, 162, 466-474. | 7.5 | 54 |
| 8 | Stream faunas and ecoregions in South Island, New Zealand: do they correspond?. <i>Fundamental and Applied Limnology</i> , 1997, 140, 289-307. | 0.7 | 53 |
| 9 | Improving the effectiveness of riparian management for aquatic invertebrates in a degraded agricultural landscape: stream size and land-use legacies. <i>Journal of Applied Ecology</i> , 2012, 49, 213-222. | 4.0 | 50 |
| 10 | Historic deforestation and the fate of endemic invertebrate species in streams. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2003, 37, 333-345. | 2.0 | 42 |
| 11 | Heavy metals: confounding factors in the response of New Zealand freshwater fish assemblages to natural and anthropogenic acidity. <i>Science of the Total Environment</i> , 2010, 408, 3240-3250. | 8.0 | 37 |
| 12 | Inferring predator-prey interactions in food webs. <i>Methods in Ecology and Evolution</i> , 2019, 10, 356-367. | 5.2 | 35 |
| 13 | Mechanisms of trophic niche compression: Evidence from landscape disturbance. <i>Journal of Animal Ecology</i> , 2020, 89, 730-744. | 2.8 | 34 |
| 14 | Response of a new zealand mayfly (<i>Deleatidium</i> spp.) to acid mine drainage: Implications for mine remediation. <i>Environmental Toxicology and Chemistry</i> , 2008, 27, 1135-1140. | 4.3 | 31 |
| 15 | An Ecoregion Classification of the South Island, New Zealand. <i>Journal of Environmental Management</i> , 1997, 51, 275-287. | 7.8 | 29 |
| 16 | Longitudinal patterns in benthic communities in an urban stream under restoration. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2005, 39, 17-28. | 2.0 | 27 |
| 17 | Meta-community theory and stream restoration: evidence that spatial position constrains stream invertebrate communities in a mine impacted landscape. <i>Restoration Ecology</i> , 2015, 23, 284-291. | 2.9 | 25 |
| 18 | Leaf litter additions enhance stream metabolism, denitrification, and restoration prospects for agricultural catchments. <i>Ecosphere</i> , 2017, 8, e02018. | 2.2 | 25 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Assemblage-based biomonitoring of freshwater ecosystem health via multimetric indices: A critical review and suggestions for improving their applicability. , 2022, 1, 100054. | | 22 |
| 20 | Anthropogenic mining alters macroinvertebrate size spectra in streams. <i>Freshwater Biology</i> , 2019, 64, 81-92. | 2.4 | 21 |
| 21 | Strategies for coexistence in two species of New Zealand Hydropsychidae (Trichoptera). <i>Hydrobiologia</i> , 1997, 350, 25-33. | 2.0 | 15 |
| 22 | Variations in benthic fauna between differing lake outlet types in New Zealand. <i>New Zealand Journal of Marine and Freshwater Research</i> , 1994, 28, 417-427. | 2.0 | 13 |
| 23 | Capacity for bioreactors and riparian rehabilitation to enhance nitrate attenuation in agricultural streams. <i>Ecological Engineering</i> , 2019, 134, 65-77. | 3.6 | 13 |
| 24 | Persistence of a significant population of rare Canterbury mudfish (<i>Neochanna burrowsius</i>) in a hydrologically isolated catchment. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2007, 41, 309-316. | 2.0 | 12 |
| 25 | Leaf breakdown, detrital resources, and food webs in streams affected by mine drainage. <i>Hydrobiologia</i> , 2013, 716, 59-73. | 2.0 | 12 |
| 26 | Physicochemical parameters and invertebrate faunas of three lake inflows and outlets in Westland, New Zealand. <i>New Zealand Journal of Marine and Freshwater Research</i> , 1992, 26, 95-102. | 2.0 | 11 |
| 27 | Comparison of fluorescent lights with differing spectral properties on catches of adult aquatic and terrestrial insects. <i>New Zealand Entomologist</i> , 2018, 41, 1-11. | 0.3 | 10 |
| 28 | Feeding ecology of <i>Aoteapsyche raruraru</i> (McFarlane) (Trichoptera: Hydropsychidae) in a New Zealand Lake Outlet. <i>Aquatic Insects</i> , 1997, 19, 51-63. | 0.9 | 8 |
| 29 | Distinctive aquatic assemblages in water-filled tree holes: a novel component of freshwater biodiversity in New Zealand temperate rainforests. <i>Insect Conservation and Diversity</i> , 2012, 5, 202-212. | 3.0 | 8 |
| 30 | Do food quantity and quality affect food webs in streams polluted by acid mine drainage?. <i>Marine and Freshwater Research</i> , 2013, 64, 1112. | 1.3 | 8 |
| 31 | Changes in stream foodweb structure across a gradient of acid mine drainage increase local community stability. <i>Ecology</i> , 2020, 101, e03102. | 3.2 | 8 |
| 32 | Discontinuities in the distribution of invertebrates in impounded south island rivers, New Zealand. <i>River Research and Applications</i> , 1992, 7, 327-335. | 0.8 | 7 |
| 33 | Life history and production of <i>Coloburiscus humeralis</i> (Ephemeroptera: Oligoneuriidae) in two South Island high-country streams, New Zealand. <i>New Zealand Journal of Marine and Freshwater Research</i> , 1993, 27, 445-451. | 2.0 | 7 |
| 34 | Evaluating practical macrophyte control tools on small agricultural waterways in Canterbury, New Zealand. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2019, 53, 182-200. | 2.0 | 6 |
| 35 | Shifts in population size structure for a drying-tolerant fish in response to extreme drought. <i>Austral Ecology</i> , 2019, 44, 658-667. | 1.5 | 6 |
| 36 | Ecological processes mediate the effects of the invasive bloom-forming diatom <i>Didymosphenia geminata</i> on stream algal and invertebrate assemblages. <i>Hydrobiologia</i> , 2020, 847, 177-190. | 2.0 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Trialling tools using hand-weeding, weed mat and artificial shading to control nuisance macrophyte growth at multiple scales in small agricultural waterways. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2020, 54, 512-526. | 2.0 | 5 |
| 38 | Partnerships Generate Co-Benefits in Agricultural Stream Restoration (Canterbury, New Zealand). <i>Case Studies in the Environment</i> , 2020, 4, . | 0.7 | 5 |
| 39 | Distribution, nymphal habitat, genetic structure and conservation of the New Zealand mayfly <i>Isothraulus abditus</i> (Insecta: Ephemeroptera) and a description of its subimago. <i>New Zealand Journal of Zoology</i> , 2019, 46, 13-30. | 1.1 | 4 |
| 40 | Analysis of the conservation status of New Zealand freshwater invertebrates: temporal changes, knowledge gaps, impediments, and management implications. <i>New Zealand Journal of Zoology</i> , 2021, 48, 81-96. | 1.1 | 4 |
| 41 | Distribution, body size, genetic structure and conservation of <i>Siphlaenigma janae</i> (Insecta: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj 5 | 1.1 | 4 |
| 42 | Acute toxicity of arsenic to larvae of four New Zealand freshwater insect taxa. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2017, 51, 443-454. | 2.0 | 2 |
| 43 | Faecal indicator bacteria in New Zealand freshwater fish: a pilot study. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2019, 53, 470-479. | 2.0 | 2 |
| 44 | Benthic Invertebrate Indices Show No Response to High Nitrate-Nitrogen in Lowland Agricultural Streams. <i>Water, Air, and Soil Pollution</i> , 2021, 232, 1. | 2.4 | 1 |
| 45 | Riparian plant species offer a range of organic resources to stream invertebrate communities through varied leaf breakdown rates. <i>New Zealand Journal of Marine and Freshwater Research</i> , 0, , 1-16. | 2.0 | 0 |