

Ashfaque A Memon

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2912852/publications.pdf>

Version: 2024-02-01

59
papers

1,612
citations

331670

21
h-index

315739

38
g-index

59
all docs

59
docs citations

59
times ranked

2640
citing authors

#	ARTICLE	IF	CITATIONS
1	Role of mitochondrial DNA copy number in incident cardiovascular diseases and the association between cardiovascular disease and type 2 diabetes: A follow-up study on middle-aged women. <i>Atherosclerosis</i> , 2022, 341, 58-62.	0.8	18
2	Novel cardiovascular biomarkers associated with peripheral arterial disease in men screened for abdominal aortic aneurysm. <i>Vasa - European Journal of Vascular Medicine</i> , 2022, 51, 167-173.	1.4	4
3	The association of mitochondrial DNA copy number with incident mental disorders in women: A population-based follow-up study. <i>Journal of Affective Disorders</i> , 2022, 308, 111-115.	4.1	4
4	Association of mitochondrial DNA copy number with prevalent and incident type 2 diabetes in women: A population-based follow-up study. <i>Scientific Reports</i> , 2021, 11, 4608.	3.3	24
5	Mitochondria-DNA copy-number and incident venous thromboembolism among middle-aged women: a population-based cohort study. <i>Journal of Thrombosis and Thrombolysis</i> , 2021, 52, 148-157.	2.1	2
6	Elucidating causal effects of type 2 diabetes on ischemic heart disease from observational data on middle-aged Swedish women: a triangular analytical approach. <i>Scientific Reports</i> , 2021, 11, 12579.	3.3	3
7	Association of Mitochondrial DNA Copy Number and Telomere Length with Prevalent and Incident Cancer and Cancer Mortality in Women: A Prospective Swedish Population-Based Study. <i>Cancers</i> , 2021, 13, 3842.	3.7	13
8	Inflammatory proteins and miRNA-144-5p in patients with depression, anxiety, or stress- and adjustment disorders after psychological treatment. <i>Cytokine</i> , 2021, 146, 155646.	3.2	10
9	The Association between Blood-Based Global DNA Methylation and Venous Thromboembolism. <i>Seminars in Thrombosis and Hemostasis</i> , 2021, 47, 662-668.	2.7	2
10	Association of circulating let-7b-5p with major depressive disorder: a nested case-control study. <i>BMC Psychiatry</i> , 2021, 21, 616.	2.6	7
11	Diagnostic potential of circulating cell-free nuclear and mitochondrial DNA for several cancer types and nonmalignant diseases: A study on suspected cancer patients. <i>Molecular Carcinogenesis</i> , 2020, 59, 1362-1370.	2.7	9
12	Macrophage Migration Inhibitory Factor as a Predictor for Long-term Improvements After Mindfulness-Based Group Therapy or Treatment as Usual for Depression, Anxiety or Stress and Adjustment Disorders. <i>Mindfulness</i> , 2020, 11, 1370-1377.	2.8	0
13	Long-term improvements after mindfulness-based group therapy of depression, anxiety and stress and adjustment disorders: A randomized controlled trial. <i>Microbial Biotechnology</i> , 2019, 13, 943-952.	1.7	11
14	Association of recurrent venous thromboembolism and circulating microRNAs. <i>Clinical Epigenetics</i> , 2019, 11, 28.	4.1	43
15	Risk prediction of recurrent venous thromboembolism: a multiple genetic risk model. <i>Journal of Thrombosis and Thrombolysis</i> , 2019, 47, 216-226.	2.1	18
16	Identification of novel diagnostic biomarkers for deep venous thrombosis. <i>British Journal of Haematology</i> , 2018, 181, 378-385.	2.5	28
17	Polymorphisms in PARK2 and MRPL37 are associated with higher risk of recurrent venous thromboembolism in a sex-specific manner. <i>Journal of Thrombosis and Thrombolysis</i> , 2018, 46, 154-165.	2.1	5
18	Macrophage Migration Inhibitory Factor and microRNA-451a in Response to Mindfulness-based Therapy or Treatment as Usual in Patients with Depression, Anxiety, or Stress and Adjustment Disorders. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 513-521.	2.1	18

#	ARTICLE	IF	CITATIONS
19	Fat mass and obesity-associated gene rs9939609 polymorphism is a potential biomarker of recurrent venous thromboembolism in male but not in female patients. <i>Gene</i> , 2018, 647, 136-142.	2.2	3
20	Evaluation of Expression Level of Apolipoprotein M as a Diagnostic Marker for Primary Venous Thromboembolism. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2018, 24, 416-422.	1.7	4
21	Alpha 2-macroglobulin 5 bp insertion/deletion polymorphism increases the risk of recurrent venous thromboembolism. <i>Gene Reports</i> , 2018, 13, 104-109.	0.8	0
22	Women's Health in the Lund Area (WHILA) study. Health problems and acute myocardial infarction in women – A 17-year follow-up study. <i>Maturitas</i> , 2018, 115, 45-50.	2.4	10
23	Risk of pulmonary embolism and deep venous thrombosis in patients with asthma: a nationwide case-control study from Sweden. <i>European Respiratory Journal</i> , 2017, 49, 1601014.	6.7	32
24	Identification of Genetic Aberrations in Thrombomodulin Gene in Patients With Recurrent Venous Thromboembolism. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017, 23, 319-328.	1.7	6
25	Association between TLR9 rs5743836 polymorphism and risk of recurrent venous thromboembolism. <i>Journal of Thrombosis and Thrombolysis</i> , 2017, 44, 130-138.	2.1	6
26	Role of IL-8, CRP and epidermal growth factor in depression and anxiety patients treated with mindfulness-based therapy or cognitive behavioral therapy in primary health care. <i>Psychiatry Research</i> , 2017, 254, 311-316.	3.3	44
27	Quantification of mitochondrial DNA copy number in suspected cancer patients by a well optimized ddPCR method. <i>Biomolecular Detection and Quantification</i> , 2017, 13, 32-39.	7.0	53
28	Leukocyte telomere length and depression, anxiety and stress and adjustment disorders in primary health care patients. <i>BMC Psychiatry</i> , 2017, 17, 148.	2.6	39
29	Association of mitochondrial DNA in peripheral blood with depression, anxiety and stress- and adjustment disorders in primary health care patients. <i>European Neuropsychopharmacology</i> , 2017, 27, 751-758.	0.7	25
30	Soluble HER3 predicts survival in bladder cancer patients. <i>Oncology Letters</i> , 2017, 15, 1783-1788.	1.8	5
31	Identification of polymorphisms in Apolipoprotein M gene and their relationship with risk of recurrent venous thromboembolism. <i>Thrombosis and Haemostasis</i> , 2016, 116, 432-441.	3.4	9
32	Palaeolithic diet decreases fasting plasma leptin concentrations more than a diabetes diet in patients with type 2 diabetes: a randomised cross-over trial. <i>Cardiovascular Diabetology</i> , 2016, 15, 80.	6.8	71
33	Epidemiology of Familial Aggregation of Venous Thromboembolism. <i>Seminars in Thrombosis and Hemostasis</i> , 2016, 42, 821-832.	2.7	10
34	Diagnostic potential of plasma microRNA signatures in patients with deep-vein thrombosis. <i>Thrombosis and Haemostasis</i> , 2016, 116, 328-336.	3.4	49
35	Thrombomodulin gene c.1418C>T polymorphism and risk of recurrent venous thromboembolism. <i>Journal of Thrombosis and Thrombolysis</i> , 2016, 42, 135-141.	2.1	12
36	Circulating human epidermal growth factor receptor 2 (<sc>HER2</sc>) is associated with hyperglycaemia and insulin resistance	1.8	18

#	ARTICLE	IF	CITATIONS
37	Women's health in the Lund area (WHILA) - Alcohol consumption and all-cause mortality among women - a 17-year follow-up study. BMC Public Health, 2015, 16, 22.	2.9	9
38	Role of family history of venous thromboembolism and thrombophilia as predictors of recurrence: a prospective follow-up study. Journal of Thrombosis and Haemostasis, 2015, 13, 2180-2186.	3.8	17
39	Plasminogen activator inhibitor-1 4G/5G polymorphism, factor V Leiden, prothrombin mutations and the risk of VTE recurrence. Thrombosis and Haemostasis, 2015, 114, 1156-1164.	3.4	14
40	Circulating microRNA-144-5p is associated with depressive disorders. Clinical Epigenetics, 2015, 7, 69.	4.1	84
41	Digested wheat gluten inhibits binding between leptin and its receptor. BMC Biochemistry, 2015, 16, 3.	4.4	8
42	Yoga's effect on inflammatory biomarkers and metabolic risk factors in a high risk population - a controlled trial in primary care. BMC Cardiovascular Disorders, 2015, 15, 91.	1.7	18
43	Factor H Autoantibodies in Patients with Antiphospholipid Syndrome and Thrombosis. Journal of Rheumatology, 2015, 42, 1786-1793.	2.0	29
44	Mindfulness group therapy in primary care patients with depression, anxiety and stress and adjustment disorders: Randomised controlled trial. British Journal of Psychiatry, 2015, 206, 128-135.	2.8	87
45	Determination of 14 Circulating microRNAs in Swedes and Iraqis with and without Diabetes Mellitus Type 2. PLoS ONE, 2014, 9, e86792.	2.5	104
46	Transforming growth factor (TGF)- β 2 levels and unprovoked recurrent venous thromboembolism. Journal of Thrombosis and Thrombolysis, 2014, 38, 348-354.	2.1	9
47	Apolipoprotein M and the risk of unprovoked recurrent venous thromboembolism. Thrombosis Research, 2014, 133, 322-326.	1.7	8
48	The association between apolipoprotein M and insulin resistance varies with country of birth. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 1174-1180.	2.6	15
49	The HER4 isoform JM-a/CYT2 relates to improved survival in bladder cancer patients but only if the estrogen receptor β is not expressed. Scandinavian Journal of Clinical and Laboratory Investigation, 2013, 73, 503-513.	1.2	5
50	The association between cytokines and insulin sensitivity in Iraqi immigrants and native Swedes. BMJ Open, 2013, 3, e003473.	1.9	8
51	Erlotinib Accumulation in Brain Metastases from Non-small Cell Lung Cancer: Visualization by Positron Emission Tomography in a Patient Harboring a Mutation in the Epidermal Growth Factor Receptor. Journal of Thoracic Oncology, 2011, 6, 1287-1289.	1.1	124
52	Calcium-induced apoptosis is delayed by HER1 receptor signalling through the Akt and PLC β 3 pathways in bladder cancer cells. Scandinavian Journal of Clinical and Laboratory Investigation, 2011, 71, 45-51.	1.2	2
53	PET imaging of patients with non-small cell lung cancer employing an EGF receptor targeting drug as tracer. British Journal of Cancer, 2011, 105, 1850-1855.	6.4	68
54	Positron Emission Tomography (PET) Imaging with [11C]-Labeled Erlotinib: A Micro-PET Study on Mice with Lung Tumor Xenografts. Cancer Research, 2009, 69, 873-878.	0.9	164

#	ARTICLE	IF	CITATIONS
55	Expression of the RAI gene is conducive to apoptosis: Studies of induction and interference. <i>Experimental Cell Research</i> , 2007, 313, 2611-2621.	2.6	23
56	The relation between survival and expression of HER1 and HER2 depends on the expression of HER3 and HER4: a study in bladder cancer patients. <i>British Journal of Cancer</i> , 2006, 94, 1703-1709.	6.4	74
57	The epidermal growth factor family has a dual role in deciding the fate of cancer cells. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2006, 66, 623-630.	1.2	10
58	Identification of differentially expressed proteins during human urinary bladder cancer progression. <i>Cancer Detection and Prevention</i> , 2005, 29, 249-255.	2.1	42
59	Expression of HER3, HER4 and their ligand heregulin-4 is associated with better survival in bladder cancer patients. <i>British Journal of Cancer</i> , 2004, 91, 2034-2041.	6.4	75