

# Zora PopoviÄ

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2932702/publications.pdf>

Version: 2024-02-01

110  
papers

1,471  
citations

361388

20  
h-index

414395

32  
g-index

112  
all docs

112  
docs citations

112  
times ranked

1432  
citing authors

#	ARTICLE	IF	CITATIONS
1	Mercury(II) complexes of heterocyclic thiones.. <i>Inorganica Chimica Acta</i> , 2000, 306, 142-152.	2.4	80
2	Mercury(II) compounds with 1,3-imidazole-2-thione and its 1-methyl analogue. Preparative and NMR spectroscopic studies. The crystal structures of di- $\frac{1}{4}$ -iodo-bis[iodo(1,3-imidazolium-2-thiolato-S)]mercury(II)], bis[bromo(1,3-imidazolium-2-thiolato-S)]mercury(II) and bis[ $\frac{1}{4}$ -(1-N-methyl-1,3-imidazole-2-thiolato-S)]mercury(II). <i>Inorganica Chimica Acta</i> , 1999, 294, 35-46.	2.4	64
3	Thiocyanate complexes of the group 12 metals with pyridine-2-carboxamide: Synthesis and structural characterization. <i>Polyhedron</i> , 2008, 27, 210-222.	2.2	63
4	The first example of coexistence of the ketoamino $\rightleftharpoons$ enolimino forms of diamine Schiff base naphthaldimine parts: the crystal and molecular structure of N,N $\epsilon$ -2-bis(1-naphthaldimine)-o-phenylenediamine chloroform (1/1) solvate at 200 K. <i>Journal of Molecular Structure</i> , 2001, 597, 39-47.	3.6	57
5	3D supramolecular architectures of copper(II) complexes with 6-methylpicolinic and 6-bromopicolinic acid: Synthesis, spectroscopic, thermal and magnetic properties. <i>Polyhedron</i> , 2008, 27, 3631-3638.	2.2	50
6	Association of copper(II) isonicotinamide moieties via different anionic bridging ligands: Two paths of ferromagnetic interaction in the azide coordination compound. <i>Polyhedron</i> , 2010, 29, 1910-1917.	2.2	47
7	Pd-porphyrin functionalized ionic liquid-modified mesoporous SBA-15: An efficient and recyclable catalyst for solvent-free Heck reaction. <i>Materials Research Bulletin</i> , 2010, 45, 1648-1653.	5.2	43
8	Preparation and spectroscopic properties of the complexes of mercuric thiocyanate with pyridine-2-thione and pyridine-2-carboxylic acid.. <i>Inorganica Chimica Acta</i> , 1999, 285, 208-216.	2.4	40
9	A functionalized ionic liquid containing phosphine-ligated palladium complex for the Sonogashira reactions under aerobic and CuI-free conditions. <i>Catalysis Communications</i> , 2012, 17, 160-163.	3.3	37
10	On tautomerism of two 5-methoxysalicylaldehyde structural isomers in the solid state: structural study of N-(o-hydroxyphenyl)-5-methoxysalicylaldehyde and N-(m-hydroxyphenyl)-5-methoxysalicylaldehyde. <i>Journal of Molecular Structure</i> , 2002, 615, 23-31.	3.6	35
11	Coordination-driven self-assembly of thiocyanate complexes of Co(ii), Ni(ii) and Cu(ii) with picolinamide: a structural and DFT study. <i>CrystEngComm</i> , 2011, 13, 5863.	2.6	35
12	Synthesis, spectroscopic and structural investigation of Zn(NCS) <sub>2</sub> (nicotinamide) <sub>2</sub> and [Hg(SCN) <sub>2</sub> (nicotinamide)] <sub>n</sub> . <i>Polyhedron</i> , 2008, 27, 465-472.	2.2	33
13	Synthesis and structure of cobalt(II) complexes with hydroxyl derivatives of pyridinecarboxylic acids: Conformation analysis of ligands in the solid state. <i>Journal of Molecular Structure</i> , 2008, 882, 47-55.	3.6	26
14	Bifunctional Ionic Liquid Catalyst Containing Sulfoacid Group and Hexafluorotitanate for Room Temperature Sulfoxidation of Sulfides to Sulfoxides Using Hydrogen Peroxide. <i>ACS Catalysis</i> , 2012, 2, 230-237.	11.2	25
15	Title is missing!. <i>Structural Chemistry</i> , 2002, 13, 425-436.	2.0	24
16	Mercury(II) Complexes with Heterocyclic Thiones $\rightleftharpoons$ Preparation and Characterization of the 1:1 and 1:2 Mercury(II) Complexes with Benzo-1,3-imidazole-2-thione. <i>European Journal of Inorganic Chemistry</i> , 2002, 2002, 171-180.	2.0	23
17	Coordination modes of 3-hydroxypicolinic acid: Synthesis and structural characterization of polymeric mercury(II) complexes. <i>Polyhedron</i> , 2007, 26, 1045-1052.	2.2	23
18	Pseudopolymorphism in Nickel(II) Complexes with 6-Methylpicolinate. Synthesis, Structural, Spectroscopic, Thermal, and Density Functional Theory Studies. <i>Crystal Growth and Design</i> , 2008, 8, 3465-3473.	3.0	23

#	ARTICLE	IF	CITATIONS
19	Nitroaniline Derivatives of 2-Oxo-1-naphthylideneaminesâ€™ Molecular Self-Assembling via Câ€™Hâ€™O Intermolecular Hydrogen Bonds and Stabilization of Oâ€™Hâ€™N and Nâ€™Hâ€™O Tautomers in Solution and Solid State. <i>Structural Chemistry</i> , 2004, 15, 587-598.	2.0	22
20	Structural study of picolinamide complexes of Ni(II), Zn(II), Cd(II), and Hg(II) nitrates in solid state and solution. <i>Journal of Coordination Chemistry</i> , 2012, 65, 1017-1032.	2.2	21
21	Stereochemistry of Hexacoordinated Zn(II), Cu(II), Ni(II), and Co(II) Complexes with Iminodiacetamide Ligands. <i>Inorganic Chemistry</i> , 2019, 58, 16445-16457.	4.0	20
22	Mercuration products of some aliphatic carboxylic acids: preparation and spectral properties. Crystal structure of the bipyridyl complex of the nitrate of monomercurated propionic acid. <i>Organometallics</i> , 1993, 12, 4708-4713.	2.3	19
23	Preparation and characterization of the 1:1 adducts of mercury(II) halides with N-benzyl- and N-p-tolyl-2-oxo-1-naphthylideneamine. The crystal and molecular structures of two isostructural di-1/4-halo-bis[halo(N-benzyl-2-oxo-1-naphthylideneamine)mercury(II)] adducts (halo=chloro, bromo). <i>Inorganica Chimica Acta</i> , 2001, 322, 65-73.	2.4	19
24	Coordination modes of 3-hydroxypicolinic acid (OH-picH): Synthesis and characterization of cadmium(II) complexes. <i>Polyhedron</i> , 2006, 25, 2353-2362.	2.2	19
25	A 1:2 complex of mercury(II) chloride with 1,3-imidazole-2-thione. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2000, 56, 61-63.	0.4	18
26	A Homogeneous Mixture Composed of Vanadate, Acid, and TEMPO Functionalized Ionic Liquids for Alcohol Oxidation by H <sub>2</sub> O <sub>2</sub> . <i>ChemCatChem</i> , 2011, 3, 1208-1213.	3.7	18
27	Supramolecular Architecture via Self-Complementary Amide Hydrogen Bonding in [Ag(pia) <sub>2</sub> ](NO <sub>3</sub> )·H <sub>2</sub> O: Synthesis, Thermal and Structural Studies. <i>Journal of Chemical Crystallography</i> , 2011, 41, 343-348.	1.1	18
28	Synthesis of an Ionic Paramagnetic Ruthenium(III) Complex and Its Application as an Efficient and Recyclable Catalyst for the Transfer Hydrogenation of Ketones. <i>European Journal of Inorganic Chemistry</i> , 2012, 2012, 3435-3440.	2.0	18
29	Addition compounds of bis(trifluoromethyl)mercury with tetraphenylphosphonium and tetraphenylarsonium halides and thiocyanates. <i>Inorganica Chimica Acta</i> , 1988, 150, 113-118.	2.4	17
30	Synthesis, X-ray structural, IR spectroscopic, thermal and DFT studies of nickel(II) and copper(II) complexes with 3-methylpicolinic acid. UV-Vis spectrophotometric study of complexation in the solution. <i>Polyhedron</i> , 2013, 52, 1349-1361.	2.2	17
31	Synthesis, structural, spectroscopic and thermal characterization of cobalt complexes with 3- and 6-methylpicolinic acid. Voltammetric and spectrophotometric study in solution. <i>Inorganica Chimica Acta</i> , 2009, 362, 2704-2714.	2.4	16
32	Synthesis and characterization of mercury(II) complexes with 2-styryl-1,3-benzothiazole (sb). Presence of two differently coordinated Hg(II) ions in the dinuclear complex Hg <sub>2</sub> Cl <sub>4</sub> (sb) <sub>3</sub> . Structural characterization of 2-styryl-1,3-benzothiazole and some of its derivatives. <i>Polyhedron</i> , 2007, 26, 5162-5170.	2.2	15
33	Self-Assembly of Two Isomorphous Thiocyanate Complexes of Co(II) and Ni(II) with 3-Hydroxypicolinamide. <i>Journal of Chemical Crystallography</i> , 2011, 41, 180-185.	1.1	15
34	Synthesis and characterization of copper(II) complexes with 3-methylpicolinic acid. Crystal and molecular structure of bis(3-methylpicolinato-N,O)Tj ETQq0 0 0 rgBT Lock 14 Tf 50 13	2.2	14
35	Polymorphism of cobalt(II) complex with 6-bromopicolinic acid: The influence of the solution pH value on the formation of polymorphs. <i>Journal of Molecular Structure</i> , 2009, 938, 174-178.	3.6	14
36	Various coordination modes in mercury(II) complexes with quinoline-2-carboxylic acid: Preparation and structural characterization. <i>Polyhedron</i> , 2009, 28, 2735-2743.	2.2	14

#	ARTICLE	IF	CITATIONS
37	Disappearing and Concomitant Polymorphism of Nickel(II) Complexes with 6-Hydroxypicolinic Acid. Structural and Density Functional Theory Studies. <i>Crystal Growth and Design</i> , 2010, 10, 3685-3693.	3.0	14
38	Preparation, structural, spectroscopic, thermal and DFT characterization of cadmium(II) complexes with quinaldic acid. <i>Inorganica Chimica Acta</i> , 2011, 378, 154-162.	2.4	14
39	Monomercurated acetic acid. The crystal structure of NO <sub>3</sub> HgCH <sub>2</sub> COOH. <i>Journal of Organometallic Chemistry</i> , 1991, 411, 19-23.	1.8	12
40	Title is missing!. <i>Structural Chemistry</i> , 2002, 13, 415-424.	2.0	12
41	Cis–trans isomerism in cobalt(II) complexes with 3-hydroxypicolinic acid. Structural, DFT and thermal studies. <i>Inorganica Chimica Acta</i> , 2010, 363, 1887-1896.	2.4	12
42	Mercuration of n-butyric, n-valeric and isovaleric acid. The crystal structure of C <sub>2</sub> H <sub>5</sub> C(HgCl)2COOH · DMSO. <i>Journal of Organometallic Chemistry</i> , 1991, 405, 59-65.	1.8	10
43	Mercury(II) complex with terpyridine: structure of the 2·1 solvate of [Hg(terpy) <sub>2</sub> ](CF <sub>3</sub> SO <sub>3</sub> ) <sub>2</sub> with acetone. <i>Journal of Chemical Crystallography</i> , 1995, 25, 453-458.	1.1	10
44	Pseudopolymorphism in nickel(II) complexes with 6-bromopicolinic acid. Synthesis, structural and thermal studies. <i>Journal of Molecular Structure</i> , 2009, 930, 121-125.	3.6	10
45	Structural preferences and isomerism in nickel(II) and copper(II) complexes with 3-hydroxypicolinic acid. <i>Polyhedron</i> , 2012, 39, 66-75.	2.2	10
46	Influence of electrostatic repulsive force and electron-withdrawing effect in ionic diphosphine on regioselectivity of rhodium-catalyzed hydroformylation of 1-octene. <i>Journal of Molecular Catalysis A</i> , 2015, 402, 37-45.	4.8	10
47	A diversity of hydrogen bond motifs in the crystal structures of nickel(II) and copper(II) complexes with N-arylalkyliminodiacetamides. <i>Polyhedron</i> , 2015, 93, 106-117.	2.2	10
48	Terpyridine complexes of Hg(OCOCF <sub>3</sub> ) <sub>2</sub> and Hg(BF <sub>4</sub> ) <sub>2</sub> . Crystal and molecular structure of mercuric trifluoroacetato-2,2,6,6-tetra-terpyridine. <i>Inorganica Chimica Acta</i> , 1991, 190, 169-172.	2.4	9
49	Mercuration Products of Some Aliphatic Ketones and Keto-Acids. 1H and 13C NMR Evidence on the Site and Degree of Mercuration. The Crystal and Molecular Structure of 1-(Acetoxymercuro) Butanone. <i>Main Group Chemistry</i> , 1996, 1, 373-385.	0.8	9
50	Hydrogen bonding interactions in the 2,2'-bipyridine–pyromellitic acid complex IR, nuclear magnetic resonance and theoretical studies. <i>Journal of the Chemical Society, Faraday Transactions</i> , 1998, 94, 1051-1056.	1.7	9
51	Supramolecular amide and thioamide synthons in hydrogen bonding patterns of N-aryl-furamides and N-aryl-thiofuramides. <i>Structural Chemistry</i> , 2006, 17, 275-285.	2.0	9
52	Synthesis, spectroscopic and structural investigation of two nickel(II) thiocyanate complexes with isonicotinic acid. <i>Journal of Molecular Structure</i> , 2008, 888, 394-400.	3.6	9
53	Synthesis and Characterization of a Copper(II) Complex with 6-Hydroxypicolinic Acid and 3-Picoline. <i>Croatica Chimica Acta</i> , 2012, 85, 479-483.	0.4	9
54	Can self-assembly of copper(ii) picolinamide building blocks be controlled?. <i>CrystEngComm</i> , 2013, 15, 8074.	2.6	9

#	ARTICLE	IF	CITATIONS
55	Influence of the non-coordinating alkyl chain type in N-alkylated iminodiacetamides on the stability and structure of their complexes with nickel(II) and cobalt(II). <i>Polyhedron</i> , 2017, 130, 115-126.	2.2	9
56	Diiodobis(1-methyl-1,3-imidazolium-2-thiolato-S)mercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2000, 56, 801-803.	0.4	8
57	A new complex of HgBr <sub>2</sub> and pyrimidine-2-thione: (pyrimidine-2-thionato- $\mu$ -S)(pyrimidinium-2-thionato- $\mu$ -S)mercury(II) tetrabromomercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2004, 60, m44-m46.	0.4	7
58	Synthesis and characterization of some novel cadmium(II) complexes with 3-hydroxypicolinic acid (3-OHpicH): Crystal and molecular structures of [CdI(3-OHpic)(3-OHpicH)(H <sub>2</sub> O)] <sub>2</sub> , [Cd(3-OHpic) <sub>2</sub> (H <sub>2</sub> O)] <sub>2</sub> and [Cd(3-OHpic) <sub>2</sub> ] <sub>n</sub> . <i>Polyhedron</i> , 2008, 27, 1479-1488.	2.2	7
59	Uncommon isonicotinamide supramolecular synthons in copper(II) complexes directed by nitrate and perchlorate anions. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2009, 65, m361-m366.	0.4	7
60	Preparation, spectroscopic, structural, and thermal characterization of vanadium complexes with 2-quinaldic acid. <i>Monatshefte für Chemie</i> , 2012, 143, 1471-1477.	1.8	7
61	Nickel(II) complexes with N-arylaminodiacetic acids: Preparation, spectroscopic, structural and thermal characterization. <i>Inorganica Chimica Acta</i> , 2013, 400, 122-129.	2.4	7
62	Preparation and characterization of palladium(II) complexes with N-arylalkyliminodiacetic acids. Catalytic activity of complexes in methoxycarbonylation of iodobenzene. <i>Journal of Organometallic Chemistry</i> , 2014, 760, 224-230.	1.8	7
63	Cobalt(II) and nickel(II) complexes with N-benzyl- and N-(p-nitrobenzyl)iminodiacetic acids. Structural and vibrational spectroscopic characterization and DFT study. <i>Inorganica Chimica Acta</i> , 2016, 453, 95-103.	2.4	7
64	Interval ridge regression (IRR) as a fast and robust method for quantitative prediction and variable selection applied to edible oil adulteration. <i>Talanta</i> , 2016, 150, 37-45.	5.5	7
65	Copper(II) perchlorate complexes with N-arylalkyliminodiacetamide ligands: X-ray structural, vibrational spectroscopic, DFT and thermogravimetric studies. <i>Inorganica Chimica Acta</i> , 2017, 462, 57-63.	2.4	7
66	Impact of Coordinated Pseudohalide Ions and Picolinamide on Supramolecular Synthons in Selected Zinc and Cadmium Complexes. <i>Acta Chimica Slovenica</i> , 2015, 62, 328-336.	0.6	7
67	A one-dimensional Cd <sup>II</sup> coordination polymer: <i>catena</i> -poly[cadmium(II)-bis(1/4-6-methylpicolinato)]. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2007, 63, m389-m391.	0.4	6
68	trans-Bis(azido- $\mu$ -N)bis(pyridine-2-carboxamide- $\mu$ -2N1,O2)nickel(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2007, 63, m507-m509.	0.4	6
69	Vibrational spectroscopic and DFT calculation studies of cobalt(II) complexes with 3-hydroxypicolinic acid. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2013, 101, 273-282.	3.9	6
70	Structural diversity of the Ag coordination sphere in complexes of silver(I) nitrate with 3-halopyridine. Characterization of the complexes in solution and in the solid state. <i>Polyhedron</i> , 2016, 109, 166-175.	2.2	6
71	Anion influence on polymerization of Cd(II) picolinamide complexes. <i>Inorganica Chimica Acta</i> , 2018, 478, 32-43.	2.4	6
72	Preparation and Characterization of Copper(II) and Nickel(II) Complexes with N-Benzyliminodiacetamide Derivatives. <i>Australian Journal of Chemistry</i> , 2016, 69, 896.	0.9	6

#	ARTICLE	IF	CITATIONS
73	Bis(7-amino-1,3-benzothiazole- $\hat{\nu}$ N <sub>3</sub> )dichlorozinc(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003, 59, m4-m6.	0.4	5
74	Complexes of zinc(II) halides with N-benzyl- and N-p-tolyl-2-oxo-1-naphthylideneamine – preparation and characterization. <i>Polyhedron</i> , 2004, 23, 1293-1302.	2.2	5
75	A new coordination mode of 6-methylnicotinic acid intrans-tetraaquabis(6-methylpyridine-3-carboxylato- $\hat{\nu}$ O)cobalt(II) tetrahydrate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2007, 63, m615-m617.	0.4	5
76	Vibrational Spectroscopic Characterization and DFT Study of Palladium(II) Complexes with N-Benzyliminodiacetic Acid Derivatives. <i>Australian Journal of Chemistry</i> , 2016, 69, 1285.	0.9	5
77	Palladium(II) complexes with N-alkyliminodiacetic acid derivatives: Preparation, structural and spectroscopic studies. <i>Inorganica Chimica Acta</i> , 2017, 462, 64-74.	2.4	5
78	Electrochemical Characteristics of 4-oxo-4 <i>H</i> -pyran-2-carboxylic Acid (Chelidonic Acid) and some of its Metal Complexes. <i>Electroanalysis</i> , 2017, 29, 538-547.	2.9	5
79	A Design of Mercury(II) Coordination Polymers with Pyridinedicarboxylic Acids: Structural, Spectroscopic and Thermal Studies. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2018, 28, 2080-2089.	3.7	5
80	Ternary palladium(II) complexes with N-benzyliminodiacetic acid derivatives and 2,2'-bipyridine: Preparation, thermogravimetric, vibrational spectroscopic, DFT, NMR studies and biological activity in vitro. <i>Inorganica Chimica Acta</i> , 2021, 516, 120131.	2.4	5
81	Mercuration of thiophene-2-carboxylic acid, $\gamma$ -thiophenylethanoic acid and 3-(2-thienyl)alanine: preparation and spectral characterization. <i>Applied Organometallic Chemistry</i> , 2000, 14, 598-603.	3.5	4
82	Diiodo(3,4,5,6-tetrahydropyrimidinium-2-thiolato-S)mercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2001, 57, 409-411.	0.4	4
83	1,3-Benzothiazole-6-carboxamidinium chloride dihydrate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003, 59, o190-o191.	0.4	4
84	Polymeric $\frac{1}{4}$ -bromo- $\frac{1}{4}$ -pyridine-3-carboxylato- $\hat{\nu}$ 3O, $\hat{\nu}$ 2:N-mercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003, 59, m165-m167.	0.4	4
85	Preparation, structural and thermal characterization of cobalt(II) and copper(II) complexes with 3-hydroxypicolinamide. <i>Transition Metal Chemistry</i> , 2011, 36, 65-71.	1.4	4
86	The role of non-covalent intermolecular interactions on the diversity of crystal packing in supramolecular dihalopyridine-silver nitrate complexes. <i>CrystEngComm</i> , 2020, 22, 7962-7974.	2.6	4
87	Self-Assembly of Cobalt(II) Coordination Polymers with Differently Halosubstituted Nicotinate Ligands and 4,4'-bipyridine – The Effect of the Halosubstituent Positions on Polymer Types. <i>European Journal of Inorganic Chemistry</i> , 2021, 2021, 1470-1480.	2.0	4
88	cis-Bis(azido- $\hat{\nu}$ N)bis(pyridine-2-carboxamide- $\hat{\nu}$ 2N1,O)nickel(II). <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2008, 64, m311-m312.	0.2	4
89	Aquabromo(6-carboxypyridine-2-carboxylato-O,N, $\hat{\nu}$ 2)mercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2002, 58, m39-m40.	0.4	3
90	2-Furancarboxanilide. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2004, 60, o631-o633.	0.2	3

#	ARTICLE	IF	CITATIONS
91	4-Amino-N-isopropylbenzamidinium chloride ethanol solvate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2005, 61, o98-o100.	0.4	3
92	Iodo(picolinato- $\lambda^2$ N,O)(picolinic acid- $\lambda^2$ N,O)mercury(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2006, 62, m272-m274.	0.4	3
93	(Nitrato- $\lambda^2$ O)bis(pyridine-2-carboxamide- $\lambda^2$ N1,O)mercury(II) nitrate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2007, 63, m557-m559.	0.4	3
94	Functionalization of plasmonic metamaterials utilizing metal-organic framework thin films. <i>Physica Scripta</i> , 2012, T149, 014051.	2.5	3
95	Impact of amide-amide hydrogen bonding on the stability of two nicotinamide complexes of silver(I). <i>Mineralogy and Petrology</i> , 2013, 107, 341-348.	1.1	3
96	New coordination modes of iminodiacetamide type ligands in palladium(II) complexes: crystallographic and DFT studies. <i>Structural Chemistry</i> , 2018, 29, 195-206.	2.0	3
97	catena-Poly[[2-methyl-1,3-benzothiazole-N)mercury(II)]-di- $\lambda^4$ -chloro]. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2001, 57, 20-21.	0.4	3
98	N-Benzyl-2-methylfuran-3-thiocarboxanilide. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2004, 60, o637-o639.	0.2	2
99	Bis{1-[N-(2-methoxyphenyl)iminomethyl]naphthalen-2(1H)-onato- $\lambda^3$ O,N, $\lambda^2$ }-zinc(II) ethanol solvate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2005, 61, m351-m352.	0.4	2
100	Copper(II) complexes with N-alkyliminodiacetamide ligands. Preparation, structural, spectroscopic and DFT studies and biological evaluation. <i>Inorganica Chimica Acta</i> , 2018, 471, 521-529.	2.4	2
101	Hydrogen-Bonded Frameworks of Mercury(II) Complexes with Pyridinedicarboxylic Acids. <i>Australian Journal of Chemistry</i> , 2018, 71, 455.	0.9	2
102	The Solvent Effect on Composition and Dimensionality of Mercury(II) Complexes with Picolinic Acid. <i>Molecules</i> , 2021, 26, 5002.	3.8	2
103	Anion-assisted supramolecular assemblies of zinc(II) complexes with isonicotinamide. <i>Heliyon</i> , 2022, 8, e09943.	3.2	2
104	On the interaction between mercury(II) salts and 3-methylpyrazoline-5-one. The first crystal structure of a cyclic organomercuric compound with a dimercurated methylenic carbon atom. <i>Structural Chemistry</i> , 2006, 17, 195-200.	2.0	1
105	10.1007/s10973-006-8123-z. <i>Journal of Thermal Analysis and Calorimetry</i> , 2007, 88, 489-493.	3.6	1
106	Synthesis, Spectral and Thermal Characterization of Bis(pyridine-3-carboxy-2-thiolato-S)mercury(II). Crystal Structure of Dibromobis(3-methoxycarbonylpyridine-2-thiolato-S)mercury(II). <i>Croatica Chemica Acta</i> , 2012, 85, 515-524.	0.4	1
107	Role of the group 12 metals in tuning the structural motif: From complexes to halometallate salts of 3-carbamoyl-1-methylpyridinium. <i>Polyhedron</i> , 2013, 52, 689-698.	2.2	1
108	Preparation and NMR spectroscopic study of palladium(II) complexes with N-arylalkyliminodiacetamide derivatives. <i>Journal of Molecular Structure</i> , 2016, 1122, 192-197.	3.6	1

#	ARTICLE	IF	CITATIONS
109	Structural and electrochemical properties of two novel CdX <sub>2</sub> (X=Br, I) picolinamide complexes. <i>Polyhedron</i> , 2021, 194, 114906.	2.2	1
110	trans-Bis(dimethyl sulfoxide- <i>l</i> -O)bis(3-oxo-3,4-dihydroquinoxaline-2-carboxylato- <i>l</i> -N1,O2)copper(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2007, 63, m181-m183.	0.4	0