

Masaki Ohsawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2946541/publications.pdf>

Version: 2024-02-01

71
papers

1,561
citations

304368

22
h-index

329751

37
g-index

72
all docs

72
docs citations

72
times ranked

2537
citing authors

#	ARTICLE	IF	CITATIONS
1	Rapid Increase in Estimated Number of Persons with Atrial Fibrillation in Japan: An Analysis from National Surveys on Cardiovascular Diseases in 1980, 1990 and 2000. <i>Journal of Epidemiology</i> , 2005, 15, 194-196.	1.1	115
2	CRP levels are elevated in smokers but unrelated to the number of cigarettes and are decreased by long-term smoking cessation in male smokers. <i>Preventive Medicine</i> , 2005, 41, 651-656.	1.6	102
3	Associations of ikigai as a positive psychological factor with all-cause mortality and cause-specific mortality among middle-aged and elderly Japanese people: Findings from the Japan Collaborative Cohort Study. <i>Journal of Psychosomatic Research</i> , 2009, 67, 67-75.	1.2	89
4	Association between Serum C-reactive Protein Levels and Microalbuminuria: A Population-Based Cross-Sectional Study in Northern Iwate, Japan. <i>Internal Medicine</i> , 2004, 43, 919-925.	0.3	79
5	Serum C-reactive protein levels can be used to predict future ischemic stroke and mortality in Japanese men from the general population. <i>Atherosclerosis</i> , 2009, 204, 234-238.	0.4	65
6	Serum Low-Density Lipoprotein to High-Density Lipoprotein Ratio as a Predictor of Future Acute Myocardial Infarction Among Men in a 2.7-Year Cohort Study of a Japanese Northern Rural Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011, 18, 89-98.	0.9	63
7	Mortality Risk Attributable to Atrial Fibrillation in Middle-Aged and Elderly People in the Japanese General Population Nineteen-Year Follow-up in NIPPON DATA80. <i>Circulation Journal</i> , 2007, 71, 814-819.	0.7	61
8	Influence of the Great East Japan Earthquake and Tsunami 2011 on Occurrence of Cerebrovascular Diseases in Iwate, Japan. <i>Stroke</i> , 2013, 44, 1518-1524.	1.0	58
9	Serum selenium levels are inversely associated with death risk among hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3331-3338.	0.4	56
10	Low-grade albuminuria and incidence of cardiovascular disease and all-cause mortality in nondiabetic and normotensive individuals. <i>Journal of Hypertension</i> , 2016, 34, 506-512.	0.3	53
11	Comparison of low-density lipoprotein cholesterol concentrations measured by a direct homogeneous assay and by the Friedewald formula in a large community population. <i>Clinica Chimica Acta</i> , 2010, 411, 1774-1780.	0.5	42
12	Factors Related to Tooth Loss Among Community-Dwelling Middle-aged and Elderly Japanese Men. <i>Journal of Epidemiology</i> , 2013, 23, 301-306.	1.1	42
13	A Community Based Epidemiological and Clinical Study of Hospitalization of Patients With Congestive Heart Failure in Northern Iwate, Japan. <i>Circulation Journal</i> , 2007, 71, 455-459.	0.7	41
14	Cardiovascular risk factors in the Japanese northeastern rural population. <i>International Journal of Cardiology</i> , 2009, 137, 226-235.	0.8	40
15	Predictive value of plasma B-type natriuretic peptide for ischemic stroke: A community-based longitudinal study. <i>Atherosclerosis</i> , 2009, 207, 298-303.	0.4	36
16	Dietary intake of n-3 polyunsaturated fatty acids is inversely associated with CRP levels, especially among male smokers. <i>Atherosclerosis</i> , 2008, 201, 184-191.	0.4	29
17	Poor self-rated health is significantly associated with elevated C-reactive protein levels in women, but not in men, in the Japanese general population. <i>Journal of Psychosomatic Research</i> , 2012, 73, 225-231.	1.2	28
18	Comparison of Predictability of Future Cardiovascular Events Between Chronic Kidney Disease (CKD) Stage Based on CKD Epidemiology Collaboration Equation and That Based on Modification of Diet in Renal Disease Equation in the Japanese General Population. <i>Circulation Journal</i> , 2013, 77, 1315-1325.	0.7	25

#	ARTICLE	IF	CITATIONS
19	Plasma B-Type Natriuretic Peptide Level and Cardiovascular Events in Chronic Kidney Disease in a Community-Based Population. <i>Circulation Journal</i> , 2010, 74, 792-797.	0.7	24
20	Prehypertension Subtype With Elevated C-Reactive Protein: Risk of Ischemic Stroke in a General Japanese Population. <i>American Journal of Hypertension</i> , 2010, 23, 1108-1113.	1.0	24
21	Seropositivity for Anti-HCV Core Antigen is Independently Associated With Increased All-Cause, Cardiovascular, and Liver Disease-Related Mortality in Hemodialysis Patients. <i>Journal of Epidemiology</i> , 2011, 21, 491-499.	1.1	24
22	Serum Selenium Levels in Hemodialysis Patients Are Significantly Lower than Those in Healthy Controls. <i>Blood Purification</i> , 2011, 32, 43-47.	0.9	24
23	Plasma B-Type Natriuretic Peptide Is Useful for Cardiovascular Risk Assessment in Community-Based Diabetes Subjects. <i>International Heart Journal</i> , 2012, 53, 176-181.	0.5	22
24	Associations of number of teeth with risks for all-cause mortality and cause-specific mortality in middle-aged and elderly men in the northern part of Japan: the Iwate-KENCO study. <i>Community Dentistry and Oral Epidemiology</i> , 2014, 42, 358-365.	0.9	22
25	Genetic polymorphism of 5,10-methylenetetrahydrofolate increases risk of myocardial infarction and is correlated to elevated levels of homocysteine in the Japanese general population. <i>Coronary Artery Disease</i> , 2000, 11, 47-51.	0.3	20
26	Incidence Rate of Cerebrovascular Diseases in Northern Japan Determined from the Iwate Stroke Registry with an Inventory Survey System. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013, 22, e317-e322.	0.7	19
27	Standardized Prevalence Ratios for Chronic Hepatitis C Virus Infection Among Adult Japanese Hemodialysis Patients. <i>Journal of Epidemiology</i> , 2010, 20, 30-39.	1.1	18
28	Cardiovascular Risk Factors in Hemodialysis Patients: Results from Baseline Data of Kaleidoscopic Approaches to Patients with End-stage Renal Disease Study. <i>Journal of Epidemiology</i> , 2005, 15, 96-105.	1.1	17
29	Serum ionic fluoride concentrations are related to renal function and menopause status but not to age in a Japanese general population. <i>Clinica Chimica Acta</i> , 2010, 411, 263-266.	0.5	16
30	High risks of all-cause and cardiovascular deaths in apparently healthy middle-aged people with preserved glomerular filtration rate and albuminuria: A prospective cohort study. <i>International Journal of Cardiology</i> , 2013, 170, 167-172.	0.8	16
31	Plasma B-type Natriuretic Peptide as a Predictor of Cardiovascular Events in Subjects with Atrial Fibrillation: A Community-Based Study. <i>PLoS ONE</i> , 2013, 8, e81243.	1.1	16
32	Renal function trajectory over time and adverse clinical outcomes. <i>Clinical and Experimental Nephrology</i> , 2016, 20, 379-393.	0.7	16
33	Associations of marital status with mortality from all causes and mortality from cardiovascular disease in Japanese haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 1013-1020.	0.4	15
34	Post-operative Constrictive Pericarditis Complicated with Lymphocytopenia and Hypoglobulinemia. <i>Internal Medicine</i> , 2004, 43, 811-815.	0.3	13
35	Predictive Value of Lipoprotein Indices for Residual Risk of Acute Myocardial Infarction and Sudden Death in Men With Low-Density Lipoprotein Cholesterol Levels ≤ 120 mg/dl. <i>American Journal of Cardiology</i> , 2013, 112, 1063-1068.	0.7	13
36	Relative and absolute risks of all-cause and cause-specific deaths attributable to atrial fibrillation in middle-aged and elderly community dwellers. <i>International Journal of Cardiology</i> , 2015, 184, 692-698.	0.8	13

#	ARTICLE	IF	CITATIONS
37	Ability of B-Type Natriuretic Peptide Testing to Predict Cardioembolic Stroke in the General Population—Comparisons With C-Reactive Protein and Urinary Albumin. <i>Circulation Journal</i> , 2018, 82, 1017-1025.	0.7	13
38	Sex-Specific Threshold Levels of Plasma B-Type Natriuretic Peptide for Prediction of Cardiovascular Event Risk in a Japanese Population Initially Free of Cardiovascular Disease. <i>American Journal of Cardiology</i> , 2011, 108, 1564-1569.	0.7	12
39	Prognostic Value of Electrocardiographic Left Ventricular Hypertrophy on Cardiovascular Risk in a Non-Hypertensive Community-Based Population. <i>American Journal of Hypertension</i> , 2018, 31, 895-901.	1.0	12
40	Poor self-rated health predicts the incidence of functional disability in elderly community dwellers in Japan: a prospective cohort study. <i>BMC Geriatrics</i> , 2020, 20, 328.	1.1	12
41	Gender-specific risk stratification with plasma B-type natriuretic peptide for future onset of congestive heart failure and mortality in the Japanese general population. <i>International Journal of Cardiology</i> , 2010, 143, 124-129.	0.8	11
42	Influence of mild-to-moderate alcohol consumption on cardiovascular diseases in men from the general population. <i>Atherosclerosis</i> , 2012, 224, 222-227.	0.4	11
43	Comparison between urine albumin-to-creatinine ratio and urine protein dipstick testing for prevalence and ability to predict the risk for chronic kidney disease in the general population (Iwate-KENCO study): a prospective community-based cohort study. <i>BMC Nephrology</i> , 2016, 17, 46.	0.8	11
44	Risk of stroke and heart failure attributable to atrial fibrillation in middle-aged and elderly people: Results from a five-year prospective cohort study of Japanese community dwellers. <i>Journal of Epidemiology</i> , 2017, 27, 360-367.	1.1	11
45	Bradycardia is associated with future cardiovascular diseases and death in men from the general population. <i>Atherosclerosis</i> , 2014, 236, 116-120.	0.4	9
46	Standardized Prevalence Ratios for Atrial Fibrillation in Adult Dialysis Patients in Japan. <i>Journal of Epidemiology</i> , 2016, 26, 272-276.	1.1	9
47	Burden of high blood pressure as a contributing factor to stroke in the Japanese community-based diabetic population. <i>Hypertension Research</i> , 2018, 41, 531-538.	1.5	9
48	The urine albumin-creatinine ratio is a predictor for incident long-term care in a general population. <i>PLoS ONE</i> , 2018, 13, e0195013.	1.1	8
49	Serum C-Reactive Protein Levels and Death and Cardiovascular Events in Mild to Moderate Chronic Kidney Disease. <i>International Heart Journal</i> , 2011, 52, 180-184.	0.5	7
50	Extent of Flood Damage Increased Cerebrovascular Disease Incidences in Iwate Prefecture after the Great East Japan Earthquake and Tsunami of 2011. <i>Cerebrovascular Diseases</i> , 2014, 37, 451-459.	0.8	7
51	Concordance of CKD stages in estimation by the CKD-EPI equation and estimation by the MDRD equation in the Japanese general population: The Iwate KENCO Study. <i>International Journal of Cardiology</i> , 2013, 165, 377-379.	0.8	6
52	Ten-Year Cerebrovascular Disease Trend Occurrence by Population-based Stroke Registry in an Aging Japan Local Prefecture. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2020, 29, 104580.	0.7	6
53	Conflicting effect of alcohol on cardiovascular risk: a clue to understand the different etiologies of coronary artery disease, stroke and peripheral artery disease. <i>Hypertension Research</i> , 2013, 36, 16-18.	1.5	5
54	Cardiovascular Risk Stratification With Plasma B-Type Natriuretic Peptide Levels in a Community-Based Hypertensive Cohort. <i>American Journal of Cardiology</i> , 2014, 113, 682-686.	0.7	5

#	ARTICLE	IF	CITATIONS
55	Slightly Elevated Serum Ionic Fluoride Levels Inhibit Insulin Secretion and Increase Glucose Levels in a General Japanese Population: a Cross-sectional Study. <i>Biological Trace Element Research</i> , 2021, 199, 2819-2825.	1.9	5
56	Age-specific incidence rates of atrial fibrillation and risk factors for the future development of atrial fibrillation in the Japanese general population. <i>Journal of Cardiology</i> , 2021, 77, 88-92.	0.8	5
57	Effect of Pre-Myocardial Infarction Angina Pectoris on Post-Myocardial Infarction Arrhythmias After Reperfusion Therapy. <i>American Journal of Cardiology</i> , 2006, 97, 1157-1161.	0.7	4
58	Trend in the Prevalence of Atrial Fibrillation During the Past 15 Years in Iwate (Northeastern Area of) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.7	4
59	Additional prognostic value of electrocardiographic left ventricular hypertrophy in traditional cardiovascular risk assessments in chronic kidney disease. <i>Journal of Hypertension</i> , 2020, 38, 1149-1157.	0.3	4
60	A Case of Sick Sinus Syndrome that Developed Torsades de Pointes, Pacing Failure and Sensing Failure During Administration of Bepridil. <i>International Heart Journal</i> , 2003, 44, 783-788.	0.6	4
61	The Potential of a Stroke Registry Using Diagnosis Procedure Combination Data from All Hospitals in a Japanese Prefecture. <i>Cerebrovascular Diseases</i> , 2022, 51, 447-452.	0.8	4
62	Association between Milk Intake and Incident Stroke among Japanese Community Dwellers: The Iwate-KENCO Study. <i>Nutrients</i> , 2021, 13, 3781.	1.7	2
63	Occurrence of Cerebrovascular Diseases Decreased after the Great East Japan Earthquake and Tsunami of 2011. <i>Cerebrovascular Diseases Extra</i> , 2020, 10, 105-115.	0.5	1
64	Age-Specific Incidence Rates of Acute Coronary Syndrome in Japan—A Special Gift From a Collaboration of Cardiologists and Epidemiologists—. <i>Circulation Journal</i> , 2021, 85, 1731-1734.	0.7	1
65	Corrigendum to “Gender-specific risk stratification with plasma B-type natriuretic peptide for future onset of congestive heart failure and mortality in the Japanese general population” <i>Int J Cardiol</i> 143 (2010) 124–129. <i>International Journal of Cardiology</i> , 2011, 146, 485.	0.8	0
66	Tightrope walking: Achieving the best possible balance between better model fit and accurate prediction: Response to letter (IJC-D-14-00417). <i>International Journal of Cardiology</i> , 2014, 174, 791-794.	0.8	0
67	Usefulness of risk grading system using albuminuria for predicting cardiovascular events and all-cause death in chronic kidney disease: A population-based prospective cohort study in Japan. <i>International Journal of Cardiology</i> , 2014, 175, 576-577.	0.8	0
68	Low educational level increases functional disability risk subsequent to heart failure in Japan: On behalf of the Iwate KENCO study group. <i>PLoS ONE</i> , 2021, 16, e0253017.	1.1	0
69	Utility of urinary albumin excretion as an index for stratifying the residual cardiovascular risk in patients undergoing antihypertensive agents treatment. <i>Journal of Hypertension</i> , 2021, Publish Ahead of Print, 2431-2438.	0.3	0
70	Abstract P331: Unfavorable Social Factors Increased the Risk of Future Disability Subsequent to Heart Failure in Elderly Japanese Community Dwellers: The Iwate KENCO Study Group. <i>Circulation</i> , 2019, 139, .	1.6	0
71	Abstract P401: Female Gender is a Significant Risk Factor for Functional Disability After Hemorrhagic Stroke but Not Ischemic Stroke Among Japanese Community-Dwelling Elderly Individuals. <i>Circulation</i> , 2019, 139, .	1.6	0