

Thomas M Krieg

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/295070/thomas-m-krieg-publications-by-year.pdf>

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75 papers	5,944 citations	39 h-index	77 g-index
86 ext. papers	6,775 ext. citations	6 avg, IF	5.41 L-index

#	Paper	IF	Citations
75	Primary systemic sclerosis heart involvement: A systematic literature review and preliminary data-driven, consensus-based WSF/HFA definition.. <i>Journal of Scleroderma and Related Disorders</i> , 2022 , 7, 24-32	2.3	5
74	A Practical Approach to the Management of Digital Ulcers in Patients With Systemic Sclerosis: A Narrative Review. <i>JAMA Dermatology</i> , 2021 , 157, 851-858	5.1	3
73	A story of fibers and stress: Matrix-embedded signals for fibroblast activation in the skin. <i>Wound Repair and Regeneration</i> , 2021 , 29, 515-530	3.6	5
72	Systemic sclerosis and the COVID-19 pandemic: World Scleroderma Foundation preliminary advice for patient management. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 724-726	2.4	38
71	Role of collagen XII in skin homeostasis and repair. <i>Matrix Biology</i> , 2020 , 94, 57-76	11.4	10
70	Scleroderma Renal Crisis: Risk Factors for an Increasingly Rare Organ Complication. <i>Journal of Rheumatology</i> , 2020 , 47, 241-248	4.1	11
69	Proteomic Analysis of Human Scleroderma Fibroblasts Response to Transforming Growth Factor- β <i>Proteomics - Clinical Applications</i> , 2019 , 13, e1800069	3.1	3
68	Dual role of laminin-511 in regulating melanocyte migration and differentiation. <i>Matrix Biology</i> , 2019 , 80, 59-71	11.4	7
67	New developments on skin fibrosis - Essential signals emanating from the extracellular matrix for the control of myofibroblasts. <i>Matrix Biology</i> , 2018 , 68-69, 522-532	11.4	43
66	TGFB1 is secreted through an unconventional pathway dependent on the autophagic machinery and cytoskeletal regulators. <i>Autophagy</i> , 2018 , 14, 465-486	10.2	47
65	Defining Skin Ulcers in Systemic Sclerosis: Systematic Literature Review and Proposed World Scleroderma Foundation (WSF) definition. <i>Journal of Scleroderma and Related Disorders</i> , 2017 , 2, 115-120	2.3	49
64	Deep Proteome Profiling Reveals Common Prevalence of MZB1-Positive Plasma B Cells in Human Lung and Skin Fibrosis. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2017 , 196, 1298-1310	10.2	60
63	Pathophysiological Mechanisms in Sclerosing Skin Diseases. <i>Frontiers in Medicine</i> , 2017 , 4, 120	4.9	5
62	Role of Integrins $\alpha 1$ and $\alpha 2$ in Wound and Tumor Angiogenesis in Mice. <i>American Journal of Pathology</i> , 2016 , 186, 3011-3027	5.8	23
61	Laminin β in the keratinocyte basement membrane is required for epidermal-dermal intercommunication. <i>Matrix Biology</i> , 2016 , 56, 24-41	11.4	20
60	Localized scleroderma: a review. <i>Journal of Scleroderma and Related Disorders</i> , 2016 , 1, 286-297	2.3	1
59	COMP-assisted collagen secretion--a novel intracellular function required for fibrosis. <i>Journal of Cell Science</i> , 2016 , 129, 706-16	5.3	43

58	Elucidating the burden of recurrent and chronic digital ulcers in systemic sclerosis: long-term results from the DUO Registry. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1770-6	2.4	49
57	Clinical characteristics and predictors of gangrene in patients with systemic sclerosis and digital ulcers in the Digital Ulcer Outcome Registry: a prospective, observational cohort. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1736-40	2.4	26
56	Deletion of the epidermis derived laminin α chain leads to defects in the regulation of late hair morphogenesis. <i>Matrix Biology</i> , 2016 , 56, 42-56	11.4	12
55	Effect of Macitentan on the Development of New Ischemic Digital Ulcers in Patients With Systemic Sclerosis: DUAL-1 and DUAL-2 Randomized Clinical Trials. <i>JAMA - Journal of the American Medical Association</i> , 2016 , 315, 1975-88	27.4	74
54	Interleukin-4 Receptor β Signaling in Myeloid Cells Controls Collagen Fibril Assembly in Skin Repair. <i>Immunity</i> , 2015 , 43, 803-16	32.3	182
53	Epidermal RelA specifically restricts contact allergen-induced inflammation and apoptosis in skin. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 2541-2550	4.3	5
52	Randomized standard-of-care-controlled trial of a silica gel fibre matrix in the treatment of chronic venous leg ulcers. <i>European Journal of Dermatology</i> , 2014 , 24, 210-6	0.8	10
51	Molecular and cellular basis of scleroderma. <i>Journal of Molecular Medicine</i> , 2014 , 92, 913-24	5.5	27
50	Genetic ablation of mast cells redefines the role of mast cells in skin wound healing and bleomycin-induced fibrosis. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 2005-2015	4.3	53
49	Role of integrin signalling through integrin-linked kinase in skin physiology and pathology. <i>Experimental Dermatology</i> , 2014 , 23, 453-6	4	7
48	Stabilization of integrin-linked kinase by the Hsp90-CHIP axis impacts cellular force generation, migration and the fibrotic response. <i>EMBO Journal</i> , 2013 , 32, 1409-24	13	51
47	Enhanced deposition of cartilage oligomeric matrix protein is a common feature in fibrotic skin pathologies. <i>Matrix Biology</i> , 2013 , 32, 325-31	11.4	37
46	Pharmacology and rationale for imatinib in the treatment of scleroderma. <i>Journal of Experimental Pharmacology</i> , 2013 , 5, 15-22	3	10
45	Collagen XII and XIV, new partners of cartilage oligomeric matrix protein in the skin extracellular matrix suprastructure. <i>Journal of Biological Chemistry</i> , 2012 , 287, 22549-59	5.4	92
44	Vascular endothelial insulin/IGF-1 signaling controls skin wound vascularization. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 421, 197-202	3.4	32
43	Biomarkers for skin involvement and fibrotic activity in scleroderma. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2012 , 26, 267-76	4.6	26
42	Dwarfism in mice lacking collagen-binding integrins α 1 and β 1 is caused by severely diminished IGF-1 levels. <i>Journal of Biological Chemistry</i> , 2012 , 287, 6431-40	5.4	23
41	Frequency of disease-associated and other nuclear autoantibodies in patients of the German Network for Systemic Scleroderma: correlation with characteristic clinical features. <i>Arthritis Research and Therapy</i> , 2011 , 13, R172	5.7	91

40	Combination therapy with an endothelin-1 receptor antagonist (bosentan) and a phosphodiesterase V inhibitor (sildenafil) for the management of severe digital ulcerations in systemic sclerosis. <i>Journal of the American Academy of Dermatology</i> , 2011 , 65, e102-e104	4.5	14
39	The extracellular matrix of the dermis: flexible structures with dynamic functions. <i>Experimental Dermatology</i> , 2011 , 20, 689-95	4	51
38	Registries in systemic sclerosis: a worldwide experience. <i>Rheumatology</i> , 2011 , 50, 60-8	3.9	31
37	Pivotal role for alpha1-antichymotrypsin in skin repair. <i>Journal of Biological Chemistry</i> , 2011 , 286, 28889-28901	3.4	32
36	Scleroderma: from pathophysiology to novel therapeutic approaches. <i>Experimental Dermatology</i> , 2010 , 19, 393-400	4	32
35	Defective granulation tissue formation in mice with specific ablation of integrin-linked kinase in fibroblasts - role of TGF β levels and RhoA activity. <i>Journal of Cell Science</i> , 2010 , 123, 3872-83	5.3	42
34	Differential roles of macrophages in diverse phases of skin repair. <i>Journal of Immunology</i> , 2010 , 184, 3964-77	5.3	742
33	Cell-matrix interactions in dermal repair and scarring. <i>Fibrogenesis and Tissue Repair</i> , 2010 , 3, 4		119
32	Scleroderma. <i>New England Journal of Medicine</i> , 2009 , 360, 1989-2003	59.2	1041
31	Alternative proteolytic processing of hepatocyte growth factor during wound repair. <i>American Journal of Pathology</i> , 2009 , 174, 2116-28	5.8	51
30	Role of tyrosine phosphatase SHP-1 in the mechanism of endorepellin angiostatic activity. <i>Blood</i> , 2009 , 114, 4897-906	2.2	59
29	Integrin alpha2beta1 is the required receptor for endorepellin angiostatic activity. <i>Journal of Biological Chemistry</i> , 2008 , 283, 2335-43	5.4	90
28	Fibroblast matrix interactions in tissue repair and fibrosis. <i>Experimental Dermatology</i> , 2008 , 17, 877-879	4	1
27	Integrin alpha2beta1 is required for regulation of murine wound angiogenesis but is dispensable for reepithelialization. <i>Journal of Investigative Dermatology</i> , 2007 , 127, 467-78	4.3	97
26	Scleroderma -- news to tell. <i>Archives of Dermatological Research</i> , 2007 , 299, 139-44	3.3	2
25	New developments in fibroblast and myofibroblast biology: implications for fibrosis and scleroderma. <i>Current Rheumatology Reports</i> , 2007 , 9, 136-43	4.9	132
24	Fibrosis in connective tissue disease: the role of the myofibroblast and fibroblast-epithelial cell interactions. <i>Arthritis Research and Therapy</i> , 2007 , 9 Suppl 2, S4	5.7	95
23	Interactions of primary fibroblasts and keratinocytes with extracellular matrix proteins: contribution of alpha2beta1 integrin. <i>Journal of Cell Science</i> , 2006 , 119, 1886-95	5.3	84

22	Mechanical tension and integrin alpha 2 beta 1 regulate fibroblast functions. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2006 , 11, 66-72	1.1	105
21	Absence of autoantibodies against correctly folded recombinant fibrillin-1 protein in systemic sclerosis patients. <i>Arthritis Research and Therapy</i> , 2005 , 7, R1221-6	5.7	22
20	Dissecting the roles of endothelin, TGF-beta and GM-CSF on myofibroblast differentiation by keratinocytes. <i>Thrombosis and Haemostasis</i> , 2004 , 92, 262-74	7	77
19	T cell-specific inactivation of the interleukin 10 gene in mice results in enhanced T cell responses but normal innate responses to lipopolysaccharide or skin irritation. <i>Journal of Experimental Medicine</i> , 2004 , 200, 1289-97	16.6	244
18	Keratin 14 Cre transgenic mice authenticate keratin 14 as an oocyte-expressed protein. <i>Genesis</i> , 2004 , 38, 176-81	1.9	115
17	Myofibroblast differentiation is induced in keratinocyte-fibroblast co-cultures and is antagonistically regulated by endogenous transforming growth factor-beta and interleukin-1. <i>American Journal of Pathology</i> , 2004 , 164, 2055-66	5.8	151
16	High expression and autoinduction of monocyte chemoattractant protein-1 in scleroderma fibroblasts. <i>European Journal of Immunology</i> , 2001 , 31, 2936-41	6.1	64
15	Apoptosis in v-myc-transfected MSU-1.1 fibroblasts is induced by cell-matrix contact and differs from that of normal dermal fibroblasts. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2001 , 37, 606-12	2.6	
14	Fibroblasts in mechanically stressed collagen lattices assume a "synthetic" phenotype. <i>Journal of Biological Chemistry</i> , 2001 , 276, 36575-85	5.4	277
13	Expression and proteolysis of vascular endothelial growth factor is increased in chronic wounds. <i>Journal of Investigative Dermatology</i> , 2000 , 115, 12-8	4.3	234
12	Bleomycin increases steady-state levels of type I collagen, fibronectin and decorin mRNAs in human skin fibroblasts. <i>Archives of Dermatological Research</i> , 2000 , 292, 556-61	3.3	41
11	Interactions of fibroblasts with the extracellular matrix: implications for the understanding of fibrosis. <i>Seminars in Immunopathology</i> , 1999 , 21, 415-29		23
10	Ultraviolet-B induction of interstitial collagenase and stromelysin-1 occurs in human dermal fibroblasts via an autocrine interleukin-6-dependent loop. <i>FEBS Letters</i> , 1999 , 449, 36-40	3.8	37
9	Highly sensitive DNA typing for detecting tumors transmitted by transplantation. <i>Transplant International</i> , 1998 , 11, 382-386	3	13
8	Mutations in the hair cortex keratin hHb6 cause the inherited hair disease monilethrix. <i>Nature Genetics</i> , 1997 , 16, 372-4	36.3	159
7	Differential regulation of transcription and transcript stability of pro-alpha 1(I) collagen and fibronectin in activated fibroblasts derived from patients with systemic scleroderma. <i>Biochemical Journal</i> , 1996 , 315 (Pt 2), 549-54	3.8	54
6	In vitro reconstituted skin as a tool for biology, pharmacology and therapy: a review. <i>Wound Repair and Regeneration</i> , 1995 , 3, 248-57	3.6	16
5	UVA-induced autocrine stimulation of fibroblast-derived collagenase/MMP-1 by interrelated loops of interleukin-1 and interleukin-6. <i>Photochemistry and Photobiology</i> , 1994 , 59, 550-6	3.6	227

4	Downregulation of collagen synthesis in fibroblasts within three-dimensional collagen lattices involves transcriptional and posttranscriptional mechanisms. <i>FEBS Letters</i> , 1993 , 318, 129-33	3.8	83
3	Interleukin-6 expression by fibroblasts grown in three-dimensional gel cultures. <i>FEBS Letters</i> , 1992 , 298, 229-32	3.8	15
2	Altered regulation of collagen metabolism in scleroderma fibroblasts grown within three-dimensional collagen gels. <i>Experimental Dermatology</i> , 1992 , 1, 185-90	4	39
1	Ultrastructure and composition of connective tissue in hyalinosi cutis et mucosae skin. <i>Journal of Investigative Dermatology</i> , 1984 , 82, 252-8	4.3	45