

Zixi Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2958034/publications.pdf>

Version: 2024-02-01

17
papers

259
citations

933447

10
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

140
citing authors

#	ARTICLE	IF	CITATIONS
1	The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in East Asia. <i>Science Advances</i> , 2021, 7, .	10.3	51
2	Liquidambar (Altingiaceae) and associated insect herbivory from the Miocene of southeastern China. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 497, 11-24.	2.3	26
3	A New Species of Ginkgo with Male Cones and Pollen Grains in situ from the Middle Jurassic of Eastern Xinjiang, China. <i>Acta Geologica Sinica</i> , 2017, 91, 9-21.	1.4	25
4	New fossil leaves and fruits of Lauraceae from the Middle Miocene of Fujian, southeastern China differentiated using a cluster analysis. <i>Historical Biology</i> , 2019, 31, 581-599.	1.4	22
5	<i>Elatides sandaolingensis</i> n. sp. (Cupressaceae sensu lato) – A new fossil conifer with cones from the Middle Jurassic of Xinjiang, northwestern China. <i>Palaeoworld</i> , 2016, 25, 239-250.	1.1	19
6	Two samaras of Rhamnaceae from the middle Miocene of southeast China. <i>Review of Palaeobotany and Palynology</i> , 2018, 259, 112-122.	1.5	17
7	A new species of <i>Garcinia</i> (Clusiaceae) from the middle Miocene of Fujian, China, and a phytogeographic analysis. <i>Geological Journal</i> , 2019, 54, 1317-1330.	1.3	17
8	A new species of <i>Ormosia</i> (Leguminosae) from the middle Miocene of Fujian, Southeast China and its biogeography. <i>Review of Palaeobotany and Palynology</i> , 2019, 270, 40-47.	1.5	16
9	A new cordaitan pollen cone and pollen grains in situ from the Early Permian of Hexi Corridor and its geotectonic significance. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2016, 463, 261-274.	2.3	14
10	A new <i>Choerospondias</i> (Anacardiaceae) endocarp from the middle Miocene of Southeast China and its paleoecological implications. <i>Review of Palaeobotany and Palynology</i> , 2020, 283, 104312.	1.5	13
11	A new <i>Cercis</i> (Leguminosae) from the middle Miocene of Fujian, China. <i>Historical Biology</i> , 2022, 34, 94-101.	1.4	8
12	First fossil record of <i>Canarium</i> (Burseraceae) from the middle Miocene of Fujian, southeastern China and its paleoecological implications. <i>Palaeoworld</i> , 2023, 32, 607-617.	1.1	8
13	Identification of two new species of <i>Meliolinites</i> associated with Lauraceae leaves from the middle Miocene of Fujian, China. <i>Mycologia</i> , 2017, 109, 1-14.	1.9	6
14	The diversity and paleoenvironmental significance of <i>Calophyllum</i> (Clusiaceae) from the Miocene of southeastern China. <i>Historical Biology</i> , 0, , 1-15.	1.4	5
15	A new species of <i>Cyathocarpus</i> with in situ spores from the lower Permian of Gansu, northwestern China. <i>Historical Biology</i> , 2019, 31, 824-835.	1.4	4
16	The geological significance of a new species of <i>Coniopteris</i> from the Middle Jurassic of northwestern China. <i>Historical Biology</i> , 2020, 32, 267-280.	1.4	4
17	<i>Lygodium</i> with in situ spores from the middle Miocene of Southeast China and its paleoclimatic implication. <i>Review of Palaeobotany and Palynology</i> , 2022, 296, 104545.	1.5	4