

Yanxiong Ke

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2959658/publications.pdf>

Version: 2024-02-01

102
papers

3,376
citations

201674

27
h-index

149698

56
g-index

109
all docs

109
docs citations

109
times ranked

3605
citing authors

#	ARTICLE	IF	CITATIONS
1	An Interweaving MOF with High Hydrogen Uptake. <i>Journal of the American Chemical Society</i> , 2006, 128, 3896-3897.	13.7	567
2	[[Tp) ₈ (H ₂ O) ₆ Cu ₁₆ Fe ₁₈ (CN) ₂₄] ⁴⁺ : A Cyanide-Bridged Face-Centered-Cubic Cluster with Single-Molecule-Magnet Behavior. <i>Angewandte Chemie - International Edition</i> , 2004, 43, 5940-5943.	13.8	219
3	Temperature-dependent supramolecular stereoisomerism in porous copper coordination networks based on a designed carboxylate ligand. <i>Chemical Communications</i> , 2005, , 5447.	4.1	176
4	Synthesis, characterization, and photoluminescence of isostructural Mn, Co, and Zn MOFs having a diamondoid structure with large tetrahedral cages and high thermal stability. <i>Chemical Communications</i> , 2005, , 2663.	4.1	161
5	Construction of Open Metal-Organic Frameworks Based on Predesigned Carboxylate Isomers: From Achiral to Chiral Nets. <i>Chemistry - A European Journal</i> , 2006, 12, 3768-3776.	3.3	151
6	Construction of Robust Open Metal-Organic Frameworks with Chiral Channels and Permanent Porosity. <i>Inorganic Chemistry</i> , 2007, 46, 2725-2734.	4.0	149
7	Synthesis and Structure of Cuboctahedral and Anticuboctahedral Cages Containing 12 Quadruply Bonded Dimolybdenum Units. <i>Inorganic Chemistry</i> , 2005, 44, 4154-4156.	4.0	101
8	Stability and Porosity Enhancement through Concurrent Ligand Extension and Secondary Building Unit Stabilization. <i>Inorganic Chemistry</i> , 2006, 45, 7566-7568.	4.0	90
9	Supramolecular Isomerism in Honeycomb Metal-Organic Frameworks Driven by CH- π Interactions: Homochiral Crystallization from an Achiral Ligand through Chiral Inducement. <i>Inorganic Chemistry</i> , 2010, 49, 8650-8652.	4.0	87
10	A Novel Analgesic Isolated from a Traditional Chinese Medicine. <i>Current Biology</i> , 2014, 24, 117-123.	3.9	85
11	(10,3)-a Noninterpenetrated Network Built from a Piedfort Ligand Pair. <i>Inorganic Chemistry</i> , 2006, 45, 1897-1899.	4.0	75
12	A novel click chitoooligosaccharide for hydrophilic interaction liquid chromatography. <i>Chemical Communications</i> , 2009, , 6973.	4.1	74
13	Fabrication of well-ordered macroporous active carbon with a microporous framework. <i>Journal of Materials Chemistry</i> , 2001, 11, 1975-1977.	6.7	70
14	Selective enrichment of glycopeptides/phosphopeptides using porous titania microspheres. <i>Chemical Communications</i> , 2010, 46, 5488.	4.1	61
15	Facile synthesis of titania-zirconia monodisperse microspheres and application for phosphopeptides enrichment. <i>Chemical Communications</i> , 2009, , 2929.	4.1	59
16	Separation of carbohydrates using hydrophilic interaction liquid chromatography. <i>Carbohydrate Research</i> , 2013, 379, 13-17.	2.3	58
17	Combination of off-line two-dimensional hydrophilic interaction liquid chromatography for polar fraction and two-dimensional hydrophilic interaction liquid chromatography-reversed-phase liquid chromatography for medium-polar fraction in a traditional Chinese medicine. <i>Journal of Chromatography A</i> , 2012, 1224, 61-69.	3.7	53
18	Purification of amide alkaloids from Piper longum L. using preparative two-dimensional normal-phase liquid chromatography-reversed-phase liquid chromatography. <i>Analyst, The</i> , 2013, 138, 3313.	3.5	50

#	ARTICLE	IF	CITATIONS
19	Preparation and chromatographic evaluation of a newly designed steviol glycoside modified-silica stationary phase in hydrophilic interaction liquid chromatography and reversed phase liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1388, 110-118.	3.7	48
20	Two-step templating route to macroporous or hollow sphere oxides. <i>Journal of Materials Chemistry</i> , 2001, 11, 2930-2933.	6.7	43
21	Novel reversed-phase high-performance liquid chromatography stationary phase with oligo(ethylene) Tj ETQq1 1 0.784314 rgBT /Over	3.7	41
22	1D zigzag chain vs. 1D helical chain: the role of the supramolecular interactions on the formation of chiral architecture. <i>CrystEngComm</i> , 2010, 12, 337-340.	2.6	38
23	“Click dipeptide” A novel stationary phase applied in two-dimensional liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 8623-8629.	3.7	37
24	Selective separation of structure-related alkaloids in <i>Rhizoma coptidis</i> with “click”-binaphthyl stationary phase and their structural elucidation with liquid chromatography-mass spectrometry. <i>Analyst, The</i> , 2011, 136, 4357.	3.5	36
25	Alkaloids analysis using off-line two-dimensional supercritical fluid chromatography – ultra-high performance liquid chromatography. <i>Analyst, The</i> , 2014, 139, 3577-3587.	3.5	36
26	Rapid and simultaneous analysis of sesquiterpene pyridine alkaloids from <i>Tripterygium wilfordii</i> Hook. f. Using supercritical fluid chromatography-diode array detector-tandem mass spectrometry. <i>Journal of Supercritical Fluids</i> , 2015, 104, 85-93.	3.2	31
27	Selective separation of flavonoid glycosides in <i>Dalbergia odorifera</i> by matrix solid-phase dispersion using titania. <i>Journal of Separation Science</i> , 2011, 34, 1347-1354.	2.5	28
28	A polyacrylamide-based silica stationary phase for the separation of carbohydrates using alcohols as the weak eluent in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1524, 153-159.	3.7	26
29	Hydrothermal synthesis, structures and spectroscopy of 2D lanthanide coordination polymers built from helical chains: [Ln ₂ (BDC) ₃ (H ₂ O) ₂] _n (Ln=Sm, 1; Ln=Eu, 2; BDC=1,3-benzenedicarboxylate). <i>Journal of Molecular Structure</i> , 2005, 734, 7-13.	3.6	25
30	Pore expansion of highly monodisperse phenylene-bridged organosilica spheres for chromatographic application. <i>Talanta</i> , 2010, 81, 824-830.	5.5	24
31	A dextran-bonded stationary phase for saccharide separation. <i>Journal of Chromatography A</i> , 2014, 1345, 57-67.	3.7	23
32	Rapid purification of diastereoisomers from <i>Piper kadsura</i> using supercritical fluid chromatography with chiral stationary phases. <i>Journal of Chromatography A</i> , 2017, 1509, 141-146.	3.7	22
33	Large-pore monodispersed mesoporous silica spheres: synthesis and application in HPLC. <i>Chemical Communications</i> , 2009, , 1085.	4.1	20
34	Retention mechanism and enrichment of glycopeptides on titanium dioxide. <i>Analytical Methods</i> , 2013, 5, 7072.	2.7	19
35	Novel chiral ionic liquids stationary phases for the enantiomer separation of chiral acid by high-performance liquid chromatography. <i>Chirality</i> , 2018, 30, 670-679.	2.6	18
36	A 3D silver coordination polymer with novel topology structure. <i>Solid State Sciences</i> , 2004, 6, 753-755.	3.2	17

#	ARTICLE	IF	CITATIONS
37	Enantioselective recognition ability of peptoids with $\hat{\pm}$ -chiral, aromatic side chains. <i>Analyst</i> , 2011, 136, 4409.	3.5	17
38	The development of an evaluation method for capture columns used in two-dimensional liquid chromatography. <i>Analytica Chimica Acta</i> , 2011, 706, 184-190.	5.4	16
39	Synthesis and evaluation of a maltose $\hat{\text{C}}$ -bonded silica gel stationary phase for hydrophilic interaction chromatography and its application in <i>Ginkgo Biloba</i> extract separation in two-dimensional systems. <i>Journal of Separation Science</i> , 2016, 39, 3339-3347.	2.5	16
40	Synthesis of mesostructured tin oxide with neutral surfactant as a template in aqueous media. <i>Materials Letters</i> , 2003, 57, 2679-2681.	2.6	15
41	A novel ionic-bonded cellulose stationary phase for saccharide separation. <i>Journal of Chromatography A</i> , 2013, 1291, 56-63.	3.7	15
42	Efficient preparative separation of 6 $\hat{\text{C}}$ -(4 $\hat{\text{C}}$ -aminophenyl)-5 $\hat{\text{C}}$ -methyl-4, 5 $\hat{\text{C}}$ -dihydro-3(2H)-pyridazinone enantiomers on polysaccharide $\hat{\text{C}}$ -based stationary phases in polar organic solvent chromatography and supercritical fluid chromatography. <i>Journal of Separation Science</i> , 2019, 42, 2482-2490.	2.5	15
43			

#	ARTICLE	IF	CITATIONS
55	Chemical separation and characterization of complex samples with herbal medicine. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 124, 115775.	11.4	11
56	Evaluation of a series of phenyl-type stationary phases in supercritical fluid chromatography with the linear solvation energy relationship model and its application to the separation of phenolic compounds. <i>Journal of Chromatography A</i> , 2020, 1614, 460700.	3.7	11
57	Synthesis and structure of 2D- and 3D- inorganic-organic coordination polymers: based on Ag-hmt subunit. <i>Crystal Research and Technology</i> , 2004, 39, 89-93.	1.3	10
58	Investigation of peptoid chiral stationary phases varied in absolute configuration. <i>Journal of Chromatography A</i> , 2013, 1281, 155-159.	3.7	10
59	Improvement of chiral stationary phases based on cinchona alkaloids bonded to crown ethers by chiral modification. <i>Journal of Separation Science</i> , 2015, 38, 3884-3890.	2.5	10
60	Preparation and evaluation of novel chiral stationary phases based on quinine derivatives comprising crown ether moieties. <i>Journal of Separation Science</i> , 2015, 38, 205-210.	2.5	10
61	Separation of Piper kadsura Using Preparative Supercritical Fluid Chromatography Combined with Preparative Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2018, 81, 1181-1187.	1.3	10
62	Separation of Ketorolac enantiomers on polysaccharide-based chiral stationary phases using a polar organic mobile phase. <i>Separation Science Plus</i> , 2018, 1, 351-358.	0.6	10
63	Synthesis and Crystal Structure of a New Copper(II) PMIDA Compound (H ₄ PMIDA=H ₂ O ₃ PCH ₂ N(CH ₂ CO ₂ H) ₂). <i>Structural Chemistry</i> , 2004, 15, 207-210.	2.0	9
64	Synthesis and structure characterization of three-coordinate silver (I) and seven-coordinate cobalt (II) coordination polymers with 4-pyridylthioacetate. <i>Journal of Molecular Structure</i> , 2004, 689, 75-80.	3.6	9
65	Synthesis and structure of the three-dimensional coordination polymer [Ag ₃ hmt ₃ (1/3 -btc)]·5H ₂ O. <i>Crystal Research and Technology</i> , 2006, 41, 98-102.	1.3	9
66	Preparation of a stationary phase with a triazine ring embedded group for reversed phase high-performance LC. <i>Journal of Separation Science</i> , 2010, 33, 2998-3004.	2.5	9
67	Preparation of a binaphthyl stationary phase and its application for separation of anthraquinones from <i>Rheum palmatum</i> L.. <i>Journal of Separation Science</i> , 2011, 34, 1133-1140.	2.5	9
68	Synthesis and structure of a 3-rings antimony germanate: Sb ₂ Ge ₃ O ₉ . <i>Solid State Sciences</i> , 2002, 4, 803-806.	3.2	8
69	Novel chiral stationary phases based on peptoid combining a quinine/quinidine moiety through a C9-position carbamate group. <i>Journal of Separation Science</i> , 2014, 37, 934-943.	2.5	8
70	Ultrasonic-Assisted Sol-Gel Synthesis of Core-Shell Silica Particles for High-Performance Liquid Chromatography. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2020, 30, 859-868.	3.7	8
71	Investigation of brush-type chiral stationary phases based on O,O'-bis(aryldiaroyl tartardiamide and O,O'-bis(arylcarbamoyl) tartardiamide. <i>Journal of Separation Science</i> , 2012, 35, 351-358.	2.5	7
72	Study of stereomeric peptoid chiral stationary phases containing different chiral side chains. <i>Journal of Chromatography A</i> , 2013, 1298, 152-156.	3.7	7

#	ARTICLE	IF	CITATIONS
73	Preparation of sub-2 μ m large-pore monodispersed mesoporous silica spheres using mixed templates and application in HPLC. <i>Microporous and Mesoporous Materials</i> , 2018, 265, 234-240.	4.4	7
74	Monodisperse core-shell silica particles as a high-performance liquid chromatography packing material: Facile in situ silica sol-gel synthesis. <i>Journal of Chromatography A</i> , 2020, 1625, 461282.	3.7	7
75	Synthesis of Al ³⁺ -doping-TiO ₂ monodisperse microspheres and their application for phosphopeptides and glycopeptides enrichment. <i>Talanta</i> , 2021, 223, 121715.	5.5	7
76	Evaluation of α -click-binaphthyl chiral stationary phases by liquid chromatography. <i>Chirality</i> , 2012, 24, 391-399.	2.6	6
77	Synthesis of Large-Pore Silica Microspheres Using Dodecylamine as a Catalyst, Template and Porogen Agent. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2019, 29, 1417-1421.	3.7	6
78	Novel chiral stationary phases based on 3,5-dimethyl phenylcarbamoylated β -cyclodextrin combining cinchona alkaloid moiety. <i>Chirality</i> , 2020, 32, 1080-1090.	2.6	6
79	Enantiomeric analysis of simendan on polysaccharide-based stationary phases by polar organic solvent chromatography. <i>Journal of Separation Science</i> , 2020, 43, 2097-2104.	2.5	6
80	Crystal structure of 2,8,14,20-tetranaphthylpyrogallol[4]arene. <i>Journal of Chemical Crystallography</i> , 2006, 36, 67-70.	1.1	5
81	Investigation of Peptoid Chiral Stationary Phases Terminated with <i>N</i> -substituted Phenylproline/leucine Amide. <i>Chinese Journal of Chemistry</i> , 2012, 30, 2791-2797.	4.9	5
82	Highly Efficient Separation of Methylated Peptides Utilizing Selective Complexation between Lysine and 18-Crown-6. <i>Analytical Chemistry</i> , 2020, 92, 15663-15670.	6.5	5
83	Regioselective and diastereodivergent organocatalytic asymmetric vinylogous Michael addition. <i>Organic Chemistry Frontiers</i> , 2021, 8, 4758-4766.	4.5	5
84	Pore size control of monodisperse mesoporous silica particles with alkyl imidazole ionic liquid templates for high performance liquid chromatography applications. <i>Colloids and Surfaces A: Physicochemical and Engineering Aspects</i> , 2022, 637, 128200.	4.7	5
85	Separation of minor steviol glycosides using hydrophilic interaction liquid chromatography (HILIC) and off-line two-dimensional reversed-phase liquid chromatography/HILIC methods. <i>Journal of Food Composition and Analysis</i> , 2022, 112, 104683.	3.9	5
86	Hydrothermal synthesis and structure of a molybdenum oxide (TMA) ₂ [Ni(H ₂ O) ₆][β -Mo ₈ O ₂₆]. <i>Solid State Sciences</i> , 2003, 5, 317-320.	3.2	4
87	HPLC and SFC enantioseparation of (\pm)-Corey lactone diol: Impact of the amylose tris(3,5-dimethylphenylcarbamate) coating amount on chiral preparation. <i>Chirality</i> , 2019, 31, 855-864.	2.6	4
88	Design, synthesis and evaluation of a series of alkylsiloxane-bonded stationary phases for expanded supercritical fluid chromatography separations. <i>Journal of Chromatography A</i> , 2019, 1593, 127-134.	3.7	4
89	A novel C ₂ symmetric chiral stationary phase with N-{(4-methylphenyl)sulfonyl}-leucine as chiral side chains. <i>Journal of Separation Science</i> , 2020, 43, 2338-2348.	2.5	4
90	Pseudomorphic synthesis of bimodal porous silica microspheres for size-exclusion chromatography of small molecules. <i>Journal of Chromatography A</i> , 2022, 1664, 462757.	3.7	4

#	ARTICLE	IF	CITATIONS
91	Self-assembly synthesis and structure of mono- and di-nuclear uranyl complex. <i>Crystal Research and Technology</i> , 2003, 38, 1004-1008.	1.3	3
92	A Three-dimensional Manganese(II) 1,2,4-Benzenetricarboxylate Hydroxide Framework with Mn ^{II} O Inorganic Sheets: Hydrothermal Synthesis and Crystal Structure. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2008, 63, 1339-1342.	0.7	3
93	Sequential enrichment of singly- and multiply-phosphorylated peptides with zwitterionic hydrophilic interaction chromatography material. <i>Journal of Chromatography A</i> , 2015, 1413, 47-59.	3.7	3
94	Pore size control of monodisperse silica particles by dual template sol-gel method. <i>Journal of Sol-Gel Science and Technology</i> , 2020, 94, 186-194.	2.4	3
95	A ternary eluent strategy to tune the peak shape of steviol glycosides in reversed-phase liquid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2021, 1173, 122673.	2.3	3
96	Enantioseparation of cloprostenol on the polysaccharide chiral stationary phase: Influence of the mobile phase on enantioselective adsorption. <i>Journal of Chromatography A</i> , 2021, 1653, 462413.	3.7	3
97	Synthesis and structure of a three-dimensional coordination polymer [Ag ₃ hmt ₃ (3-btc)] ₂ ·1/25H ₂ O. <i>Journal of Structural Chemistry</i> , 2004, 45, 541-546.	1.0	2
98	A subtraction fitting method for independent determination of enantioselective and nonselective adsorption isotherms based on the single-component isotherms in the framework of the two-site model. <i>Journal of Chromatography A</i> , 2020, 1632, 461608.	3.7	2
99	Synthesis of mesostructured lamellar lead sulfide in acidic media. <i>Journal of Materials Research</i> , 2003, 18, 549-551.	2.6	1
100	Adsorption mechanism of triterpenoid saponins in reversed-phase liquid chromatography and hydrophilic interaction liquid chromatography: Mogroside V as test substance. <i>Journal of Chromatography A</i> , 2020, 1620, 461010.	3.7	1
101	Hydrothermal Synthesis and Structure of a Molybdenum Oxide (TMA) ₂ [Ni(H ₂ O) ₆] [M ₂ -Mo ₈ O ₂₆]. <i>ChemInform</i> , 2003, 34, no.	0.0	0
102	This paper has been retracted. Template-Induced Formation of a 3D Zinc Metal-Organic Framework Possessing a Rare (10,3)-d Net. <i>Crystal Growth and Design</i> , 2008, .	3.0	0