Douglas K Bishop

List of Publications by Citations

Source: https://exaly.com/author-pdf/29606/douglas-k-bishop-publications-by-citations.pdf

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

25 3,942 37 39 h-index g-index citations papers 15.6 4,459 39 5.33 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
37	DMC1: a meiosis-specific yeast homolog of E. coli recA required for recombination, synaptonemal complex formation, and cell cycle progression. <i>Cell</i> , 1992 , 69, 439-56	56.2	1018
36	RecA homologs Dmc1 and Rad51 interact to form multiple nuclear complexes prior to meiotic chromosome synapsis. <i>Cell</i> , 1994 , 79, 1081-92	56.2	405
35	Early decision; meiotic crossover interference prior to stable strand exchange and synapsis. <i>Cell</i> , 2004 , 117, 9-15	56.2	288
34	A meiotic recombination checkpoint controlled by mitotic checkpoint genes. <i>Nature</i> , 1996 , 383, 840-3	50.4	284
33	Xrcc3 is required for assembly of Rad51 complexes in vivo. <i>Journal of Biological Chemistry</i> , 1998 , 273, 21482-8	5.4	223
32	Rad51 is an accessory factor for Dmc1-mediated joint molecule formation during meiosis. <i>Science</i> , 2012 , 337, 1222-5	33.3	194
31	Saccharomyces cerevisiae recA homologues RAD51 and DMC1 have both distinct and overlapping roles in meiotic recombination. <i>Genes To Cells</i> , 1997 , 2, 615-29	2.3	154
30	DNA strand exchange and RecA homologs in meiosis. <i>Cold Spring Harbor Perspectives in Biology</i> , 2014 , 7, a016659	10.2	132
29	Saccharomyces cerevisiae checkpoint genes MEC1, RAD17 and RAD24 are required for normal meiotic recombination partner choice. <i>Genetics</i> , 1999 , 153, 607-20	4	114
28	Saccharomyces cerevisiae Dmc1 protein promotes renaturation of single-strand DNA (ssDNA) and assimilation of ssDNA into homologous super-coiled duplex DNA. <i>Journal of Biological Chemistry</i> , 2001 , 276, 41906-12	5.4	109
27	Synthesis-dependent strand annealing in meiosis. <i>PLoS Biology</i> , 2007 , 5, e299	9.7	102
26	Meiotic crossover control by concerted action of Rad51-Dmc1 in homolog template bias and robust homeostatic regulation. <i>PLoS Genetics</i> , 2013 , 9, e1003978	6	98
25	A comparative analysis of Dmc1 and Rad51 nucleoprotein filaments. <i>Nucleic Acids Research</i> , 2008 , 36, 4057-66	20.1	84
24	High copy number suppression of the meiotic arrest caused by a dmc1 mutation: REC114 imposes an early recombination block and RAD54 promotes a DMC1-independent DSB repair pathway. <i>Genes To Cells</i> , 1999 , 4, 425-44	2.3	76
23	Gradual implementation of the meiotic recombination program via checkpoint pathways controlled by global DSB levels. <i>Molecular Cell</i> , 2015 , 57, 797-811	17.6	67
22	Swi2/Snf2-related translocases prevent accumulation of toxic Rad51 complexes during mitotic growth. <i>Molecular Cell</i> , 2010 , 39, 862-72	17.6	65
21	Crossover interference in Saccharomyces cerevisiae requires a TID1/RDH54- and DMC1-dependent pathway. <i>Genetics</i> , 2003 , 163, 1273-86	4	59

(2018-2015)

20	Small Rad51 and Dmc1 Complexes Often Co-occupy Both Ends of a Meiotic DNA Double Strand Break. <i>PLoS Genetics</i> , 2015 , 11, e1005653	6	57
19	Tid1/Rdh54 promotes dissociation of Dmc1 from nonrecombinogenic sites on meiotic chromatin. <i>Genes and Development</i> , 2006 , 20, 2593-604	12.6	55
18	Non-enzymatic roles of human RAD51 at stalled replication forks. <i>Nature Communications</i> , 2019 , 10, 44	10 7.4	48
17	The Mei5-Sae3 protein complex mediates Dmc1 activity in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2009 , 284, 11766-70	5.4	38
16	Nine novel conserved motifs in BRCA1 identified by the chicken orthologue. <i>Oncogene</i> , 2001 , 20, 4433-	89.2	36
15	RAD54 family translocases counter genotoxic effects of RAD51 in human tumor cells. <i>Nucleic Acids Research</i> , 2015 , 43, 3180-96	20.1	35
14	Caffeine impairs resection during DNA break repair by reducing the levels of nucleases Sae2 and Dna2. <i>Nucleic Acids Research</i> , 2015 , 43, 6889-901	20.1	35
13	The RAD51-stimulatory compound RS-1 can exploit the RAD51 overexpression that exists in cancer cells and tumors. <i>Cancer Research</i> , 2014 , 74, 3546-55	10.1	32
12	The third exon of the budding yeast meiotic recombination gene HOP2 is required for calcium-dependent and recombinase Dmc1-specific stimulation of homologous strand assimilation. <i>Journal of Biological Chemistry</i> , 2014 , 289, 18076-86	5.4	25
11	DNA damage response clamp 9-1-1 promotes assembly of ZMM proteins for formation of crossovers and synaptonemal complex. <i>Journal of Cell Science</i> , 2015 , 128, 1494-506	5.3	24
10	The ATPase activity of E. coli RecA prevents accumulation of toxic complexes formed by erroneous binding to undamaged double stranded DNA. <i>Nucleic Acids Research</i> , 2018 , 46, 9510-9523	20.1	16
9	Caffeine inhibits gene conversion by displacing Rad51 from ssDNA. <i>Nucleic Acids Research</i> , 2015 , 43, 690	02018	11
8	RPA resolves conflicting activities of accessory proteins during reconstitution of Dmc1-mediated meiotic recombination. <i>Nucleic Acids Research</i> , 2019 , 47, 747-761	20.1	11
7	Surface Spreading and Immunostaining of Yeast Chromosomes. <i>Journal of Visualized Experiments</i> , 2015 , e53081	1.6	10
6	Multiple mechanisms of meiotic recombination. <i>Cell</i> , 2006 , 127, 1095-7	56.2	8
5	Distinct Functions in Regulation of Meiotic Crossovers for DNA Damage Response Clamp Loader Rad24(Rad17) and Mec1(ATR) Kinase. <i>Genetics</i> , 2019 , 213, 1255-1269	4	7
4	A mutant form of Dmc1 that bypasses the requirement for accessory protein Mei5-Sae3 reveals independent activities of Mei5-Sae3 and Rad51 in Dmc1 filament stability. <i>PLoS Genetics</i> , 2019 , 15, e100	08217	3
3	Purification of Saccharomyces cerevisiae Homologous Recombination Proteins Dmc1 and Rdh54/Tid1 and a Fluorescent D-Loop Assay. <i>Methods in Enzymology</i> , 2018 , 600, 307-320	1.7	3

How strand exchange protein function benefits from ATP hydrolysis. *Current Opinion in Genetics and Development*, **2021**, 71, 120-128

4.9 1

Meiosis in Quarantine discussions lead to an action plan to increase diversity and inclusion within the genetics community. *PLoS Genetics*, **2021**, 17, e1009648

6