
Liliana F Devizzi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2979633/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Prognostic Scoring System for Primary CNS Lymphomas: The International Extranodal Lymphoma Study Group Experience. Journal of Clinical Oncology, 2003, 21, 266-272.	0.8	688
2	Peripheral T-cell lymphoma unspecified (PTCL-U): a new prognostic model from a retrospective multicentric clinical study. Blood, 2004, 103, 2474-2479.	0.6	549
3	Nongastric marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue. Blood, 2003, 101, 2489-2495.	0.6	494
4	Clinical activity of rituximab in extranodal marginal zone B-cell lymphoma of MALT type. Blood, 2003, 102, 2741-2745.	0.6	391
5	Primary diffuse large B-cell lymphoma of the breast: prognostic factors and outcomes of a study by the International Extranodal Lymphoma Study Group. Annals of Oncology, 2008, 19, 233-241.	0.6	203
6	Successful in vivo purging of CD34-containing peripheral blood harvests in mantle cell and indolent lymphoma: evidence for a role of both chemotherapy and rituximab infusion. Blood, 2000, 96, 864-869.	0.6	201
7	Final Results of the IELSG-19 Randomized Trial of Mucosa-Associated Lymphoid Tissue Lymphoma: Improved Event-Free and Progression-Free Survival With Rituximab Plus Chlorambucil Versus Either Chlorambucil or Rituximab Monotherapy. Journal of Clinical Oncology, 2017, 35, 1905-1912.	0.8	143
8	Relevance of intraocular involvement in the management of primary central nervous system lymphomas. Annals of Oncology, 2002, 13, 531-538.	0.6	120
9	Vaccination with autologous tumor-loaded dendritic cells induces clinical and immunologic responses in indolent B-cell lymphoma patients with relapsed and measurable disease: a pilot study. Blood, 2009, 113, 18-27.	0.6	99
10	High-Dose Yttrium-90–lbritumomab Tiuxetan With Tandem Stem-Cell Reinfusion: An Outpatient Preparative Regimen for Autologous Hematopoietic Cell Transplantation. Journal of Clinical Oncology, 2008, 26, 5175-5182.	0.8	68
11	Prolonged survival in poor-risk diffuse large B-cell lymphoma following front-line treatment with rituximab-supplemented, early-intensified chemotherapy with multiple autologous hematopoietic stem cell support: a multicenter study by GITIL (Gruppo Italiano Terapie Innovative nei Linfomi). Leukemia, 2007, 21, 1802-1811.	3.3	66
12	Clinical activity of bortezomib in relapsed/refractory MALT lymphomas: results of a phase II study of the International Extranodal Lymphoma Study Group (IELSG). Annals of Oncology, 2011, 22, 689-695.	0.6	64
13	Helicobacter pylori eradication as exclusive treatment for limited-stage gastric diffuse large B-cell lymphoma: results of a multicenter phase 2 trial. Blood, 2012, 120, 3858-3860.	0.6	58
14	The lymphocyte to monocyte ratio improves the IPlâ€risk definition of diffuse large Bâ€cell lymphoma when rituximab is added to chemotherapy. American Journal of Hematology, 2013, 88, 1062-1067.	2.0	54
15	Genetic and phenotypic attributes of splenic marginal zone lymphoma. Blood, 2022, 139, 732-747.	0.6	49
16	Clinical activity of everolimus in relapsed/refractory marginal zone <scp>B</scp> â€cell lymphomas: results of a phase <scp>II</scp> study of the <scp>I</scp> nternational <scp>E</scp> xtranodal <scp>L</scp> ymphoma <scp>S</scp> tudy <scp>G</scp> roup. British Journal of Haematology, 2014, 166, 69-76.	1.2	39
17	Ifosfamide and vinorelbine: an active regimen for patients with relapsed or refractory Hodgkin's disease. British Journal of Haematology, 1998, 103, 533-535.	1.2	37
18	Phase <scp>II</scp> study of sorafenib in patients with relapsed or refractory lymphoma. British Journal of Haematology, 2012, 158, 108-119.	1.2	36

LILIANA F DEVIZZI

#	Article	IF	CITATIONS
19	High response rate and manageable toxicity with an intensive, short-term chemotherapy programme for Burkitt's lymphoma in adults. British Journal of Haematology, 2004, 126, 815-820.	1.2	35
20	Addition of Rituximab to Involved-Field Radiation Therapy Prolongs Progression-free Survival in Stage I-II Follicular Lymphoma: Results of a Multicenter Study. International Journal of Radiation Oncology Biology Physics, 2016, 94, 783-791.	0.4	35
21	Long-term results of high-dose chemotherapy with autologous bone marrow or peripheral stem cell transplant as first salvage treatment for relapsed or refractory Hodgkin lymphoma: a single institution experience. Leukemia and Lymphoma, 2010, 51, 1251-1259.	0.6	34
22	CLIPI: a new prognostic index for indolent cutaneous B cell lymphoma proposed by the International Extranodal Lymphoma Study Group (IELSG 11). Annals of Hematology, 2011, 90, 401-408.	0.8	28
23	High-dose sequential chemotherapy and in vivo rituximab-purged stem cell autografting in mantle cell lymphoma: a 10-year update of the R-HDS regimen. Bone Marrow Transplantation, 2009, 43, 509-511.	1.3	25
24	Myeloablative doses of yttriumâ€90â€ibritumomab tiuxetan and the risk of secondary myelodysplasia/acute myelogenous leukemia. Cancer, 2011, 117, 5074-5084.	2.0	23
25	Early stage Hodgkin's disease: Ten-year results of a non-randomised study with radiotherapy alone or combined with MOPP. European Journal of Cancer, 1993, 29, 24-29.	1.3	22
26	Lenalidomide monotherapy in heavily pretreated patients with non-Hodgkin lymphoma: an Italian observational multicenter retrospective study in daily clinical practice. Leukemia and Lymphoma, 2015, 56, 1671-1676.	0.6	22
27	Dosimetry in Myeloablative 90Y-Labeled Ibritumomab Tiuxetan Therapy: Possibility of Increasing Administered Activity on the Base of Biological Effective Dose Evaluation. Preliminary Results. Cancer Biotherapy and Radiopharmaceuticals, 2007, 22, 113-120.	0.7	20
28	The modified International Prognostic Index can predict the outcome of localized primary intestinal lymphoma of both extranodal marginal zone B-cell and diffuse large B-cell histologies. British Journal of Haematology, 2002, 118, 218-228.	1.2	19
29	Absorbed dose and biologically effective dose in patients with high-risk non-Hodgkin's lymphoma treated with high-activity myeloablative 90Y-ibritumomab tiuxetan (Zevalin®). European Journal of Nuclear Medicine and Molecular Imaging, 2009, 36, 1745-1757.	3.3	19
30	Long-Term Results of Autologous Hematopoietic Stem-Cell Transplantation After High-Dose ⁹⁰ Y-Ibritumomab Tiuxetan for Patients With Poor-Risk Non-Hodgkin Lymphoma Not Eligible for High-Dose BEAM. Journal of Clinical Oncology, 2013, 31, 2974-2976.	0.8	14
31	Results of a randomized trial comparing high-dose chemotherapy plus Auto-SCT and R-FC in CLL at diagnosis. Bone Marrow Transplantation, 2014, 49, 485-491.	1.3	10
32	Lenalidomide in Pretreated Patients with Diffuse Large B ell Lymphoma: An Italian Observational Multicenter Retrospective Study in Daily Clinical Practice. Oncologist, 2019, 24, 1246-1252.	1.9	10
33	Extended-field radiotherapy in favorable stage IA–IIA Hodgkin's disease (Prognostic role of stage). International Journal of Radiation Oncology Biology Physics, 1994, 30, 813-819.	0.4	9
34	Efficacy and safety of high-dose chemotherapy with in vivo purged auto-SCT in relapsed follicular lymphoma: long-term follow-up. Bone Marrow Transplantation, 2010, 45, 1119-1120.	1.3	9
35	The efficacy of lenalidomide combination therapy in heavily pretreated non-Hodgkin lymphoma patients: an Italian observational, multicenter, retrospective study. Leukemia and Lymphoma, 2017, 58, 226-229.	0.6	6
36	Radioimmunotherapy and secondary leukemia: A case report. Leukemia Research, 2010, 34, e1-e4.	0.4	4

LILIANA F DEVIZZI

#	Article	IF	CITATIONS
37	High-dose chemotherapy followed by autologous transplantation may overcome the poor prognosis of diffuse large B-cell lymphoma patients with MYC/BCL2 co-expression. Blood Cancer Journal, 2016, 6, e491-e491.	2.8	4
38	Fifteen-year follow-up of relapsed indolent non-Hodgkin lymphoma patients vaccinated with tumor-loaded dendritic cells. , 2021, 9, e002240.		4
39	Study of the HLA-DPβ1 Locus by the Polymerase Chain Reaction Technique in Patients with Hodgkin's Disease. Tumori, 1993, 79, 133-136.	0.6	2
40	MALT lymphoma and Kaposi Sarcoma in an HIV-negative patient. American Journal of Hematology, 2010, 85, 815-817.	2.0	2
41	Successful second autologous engraftment after long duration storage of hematopoietic stem cells. Bone Marrow Transplantation, 2013, 48, 1480-1481.	1.3	2
42	Successful in vivo purging of CD34-containing peripheral blood harvests in mantle cell and indolent lymphoma: evidence for a role of both chemotherapy and rituximab infusion. Blood, 2000, 96, 864-869.	0.6	1
43	Lack of carboplatin activity in malignant lymphomas. European Journal of Cancer, 1996, 32, 2370-2371.	1.3	0
44	Lenalidomide Combination Therapy in Relapsed/Refractory Diffuse Large B Cell Lymphoma: The Italian Real-Life Experience. Clinical Lymphoma, Myeloma and Leukemia, 2019, 19, e321-e323.	0.2	0
45	Radioimmunotherapy in the Transplant Setting. , 2018, , 431-441.		0