

# Thomas Mentzel

## List of Publications by Citations

Source: <https://exaly.com/author-pdf/2982656/thomas-mentzel-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

76  
papers

4,695  
citations

38  
h-index

68  
g-index

90  
ext. papers

5,322  
ext. citations

4.2  
avg, IF

5.35  
L-index

#	Paper	IF	Citations
76	Myxofibrosarcoma. Clinicopathologic analysis of 75 cases with emphasis on the low-grade variant. <i>American Journal of Surgical Pathology</i> , 1996, 20, 391-405	6.7	444
75	Fibrosarcomatous ("high-grade") dermatofibrosarcoma protuberans: clinicopathologic and immunohistochemical study of a series of 41 cases with emphasis on prognostic significance. <i>American Journal of Surgical Pathology</i> , 1998, 22, 576-87	6.7	305
74	Cellular Benign Fibrous Histiocytoma. <i>American Journal of Surgical Pathology</i> , 1994, 18, 668-676	6.7	194
73	Postradiation cutaneous angiosarcoma after treatment of breast carcinoma is characterized by MYC amplification in contrast to atypical vascular lesions after radiotherapy and control cases: clinicopathological, immunohistochemical and molecular analysis of 66 cases. <i>Modern Pathology</i> , 2012, 25, 75-85	9.8	192
72	Solitary fibrous tumour: clinicopathological, immunohistochemical, and ultrastructural analysis of 12 cases arising in soft tissues, nasal cavity and nasopharynx, urinary bladder and prostate. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1997, 430, 445-53	5.1	171
71	Current treatment options in dermatofibrosarcoma protuberans. <i>Journal of Cancer Research and Clinical Oncology</i> , 2009, 135, 653-65	4.9	167
70	Pleomorphic rhabdomyosarcoma in adults: a clinicopathologic study of 38 cases with emphasis on morphologic variants and recent skeletal muscle-specific markers. <i>Modern Pathology</i> , 2001, 14, 595-603	9.8	165
69	Lipoblastoma and lipoblastomatosis: a clinicopathological study of 14 cases. <i>Histopathology</i> , 1993, 23, 527-33	7.3	142
68	Cutaneous clear cell sarcoma: a clinicopathologic, immunohistochemical, and molecular analysis of 12 cases emphasizing its distinction from dermal melanoma. <i>American Journal of Surgical Pathology</i> , 2010, 34, 216-22	6.7	132
67	Spindle cell liposarcoma, a hitherto unrecognized variant of liposarcoma. Analysis of six cases. <i>American Journal of Surgical Pathology</i> , 1994, 18, 913-21	6.7	131
66	Cellular angiofibroma: analysis of 25 cases emphasizing its relationship to spindle cell lipoma and mammary-type myofibroblastoma. <i>Modern Pathology</i> , 2011, 24, 82-9	9.8	123
65	Perineurioma (storiform perineurial fibroma): clinico-pathological analysis of four cases. <i>Histopathology</i> , 1994, 25, 261-7	7.3	107
64	Sclerosing, pseudovascular rhabdomyosarcoma in adults. Clinicopathological and immunohistochemical analysis of three cases. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2000, 436, 305-11	5.1	103
63	Cutaneous clear cell myomelanocytic tumour: a new member of the growing family of perivascular epithelioid cell tumours (PEComas). Clinicopathological and immunohistochemical analysis of seven cases. <i>Histopathology</i> , 2005, 46, 498-504	7.3	98
62	Myxoid dermatofibrosarcoma protuberans: clinicopathologic, immunohistochemical, and molecular analysis of eight cases. <i>American Journal of Dermatopathology</i> , 2007, 29, 443-8	0.9	93
61	Lipomatous tumours of soft tissues: an update. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1995, 427, 353-63	5.1	92
60	Hobnail hemangioma ("targetoid hemosiderotic hemangioma"): clinicopathologic and immunohistochemical analysis of 62 cases. <i>Journal of Cutaneous Pathology</i> , 1999, 26, 279-86	1.7	90

59	Primary cutaneous leiomyosarcoma: clinicopathological analysis of 36 cases. <i>Histopathology</i> , <b>2010</b> , 56, 251-62	7.3	87
58	Primary liposarcoma of the skin: a rare neoplasm with unusual high grade features. <i>American Journal of Dermatopathology</i> , <b>1998</b> , 20, 332-8	0.9	84
57	Granular cell change in smooth muscle tumours of skin and soft tissue. <i>Histopathology</i> , <b>1994</b> , 24, 223-31	7.3	83
56	Well-differentiated spindle cell liposarcoma (Atypical spindle cell lipomatous tumor) does not belong to the spectrum of atypical lipomatous tumor but has a close relationship to spindle cell lipoma: clinicopathologic, immunohistochemical, and molecular analysis of six cases. <i>Modern Pathology</i> , <b>2010</b> , 23, 729-36	9.8	82
55	TERT promoter mutations are frequent in atypical fibroxanthomas and pleomorphic dermal sarcomas. <i>Modern Pathology</i> , <b>2014</b> , 27, 502-8	9.8	81
54	Benign vascular proliferations in irradiated skin. <i>American Journal of Surgical Pathology</i> , <b>2002</b> , 26, 328-376.7	7	79
53	Targeted massively parallel sequencing of angiosarcomas reveals frequent activation of the mitogen activated protein kinase pathway. <i>Oncotarget</i> , <b>2015</b> , 6, 36041-52	3.3	71
52	Proliferating trichilemmal tumors: clinicopathologic evaluation is a guide to biologic behavior. <i>Journal of Cutaneous Pathology</i> , <b>2003</b> , 30, 492-8	1.7	71
51	Plaque-like CD34-positive dermal fibroma ("medallion-like dermal dendrocyte hamartoma"): clinicopathologic, immunohistochemical, and molecular analysis of 5 cases emphasizing its distinction from superficial, plaque-like dermatofibrosarcoma protuberans. <i>American Journal of Surgical Pathology</i> , <b>2010</b> , 34, 190-201	6.7	67
50	Spindle cell rhabdomyosarcoma in adults: clinicopathological and immunohistochemical analysis of seven new cases. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2006</b> , 449, 554-60	5.1	67
49	Benign fibrous histiocytoma (dermatofibroma) of the face: clinicopathologic and immunohistochemical study of 34 cases associated with an aggressive clinical course. <i>American Journal of Dermatopathology</i> , <b>2001</b> , 23, 419-26	0.9	66
48	Intradermal spindle cell/pleomorphic lipoma: a distinct subset. <i>American Journal of Dermatopathology</i> , <b>2000</b> , 22, 496-502	0.9	64
47	Dermatomyofibroma: clinicopathologic and immunohistochemical analysis of 56 cases and reappraisal of a rare and distinct cutaneous neoplasm. <i>American Journal of Dermatopathology</i> , <b>2009</b> , 31, 44-9	0.9	60
46	"Atypical" Pleomorphic Lipomatous Tumor: A Clinicopathologic, Immunohistochemical and Molecular Study of 21 Cases, Emphasizing its Relationship to Atypical Spindle Cell Lipomatous Tumor and Suggesting a Morphologic Spectrum (Atypical Spindle Cell/Pleomorphic Lipomatous Tumor). <i>American Journal of Surgical Pathology</i> , <b>2017</b> , 41, 1443-1455	6.7	58
45	The T1796A mutation of the BRAF gene is absent in Spitz nevi. <i>Journal of Cutaneous Pathology</i> , <b>2004</b> , 31, 266-70	1.7	56
44	Intraneurial angiosarcoma and angiosarcoma arising in benign and malignant peripheral nerve sheath tumours: clinicopathological and immunohistochemical analysis of four cases. <i>Histopathology</i> , <b>1999</b> , 35, 114-20	7.3	55
43	Atypical spindle cell lipoma: a clinicopathologic, immunohistochemical, and molecular study emphasizing its relationship to classical spindle cell lipoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2014</b> , 465, 97-108	5.1	49
42	Dedifferentiated myxoid liposarcoma: a clinicopathological study suggesting a closer relationship between myxoid and well-differentiated liposarcoma. <i>Histopathology</i> , <b>1997</b> , 30, 457-63	7.3	49

41	Dermatomyofibroma: additional observations on a distinctive cutaneous myofibroblastic tumour with emphasis on differential diagnosis. <i>British Journal of Dermatology</i> , <b>1993</b> , 129, 69-73	4	47
40	Ancient hematoma: a unifying concept for a post-traumatic lesion mimicking an aggressive soft tissue neoplasm. <i>Modern Pathology</i> , <b>1997</b> , 10, 334-40	9.8	42
39	Spindle cell predominant trichodiscoma: a fibrofolliculoma/trichodiscoma variant considered formerly to be a neurofollicular hamartoma: a clinicopathological and immunohistochemical analysis of 17 cases. <i>American Journal of Dermatopathology</i> , <b>2006</b> , 28, 1-8	0.9	40
38	Atypical fibroxanthoma and pleomorphic dermal sarcoma harbor frequent NOTCH1/2 and FAT1 mutations and similar DNA copy number alteration profiles. <i>Modern Pathology</i> , <b>2018</b> , 31, 418-428	9.8	38
37	Malignant dermatofibroma: clinicopathological, immunohistochemical, and molecular analysis of seven cases. <i>Modern Pathology</i> , <b>2013</b> , 26, 256-67	9.8	37
36	Atypical Fibroxanthoma Revisited. <i>Surgical Pathology Clinics</i> , <b>2017</b> , 10, 319-335	3.9	36
35	Sarcomas of the skin in the elderly. <i>Clinics in Dermatology</i> , <b>2011</b> , 29, 80-90	3	36
34	Histological and genetic evidence for a variant of superficial spreading melanoma composed predominantly of large nests. <i>Modern Pathology</i> , <b>2012</b> , 25, 838-45	9.8	31
33	Cellular benign fibrous histiocytoma. Clinicopathologic analysis of 74 cases of a distinctive variant of cutaneous fibrous histiocytoma with frequent recurrence. <i>American Journal of Surgical Pathology</i> , <b>1994</b> , 18, 668-76	6.7	31
32	Re: Infrequent SMARCB1/INI1 gene alteration in epithelioid sarcoma: a useful tool in distinguishing epithelioid sarcoma from malignant rhabdoid tumor: Direct evidence of mutational inactivation of SMARCB1/INI1 in epithelioid sarcoma. <i>Human Pathology</i> , <b>2009</b> , 40, 1361-2; author reply 1362-4	3.7	28
31	Biological continuum of benign, atypical, and malignant mesenchymal neoplasms - does it exist?. <i>Journal of Pathology</i> , <b>2000</b> , 190, 523-5	9.4	28
30	Genome-wide methylation profiling and copy number analysis in atypical fibroxanthomas and pleomorphic dermal sarcomas indicate a similar molecular phenotype. <i>Clinical Sarcoma Research</i> , <b>2019</b> , 9, 2	2.5	25
29	Myxoid epithelioid sarcoma: a diagnostic challenge. A report on six cases. <i>Histopathology</i> , <b>2010</b> , 57, 753-73	7.3	22
28	Haemorrhagic dermatomyofibroma (plaque-like dermal fibromatosis): clinicopathological and immunohistochemical analysis of three cases resembling plaque-stage Kaposi's sarcoma. <i>Histopathology</i> , <b>2003</b> , 42, 594-8	7.3	21
27	The spectrum of rare morphological variants of cutaneous epithelioid angiosarcoma. <i>Histopathology</i> , <b>2015</b> , 66, 856-63	7.3	20
26	Radiation-induced vascular lesions of the skin: an overview. <i>Advances in Anatomic Pathology</i> , <b>2013</b> , 20, 407-15	5.1	15
25	Mixed-type liposarcoma: clinicopathological, immunohistochemical, and molecular analysis of a case arising in deep soft tissues of the lower extremity. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2008</b> , 453, 197-201	5.1	15
24	Malignant mesenchymomas of soft tissue associated with numerous osteoclast-like giant cells mimicking the so-called giant cell variant of "malignant fibrous histiocytoma". <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>1994</b> , 424, 539-45	5.1	15

23	Reduced H3K27me3 expression in radiation-associated angiosarcoma of the breast. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2018</b> , 472, 361-368	5.1	12
22	S-100 protein expressing spindle cells in spindle cell lipoma: a diagnostic pitfall. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2016</b> , 469, 435-8	5.1	12
21	Fibroblastic connective tissue nevus: Clinicopathological and immunohistochemical study of 14 cases. <i>Journal of Cutaneous Pathology</i> , <b>2017</b> , 44, 827-834	1.7	11
20	S1 guidelines for dermatofibrosarcoma protuberans (DFSP) - update 2018. <i>JDDG - Journal of the German Society of Dermatology</i> , <b>2019</b> , 17, 663-668	1.2	10
19	Cutaneous mesenchymal tumours: an update. <i>Pathology</i> , <b>2014</b> , 46, 149-59	1.6	10
18	Gastrointestinal stromal tumour with skeinoid fibres and bidirectional immunohistochemical differentiation. <i>Histopathology</i> , <b>1996</b> , 29, 175-7	7.3	10
17	Leiomyosarcoma with prominent osteoclast-like giant cells. Analysis of eight cases closely mimicking the so-called giant cell variant of malignant fibrous histiocytoma. <i>American Journal of Surgical Pathology</i> , <b>1994</b> , 18, 258-65	6.7	10
16	An adult spindle cell rhabdomyosarcoma in the head and neck region with long-term survival: a case report. <i>Journal of Medical Case Reports</i> , <b>2014</b> , 8, 208	1.2	9
15	Intra-abdominal lymphangioma in a newborn. <i>Journal of Clinical Ultrasound</i> , <b>1998</b> , 26, 320-2	1	9
14	Metastasizing phyllodes tumour with malignant fibrous histiocytoma-like areas. <i>Histopathology</i> , <b>1991</b> , 19, 557-60	7.3	5
13	GNA14, GNA11, and GNAQ Mutations Are Frequent in Benign but Not Malignant Cutaneous Vascular Tumors. <i>Frontiers in Genetics</i> , <b>2021</b> , 12, 663272	4.5	5
12	S1-Leitlinie Dermatofibrosarcoma protuberans (DFSP) - Update 2018. <i>JDDG - Journal of the German Society of Dermatology</i> , <b>2019</b> , 17, 663-668	1.2	4
11	Braided pattern in a dermatofibrosarcoma protuberans: a potential mimicker of neural neoplasms. <i>American Journal of Dermatopathology</i> , <b>2014</b> , 36, 920-4	0.9	3
10	Malignant mesenchymal neoplasms of the dermis and subcutis mimicking benign lesions: a case-based review. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , <b>2017</b> , 471, 565-574	5.1	2
9	Molecular pathology diagnostics in cutaneous mesenchymal tumors. <i>JDDG - Journal of the German Society of Dermatology</i> , <b>2013</b> , 11 Suppl 4, 36-49	1.2	2
8	S1-guideline atypical fibroxanthoma (AFX) and pleomorphic dermal sarcoma (PDS).. <i>JDDG - Journal of the German Society of Dermatology</i> , <b>2022</b> ,	1.2	2
7	Myxoid pleomorphic liposarcoma-a clinicopathologic, immunohistochemical, molecular genetic and epigenetic study of 12 cases, suggesting a possible relationship with conventional pleomorphic liposarcoma. <i>Modern Pathology</i> , <b>2021</b> , 34, 2043-2049	9.8	2
6	Pulmonary leiomyosarcoma mimicking glomus tumor at first biopsy and surgically treated with isolated left main bronchus resection: rare clinical documentation. <i>Irish Journal of Medical Science</i> , <b>2013</b> , 182, 735-8	1.9	1

- 5 Molecular Diagnostics of Vascular Tumors of the Skin. *American Journal of Dermatopathology*, **2020**, 42, 799-810 0.9 1
- 4 Livid macules on the forehead after photodynamic therapy. *JDDG - Journal of the German Society of Dermatology*, **2015**, 13, 924-6 1.2
- 3 Molekularpathologische Diagnostik von mesenchymalen Tumoren der Haut. *JDDG - Journal of the German Society of Dermatology*, **2013**, 11, 37-49 1.2
- 2 Mesenchymale Tumoren der Haut **2015**, 1-76
- 1 Mesenchymale Tumoren der Haut **2016**, 737-796