Francesco Simone Ruggeri

List of Publications by Citations

 $\textbf{Source:} \ https://exaly.com/author-pdf/2985017/francesco-simone-ruggeri-publications-by-citations.pdf$

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75 2,641 papers citations

11 28 h-index

50 g-index

90 ext. papers

3,548 ext. citations

10.4 avg, IF 5.15 L-index

#	Paper	IF	Citations
75	FUS Phase Separation Is Modulated by a Molecular Chaperone and Methylation of Arginine Cation-Interactions. <i>Cell</i> , 2018 , 173, 720-734.e15	56.2	409
74	Infrared nanospectroscopy characterization of oligomeric and fibrillar aggregates during amyloid formation. <i>Nature Communications</i> , 2015 , 6, 7831	17.4	169
73	Measurement of intrinsic properties of amyloid fibrils by the peak force QNM method. <i>Nanoscale</i> , 2012 , 4, 4426-9	7.7	149
72	Cholesterol catalyses AB2 aggregation through a heterogeneous nucleation pathway in the presence of lipid membranes. <i>Nature Chemistry</i> , 2018 , 10, 673-683	17.6	126
71	The H50Q mutation enhances Esynuclein aggregation, secretion, and toxicity. <i>Journal of Biological Chemistry</i> , 2014 , 289, 21856-76	5.4	126
70	Influence of the Esheet content on the mechanical properties of aggregates during amyloid fibrillization. <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 2462-6	16.4	106
69	Nanoscale studies link amyloid maturity with polyglutamine diseases onset. <i>Scientific Reports</i> , 2016 , 6, 31155	4.9	93
68	Detecting nanoscale vibrations as signature of life. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 378-81	11.5	80
67	Different soluble aggregates of AB2 can give rise to cellular toxicity through different mechanisms. <i>Nature Communications</i> , 2019 , 10, 1541	17.4	71
66	Trodusquemine enhances Alaggregation but suppresses its toxicity by displacing oligomers from cell membranes. <i>Nature Communications</i> , 2019 , 10, 225	17.4	69
65	Nanoplasmonic mid-infrared biosensor for protein secondary structure detection. <i>Light: Science and Applications</i> , 2017 , 6, e17029	16.7	66
64	Silk micrococoons for protein stabilisation and molecular encapsulation. <i>Nature Communications</i> , 2017 , 8, 15902	17.4	65
63	AFM-Based Single Molecule Techniques: Unraveling the Amyloid Pathogenic Species. <i>Current Pharmaceutical Design</i> , 2016 , 22, 3950-70	3.3	58
62	Atomic force microscopy for single molecule characterisation of protein aggregation. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 664, 134-148	4.1	57
61	Determination of Polypeptide Conformation with Nanoscale Resolution in Water. <i>ACS Nano</i> , 2018 , 12, 6612-6619	16.7	52
60	Identification and nanomechanical characterization of the fundamental single-strand protofilaments of amyloid Esynuclein fibrils. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 7230-7235	11.5	51
59	Nanobodies raised against monomeric ?-synuclein inhibit fibril formation and destabilize toxic oligomeric species. <i>BMC Biology</i> , 2017 , 15, 57	7.3	46

58	Nanoscale spatially resolved infrared spectra from single microdroplets. <i>Lab on A Chip</i> , 2014 , 14, 1315-9	7.2	41
57	One-pot semisynthesis of exon 1 of the Huntingtin protein: new tools for elucidating the role of posttranslational modifications in the pathogenesis of Huntington's disease. <i>Angewandte Chemie - International Edition</i> , 2014 , 53, 1928-33	16.4	39
56	Mutant Exon1 Huntingtin Aggregation is Regulated by T3 Phosphorylation-Induced Structural Changes and Crosstalk between T3 Phosphorylation and Acetylation at K6. <i>Angewandte Chemie - International Edition</i> , 2017 , 56, 5202-5207	16.4	38
55	A critical concentration of N-terminal pyroglutamylated amyloid beta drives the misfolding of Ab1-42 into more toxic aggregates. <i>International Journal of Biochemistry and Cell Biology</i> , 2016 , 79, 261-	270	36
54	N-terminal Huntingtin (Htt) phosphorylation is a molecular switch regulating Htt aggregation, helical conformation, internalization, and nuclear targeting. <i>Journal of Biological Chemistry</i> , 2018 , 293, 18540-18558	5.4	36
53	Soluble aggregates present in cerebrospinal fluid change in size and mechanism of toxicity during Alzheimer's disease progression. <i>Acta Neuropathologica Communications</i> , 2019 , 7, 120	7.3	35
52	Single molecule secondary structure determination of proteins through infrared absorption nanospectroscopy. <i>Nature Communications</i> , 2020 , 11, 2945	17.4	34
51	Stabilization and Characterization of Cytotoxic AlDligomers Isolated from an Aggregation Reaction in the Presence of Zinc Ions. <i>ACS Chemical Neuroscience</i> , 2018 , 9, 2959-2971	5.7	33
50	Biomolecular condensates undergo a generic shear-mediated liquid-to-solid transition. <i>Nature Nanotechnology</i> , 2020 , 15, 841-847	28.7	33
49	Identification of Oxidative Stress in Red Blood Cells with Nanoscale Chemical Resolution by Infrared Nanospectroscopy. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	32
48	Small-molecule sequestration of amyloid-las a drug discovery strategy for Alzheimer's disease. <i>Science Advances</i> , 2020 , 6,	14.3	28
47	Transthyretin Inhibits Primary and Secondary Nucleations of Amyloid-IPeptide Aggregation and Reduces the Toxicity of Its Oligomers. <i>Biomacromolecules</i> , 2020 , 21, 1112-1125	6.9	28
46	Stabilized tilted-octahedra halide perovskites inhibit local formation of performance-limiting phases <i>Science</i> , 2021 , 374, 1598-1605	33.3	28
45	Rational design of a conformation-specific antibody for the quantification of Albligomers. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 13509-13518	3 ^{11.5}	26
44	The Influence of Pathogenic Mutations in Esynuclein on Biophysical and Structural Characteristics of Amyloid Fibrils. <i>ACS Nano</i> , 2020 , 14, 5213-5222	16.7	24
43	Fabrication and Characterization of Reconstituted Silk Microgels for the Storage and Release of Small Molecules. <i>Macromolecular Rapid Communications</i> , 2019 , 40, e1800898	4.8	23
42	Water-Dispersible Polydopamine-Coated Nanofibers for Stimulation of Neuronal Growth and Adhesion. <i>Advanced Healthcare Materials</i> , 2018 , 7, e1701485	10.1	23
41	Trodusquemine displaces protein misfolded oligomers from cell membranes and abrogates their cytotoxicity through a generic mechanism. <i>Communications Biology</i> , 2020 , 3, 435	6.7	23

40	Molecular determinants of the interaction of EGCG with ordered and disordered proteins. <i>Biopolymers</i> , 2018 , 109, e23117	2.2	20
39	Concentration-dependent and surface-assisted self-assembly properties of a bioactive estrogen receptor Ederived peptide. <i>Journal of Peptide Science</i> , 2015 , 21, 95-104	2.1	20
38	Micro- and nanoscale hierarchical structure of core-shell protein microgels. <i>Journal of Materials Chemistry B</i> , 2016 , 4, 7989-7999	7.3	19
37	Solution fibre spinning technique for the fabrication of tuneable decellularised matrix-laden fibres and fibrous micromembranes. <i>Acta Biomaterialia</i> , 2018 , 78, 111-122	10.8	19
36	Microfluidic deposition for resolving single-molecule protein architecture and heterogeneity. <i>Nature Communications</i> , 2018 , 9, 3890	17.4	19
35	Influence of the Esheet Content on the Mechanical Properties of Aggregates during Amyloid Fibrillization. <i>Angewandte Chemie</i> , 2015 , 127, 2492-2496	3.6	15
34	Ultrathin Polydopamine Films with Phospholipid Nanodiscs Containing a Glycophorin A Domain. <i>Advanced Functional Materials</i> , 2020 , 30, 2000378	15.6	14
33	Amyloid single-cell cytotoxicity assays by nanomotion detection. <i>Cell Death Discovery</i> , 2017 , 3, 17053	6.9	13
32	Infrared nanospectroscopic mapping of a single metaphase chromosome. <i>Nucleic Acids Research</i> , 2019 , 47, e108	20.1	11
31	Squalamine and Its Derivatives Modulate the Aggregation of Amyloid-Land Esynuclein and Suppress the Toxicity of Their Oligomers. <i>Frontiers in Neuroscience</i> , 2021 , 15, 680026	5.1	11
30	Infrared nanospectroscopy reveals the molecular interaction fingerprint of an aggregation inhibitor with single AB2 oligomers. <i>Nature Communications</i> , 2021 , 12, 688	17.4	11
29	Controlled self-assembly of plant proteins into high-performance multifunctional nanostructured films. <i>Nature Communications</i> , 2021 , 12, 3529	17.4	10
28	Sequence-Optimized Peptide Nanofibers as Growth Stimulators for Regeneration of Peripheral Neurons. <i>Advanced Functional Materials</i> , 2019 , 29, 1809112	15.6	9
27	Rationally Designed Antibodies as Research Tools to Study the Structure-Toxicity Relationship of Amyloid-IDligomers. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	7
26	Environmental Control of Amyloid Polymorphism by Modulation of Hydrodynamic Stress. <i>ACS Nano</i> , 2021 , 15, 944-953	16.7	7
25	Effects of sedimentation, microgravity, hydrodynamic mixing and air-water interface on Esynuclein amyloid formation. <i>Chemical Science</i> , 2020 , 11, 3687-3693	9.4	7
24	Evolution of Conformation, Nanomechanics, and Infrared Nanospectroscopy of Single Amyloid Fibrils Converting into Microcrystals. <i>Advanced Science</i> , 2021 , 8, 2002182	13.6	6
23	A dopamine metabolite stabilizes neurotoxic amyloid-lībligomers. <i>Communications Biology</i> , 2021 , 4, 19	6.7	6

(2021-2019)

22	Characterizing Individual Protein Aggregates by Infrared Nanospectroscopy and Atomic Force Microscopy. <i>Journal of Visualized Experiments</i> , 2019 ,	1.6	5
21	One-Pot Semisynthesis of Exon 1 of the Huntingtin Protein: New Tools for Elucidating the Role of Posttranslational Modifications in the Pathogenesis of Huntington Disease. <i>Angewandte Chemie</i> , 2014 , 126, 1959-1964	3.6	5
20	Influence of the electro-optical properties of an Esi:H single layer on the performances of a pin solar cell. <i>Thin Solid Films</i> , 2012 , 520, 4036-4040	2.2	5
19	Small molecule sequestration of amyloid-las a drug discovery strategy for Alzheimer disease		4
18	A rationally designed bicyclic peptide remodels AB2 aggregation in vitro and reduces its toxicity in a worm model of Alzheimer's disease. <i>Scientific Reports</i> , 2020 , 10, 15280	4.9	4
17	Supramolecular Peptide Nanofibrils with Optimized Sequences and Molecular Structures for Efficient Retroviral Transduction. <i>Advanced Functional Materials</i> , 2021 , 31, 2009382	15.6	4
16	Alpha Synuclein only Forms Fibrils In Vitro when Larger than its Critical Size of 70 Monomers. <i>ChemBioChem</i> , 2021 , 22, 2867-2871	3.8	4
15	Frontispiece: Mutant Exon1 Huntingtin Aggregation is Regulated by T3 Phosphorylation-Induced Structural Changes and Crosstalk between T3 Phosphorylation and Acetylation at K6. <i>Angewandte Chemie - International Edition</i> , 2017 , 56,	16.4	3
14	Analysis of B -crystallin polydispersity in solution through native microfluidic electrophoresis. <i>Analyst, The</i> , 2019 , 144, 4413-4424	5	3
13	Thermoresponsive, Pyrrolidone-Based Antifouling Polymer Brushes. Advanced Materials Interfaces,2101	741 <i>6</i> 7	3
12	The Hsc70 Disaggregation Machinery Removes Monomer Units Directly from Esynuclein Fibril Ends		3
11	The Nt17 domain and its helical conformation regulate the aggregation, cellular properties and neurotoxicity of mutant huntingtin exon 1		3
10	Imaging protein aggregates in the serum and cerebrospinal fluid in Parkinson's disease. <i>Brain</i> , 2021 ,	11.2	3
9	Mutant Exon1 Huntingtin Aggregation is Regulated by T3 Phosphorylation-Induced Structural Changes and Crosstalk between T3 Phosphorylation and Acetylation at K6. <i>Angewandte Chemie</i> , 2017 , 129, 5286-5291	3.6	2
8	The Hsc70 disaggregation machinery removes monomer units directly from Esynuclein fibril ends. <i>Nature Communications</i> , 2021 , 12, 5999	17.4	2
7	In situ Sub-Cellular Identification of Functional Amyloids in Bacteria and Archaea by Infrared Nanospectroscopy <i>Small Methods</i> , 2021 , 5, e2001002	12.8	2
6	Soluble amyloid beta-containing aggregates are present throughout the brain at early stages of Alzheimer's disease. <i>Brain Communications</i> , 2021 , 3, fcab147	4.5	2
5	The Nt17 Domain and its Helical Conformation Regulate the Aggregation, Cellular Properties and Neurotoxicity of Mutant Huntingtin Exon 1. <i>Journal of Molecular Biology</i> , 2021 , 433, 167222	6.5	2

4	Frontispiz: Mutant Exon1 Huntingtin Aggregation is Regulated by T3 Phosphorylation-Induced Structural Changes and Crosstalk between T3 Phosphorylation and Acetylation at K6. <i>Angewandte Chemie</i> , 2017 , 129,	3.6	1
3	N-terminal phosphorylation of Huntingtin: A molecular switch for regulating Htt aggregation, helical conformation, internalization and nuclear targeting		1
2	Berichtigung: One-Pot Semisynthesis of Exon 1 of the Huntingtin Protein: New Tools for Elucidating the Role of Posttranslational Modifications in the Pathogenesis of Huntington Disease. <i>Angewandte Chemie</i> , 2014 , 126, 7517-7517	3.6	
1	Unraveling the Physicochemical Determinants of Protein Liquid-liquid Phase Separation by Nanoscale Infrared Vibrational Spectroscopy. <i>Bio-protocol</i> , 2021 , 11, e4122	0.9	