

# Marcel Wiermer

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2985057/publications.pdf>

Version: 2024-02-01

34  
papers

3,671  
citations

279701

23  
h-index

395590

33  
g-index

38  
all docs

38  
docs citations

38  
times ranked

4409  
citing authors

#	ARTICLE	IF	CITATIONS
1	SEED LIPID DROPLET PROTEIN1, SEED LIPID DROPLET PROTEIN2, and LIPID DROPLET PLASMA MEMBRANE ADAPTOR mediate lipid dropletâ€ plasma membrane tethering. <i>Plant Cell</i> , 2022, 34, 2424-2448.	3.1	12
2	Cell wall-localized BETA-XYLOSIDASE4 contributes to immunity of Arabidopsis against <i>Botrytis cinerea</i> . <i>Plant Physiology</i> , 2022, 189, 1794-1813.	2.3	14
3	EXTRA LARGE G-PROTEIN2 mediates cell death and hyperimmunity in the <i>chitin elicitor receptor kinase 1-4</i> mutant. <i>Plant Physiology</i> , 2022, 189, 2413-2431.	2.3	5
4	<sc>NLR</sc> we there yet? Nucleocytoplasmic coordination of <sc>NLR</sc>-mediated immunity. <i>New Phytologist</i> , 2022, 236, 24-42.	3.5	12
5	SCF<sup>SNIPER7</sup> controls protein turnover of unfoldase CDC48A to promote plant immunity. <i>New Phytologist</i> , 2021, 229, 2795-2811.	3.5	13
6	Functional requirement of the <i>Arabidopsis</i> importinâ€ nuclear transport receptor family in autoimmunity mediated by the NLR protein SNC1. <i>Plant Journal</i> , 2021, 105, 994-1009.	2.8	20
7	Nucleocytoplasmic Communication in Healthy and Diseased Plant Tissues. <i>Frontiers in Plant Science</i> , 2021, 12, 719453.	1.7	9
8	Isochorismate-derived biosynthesis of the plant stress hormone salicylic acid. <i>Science</i> , 2019, 365, 498-502.	6.0	273
9	STOREKEEPER RELATED1/G-Element Binding Protein (STKR1) Interacts with Protein Kinase SnRK1. <i>Plant Physiology</i> , 2018, 176, 1773-1792.	2.3	31
10	MOS6 and TN13 in plant immunity. <i>Plant Signaling and Behavior</i> , 2018, 13, e1454816.	1.2	2
11	Nucleoporin NUP88/MOS7 is required for manifestation of phenotypes associated with the Arabidopsis CHITIN ELICITOR RECEPTOR KINASE1 mutant <i>cerk1</i> . <i>Plant Signaling and Behavior</i> , 2017, 12, e1313378.	1.2	1
12	The putative kinase substrate MUSE7 negatively impacts the accumulation of <sc>NLR</sc> proteins. <i>Plant Journal</i> , 2017, 89, 1174-1183.	2.8	4
13	The truncated NLR protein TIRâ€NBS13 is a MOS6/IMPORTINâ€ interaction partner required for plant immunity. <i>Plant Journal</i> , 2017, 92, 808-821.	2.8	43
14	E3 ligase SAUL1 serves as a positive regulator of PAMP-triggered immunity and its homeostasis is monitored by immune receptor SOC3. <i>New Phytologist</i> , 2017, 215, 1516-1532.	3.5	69
15	Nucleoporin-regulated MAP kinase signaling in immunity to a necrotrophic fungal pathogen. <i>Plant Physiology</i> , 2016, 172, pp.00832.2016.	2.3	31
16	Probing formation of cargo/importinâ€ transport complexes in plant cells using a pathogen effector. <i>Plant Journal</i> , 2015, 81, 40-52.	2.8	48
17	Two Activities of Long-Chain Acyl-Coenzyme A Synthetase Are Involved in Lipid Trafficking between the Endoplasmic Reticulum and the Plastid in Arabidopsis. <i>Plant Physiology</i> , 2015, 167, 351-366.	2.3	109
18	An E4 Ligase Facilitates Polyubiquitination of Plant Immune Receptor Resistance Proteins in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2014, 26, 485-496.	3.1	57

#	ARTICLE	IF	CITATIONS
19	Mitochondrial AtPAM16 is required for plant survival and the negative regulation of plant immunity. <i>Nature Communications</i> , 2013, 4, 2558.	5.8	64
20	Analyses of <i>wrky18 wrky40</i> Plants Reveal Critical Roles of SA/EDS1 Signaling and Indole-Glucosinolate Biosynthesis for <i>Golovinomyces orontii</i> Resistance and a Loss-of Resistance Towards <i>Pseudomonas syringae</i> pv. <i>tomato</i> AvrRPS4. <i>Molecular Plant-Microbe Interactions</i> , 2013, 26, 758-767.	1.4	91
21	Hop-on hop-off: importin- $\alpha$ -guided tours to the nucleus in innate immune signaling. <i>Frontiers in Plant Science</i> , 2013, 4, 149.	1.7	58
22	The Salmonella Type III Effector SspH2 Specifically Exploits the NLR Co-chaperone Activity of SGT1 to Subvert Immunity. <i>PLoS Pathogens</i> , 2013, 9, e1003518.	2.1	80
23	Nucleoporins Nup160 and Seh1 are required for disease resistance in Arabidopsis. <i>Plant Signaling and Behavior</i> , 2012, 7, 1212-1214.	1.2	29
24	The cyclin-FL homolog MOS12 and the MOS4-associated complex are required for the proper splicing of plant <i>resistance</i> genes. <i>Plant Journal</i> , 2012, 70, 916-928.	2.8	86
25	Putative members of the Arabidopsis Nup107-160 nuclear pore subcomplex contribute to pathogen defense. <i>Plant Journal</i> , 2012, 70, 796-808.	2.8	74
26	Balanced Nuclear and Cytoplasmic Activities of EDS1 Are Required for a Complete Plant Innate Immune Response. <i>PLoS Pathogens</i> , 2010, 6, e1000970.	2.1	202
27	Nucleoporin MOS7/Nup88 contributes to plant immunity and nuclear accumulation of defense regulators. <i>Nucleus</i> , 2010, 1, 332-336.	0.6	30
28	Arabidopsis resistance protein SNC1 activates immune responses through association with a transcriptional corepressor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 13960-13965.	3.3	205
29	Nuclear Pore Complex Component MOS7/Nup88 Is Required for Innate Immunity and Nuclear Accumulation of Defense Regulators in Arabidopsis. <i>Plant Cell</i> , 2009, 21, 2503-2516.	3.1	233
30	Should I stay or should I go? Nucleocytoplasmic trafficking in plant innate immunity. <i>Cellular Microbiology</i> , 2007, 9, 1880-1890.	1.1	56
31	Plant immunity: the EDS1 regulatory node. <i>Current Opinion in Plant Biology</i> , 2005, 8, 383-389.	3.5	542
32	Arabidopsis SENESCENCE-ASSOCIATED GENE101 Stabilizes and Signals within an ENHANCED DISEASE SUSCEPTIBILITY1 Complex in Plant Innate Immunity. <i>Plant Cell</i> , 2005, 17, 2601-2613.	3.1	413
33	Pre- and Postinvasion Defenses Both Contribute to Nonhost Resistance in Arabidopsis. <i>Science</i> , 2005, 310, 1180-1183.	6.0	753
34	Marshalling the Troops: Intracellular Dynamics in Plant Pathogen Defense. , 0 , 177-219.		1