

# Miguel Silva-Ramos

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2985086/publications.pdf>

Version: 2024-02-01

24  
papers

426  
citations

840776

11  
h-index

839539

18  
g-index

24  
all docs

24  
docs citations

24  
times ranked

687  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prostate Cancer Prevention Trial and European Randomized Study of Screening for Prostate Cancer Risk Calculators: A Performance Comparison in a Contemporary Screened Cohort. <i>European Urology</i> , 2010, 58, 551-558.	1.9	82
2	ATP released via pannexin-1 hemichannels mediates bladder overactivity triggered by urothelial P2Y6 receptors. <i>Biochemical Pharmacology</i> , 2014, 87, 371-379.	4.4	62
3	Urinary ATP May Be a Dynamic Biomarker of Detrusor Overactivity in Women with Overactive Bladder Syndrome. <i>PLoS ONE</i> , 2013, 8, e64696.	2.5	57
4	Inhibition of cholinergic neurotransmission by $\beta^2$ -adrenoceptors depends on adenosine release and A <sub>1</sub> -receptor activation in human and rat urinary bladders. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 313, F388-F403.	2.7	35
5	Activation of P2Y <sub>6</sub> Receptors Facilitates Nonneuronal Adenosine Triphosphate and Acetylcholine Release from Urothelium with the Lamina Propria of Men with Bladder Outlet Obstruction. <i>Journal of Urology</i> , 2015, 194, 1146-1154.	0.4	28
6	Activation of P2Y <sub>6</sub> receptors increases the voiding frequency in anaesthetized rats by releasing ATP from the bladder urothelium. <i>British Journal of Pharmacology</i> , 2014, 171, 3404-3419.	5.4	24
7	The CAG repeat within the androgen receptor gene and its relationship to cryptorchidism. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2006, 32, 330-335.	1.5	23
8	Impairment of ATP hydrolysis decreases adenosine A <sub>1</sub> receptor tonus favoring cholinergic nerve hyperactivity in the obstructed human urinary bladder. <i>Purinergic Signalling</i> , 2015, 11, 595-606.	2.2	19
9	Increased Urinary Adenosine Triphosphate in Patients With Bladder Outlet Obstruction Due to Benign Prostate Hyperplasia. <i>Prostate</i> , 2016, 76, 1353-1363.	2.3	18
10	CT-based renal volume and graft function after living-donor kidney transplantation: Is there a volume threshold to avoid?. <i>International Urology and Nephrology</i> , 2015, 47, 851-859.	1.4	16
11	The quality of reporting of randomized controlled trials in pelvic organ prolapse. <i>International Urogynecology Journal</i> , 2011, 22, 1117-1125.	1.4	11
12	Comparative study of gene expression in patients with varicocele by microarray technology. <i>Andrologia</i> , 2012, 44, 260-265.	2.1	11
13	Living Versus Cadaveric-donor Renal Transplant Recipients: A Comparison on Sexual Function. <i>Transplantation Proceedings</i> , 2013, 45, 1066-1069.	0.6	11
14	$\beta^2$ Adrenoceptor-induced cholinergic inhibition in human and rat urinary bladders involves the exchange protein directly activated by cyclic AMP 1 favoring adenosine release. <i>British Journal of Pharmacology</i> , 2020, 177, 1589-1608.	5.4	11
15	Lower donated kidney volume is associated with increased risk of lower graft function and acute rejection at 1 year after living donor kidney—a retrospective study. <i>Transplant International</i> , 2020, 33, 1711-1722.	1.6	7
16	Does 3D Ultrasound Enhance the Diagnosis of Bladder Tumours in Patients with Haematuria?. <i>ISRN Urology</i> , 2012, 2012, 1-5.	1.5	5
17	Remaining kidney volume indexed to weight as a strong predictor of estimated glomerular filtration rate at 1 year and mid-term renal function after living-donor nephrectomy—a retrospective observational study. <i>Transplant International</i> , 2020, 33, 1262-1273.	1.6	4
18	Marginal zone lymphoma of both spleen and kidney displaying transformation into large B-cell lymphoma. <i>International Urology and Nephrology</i> , 2006, 38, 431-437.	1.4	2

#	ARTICLE	IF	CITATIONS
19	249: Decreased Hydrolysis of ATP Acting Via P2X Receptors Favour Cholinergic Nerve Hyperactivity in Human Obstructed Bladders. <i>Journal of Urology</i> , 2007, 177, 83-83.	0.4	0
20	Hyperactivity of cholinergic nerves associated with decreased hydrolysis of ATP in obstructed human bladders. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2007, 135, 22.	2.8	0
21	ARE ONLINE-AVAILABLE PROSTATE CANCER RISK CALCULATORS AN ADDED VALUE?. <i>Journal of Urology</i> , 2009, 181, 750-750.	0.4	0
22	Dual role of P2Y6 receptors in the human urinary bladder: On the role of ATP released from the urothelium. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2013, 177, 303-304.	2.8	0
23	Overactive bladder syndrome: Evidence that urinary ATP is a dynamic biomarker of detrusor overactivity. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2013, 177, 21-22.	2.8	0
24	Transvaginal-Assisted Laparoscopic Nephrectomy: A Step Toward Natural Orifice Transluminal Endoscopic Surgery. <i>Videourology (New Rochelle, N Y)</i> , 2011, 25, .	0.1	0