

Miguel Silva-Ramos

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2985086/publications.pdf>

Version: 2024-02-01

24
papers

426
citations

840776

11
h-index

839539

18
g-index

24
all docs

24
docs citations

24
times ranked

687
citing authors

#	ARTICLE	IF	CITATIONS
1	β_3 Adrenoceptor-induced cholinergic inhibition in human and rat urinary bladders involves the exchange protein directly activated by cyclic AMP 1 favoring adenosine release. British Journal of Pharmacology, 2020, 177, 1589-1608.	5.4	11
2	Lower donated kidney volume is associated with increased risk of lower graft function and acute rejection at 1 year after living donor kidney—a retrospective study. Transplant International, 2020, 33, 1711-1722.	1.6	7
3	Remaining kidney volume indexed to weight as a strong predictor of estimated glomerular filtration rate at 1 year and mid-term renal function after living-donor nephrectomy—a retrospective observational study. Transplant International, 2020, 33, 1262-1273.	1.6	4
4	Inhibition of cholinergic neurotransmission by β_3 -adrenoceptors depends on adenosine release and A_1 -receptor activation in human and rat urinary bladders. American Journal of Physiology - Renal Physiology, 2017, 313, F388-F403.	2.7	35
5	Increased Urinary Adenosine Triphosphate in Patients With Bladder Outlet Obstruction Due to Benign Prostate Hyperplasia. Prostate, 2016, 76, 1353-1363.	2.3	18
6	Impairment of ATP hydrolysis decreases adenosine A_1 receptor tonus favoring cholinergic nerve hyperactivity in the obstructed human urinary bladder. Purinergic Signalling, 2015, 11, 595-606.	2.2	19
7	CT-based renal volume and graft function after living-donor kidney transplantation: Is there a volume threshold to avoid?. International Urology and Nephrology, 2015, 47, 851-859.	1.4	16
8	Activation of P_2Y_6 Receptors Facilitates Nonneuronal Adenosine Triphosphate and Acetylcholine Release from Urothelium with the Lamina Propria of Men with Bladder Outlet Obstruction. Journal of Urology, 2015, 194, 1146-1154.	0.4	28
9	Activation of P_2Y_6 receptors increases the voiding frequency in anaesthetized rats by releasing ATP from the bladder urothelium. British Journal of Pharmacology, 2014, 171, 3404-3419.	5.4	24
10	ATP released via pannexin-1 hemichannels mediates bladder overactivity triggered by urothelial P_2Y_6 receptors. Biochemical Pharmacology, 2014, 87, 371-379.	4.4	62
11	Living Versus Cadaveric-donor Renal Transplant Recipients: A Comparison on Sexual Function. Transplantation Proceedings, 2013, 45, 1066-1069.	0.6	11
12	Dual role of P_2Y_6 receptors in the human urinary bladder: On the role of ATP released from the urothelium. Autonomic Neuroscience: Basic and Clinical, 2013, 177, 303-304.	2.8	0
13	Overactive bladder syndrome: Evidence that urinary ATP is a dynamic biomarker of detrusor overactivity. Autonomic Neuroscience: Basic and Clinical, 2013, 177, 21-22.	2.8	0
14	Urinary ATP May Be a Dynamic Biomarker of Detrusor Overactivity in Women with Overactive Bladder Syndrome. PLoS ONE, 2013, 8, e64696.	2.5	57
15	Does 3D Ultrasound Enhance the Diagnosis of Bladder Tumours in Patients with Haematuria?. ISRN Urology, 2012, 2012, 1-5.	1.5	5
16	Comparative study of gene expression in patients with varicocele by microarray technology. Andrologia, 2012, 44, 260-265.	2.1	11
17	The quality of reporting of randomized controlled trials in pelvic organ prolapse. International Urogynecology Journal, 2011, 22, 1117-1125.	1.4	11
18	Transvaginal-Assisted Laparoscopic Nephrectomy: A Step Toward Natural Orifice Transluminal Endoscopic Surgery. Videourology (New Rochelle, N Y), 2011, 25, .	0.1	0

#	ARTICLE	IF	CITATIONS
19	Prostate Cancer Prevention Trial and European Randomized Study of Screening for Prostate Cancer Risk Calculators: A Performance Comparison in a Contemporary Screened Cohort. <i>European Urology</i> , 2010, 58, 551-558.	1.9	82
20	ARE ONLINE-AVAILABLE PROSTATE CANCER RISK CALCULATORS AN ADDED VALUE?. <i>Journal of Urology</i> , 2009, 181, 750-750.	0.4	0
21	249: Decreased Hydrolysis of ATP Acting Via P2X Receptors Favour Cholinergic Nerve Hyperactivity in Human Obstructed Bladders. <i>Journal of Urology</i> , 2007, 177, 83-83.	0.4	0
22	Hyperactivity of cholinergic nerves associated with decreased hydrolysis of ATP in obstructed human bladders. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2007, 135, 22.	2.8	0
23	Marginal zone lymphoma of both spleen and kidney displaying transformation into large B-cell lymphoma. <i>International Urology and Nephrology</i> , 2006, 38, 431-437.	1.4	2
24	The CAG repeat within the androgen receptor gene and its relationship to cryptorchidism. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2006, 32, 330-335.	1.5	23