

Gerald Zernig

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2992256/gerald-zernig-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85
papers

2,301
citations

31
h-index

44
g-index

92
ext. papers

2,569
ext. citations

4.1
avg. IF

4.48
L-index

#	Paper	IF	Citations
85	Venlafaxine and O-desmethylvenlafaxine serum levels are positively associated with antidepressant response in elder depressed out-patients. <i>World Journal of Biological Psychiatry</i> , 2021 , 1-8	3.8	2
84	Extremely High-Dosage Zolpidem Poisoning With Favorable Outcome. <i>Journal of Clinical Psychopharmacology</i> , 2021 , 41, 222-223	1.7	1
83	Therapeutic Drug Monitoring of Long-Acting Injectable Antipsychotic Drugs. <i>Therapeutic Drug Monitoring</i> , 2021 , 43, 79-102	3.2	9
82	Switch to 3-Month Long-Acting Injectable Paliperidone May Decrease Plasma Levels: A Case Series. <i>Journal of Clinical Psychopharmacology</i> , 2021 , 41, 694-696	1.7	0
81	Antidepressant efficacy is correlated with plasma levels: mega-analysis and further evidence.. <i>International Clinical Psychopharmacology</i> , 2021 , 37,	2.2	2
80	Blood Levels to Optimize Antipsychotic Treatment in Clinical Practice: A Joint Consensus Statement of the American Society of Clinical Psychopharmacology and the Therapeutic Drug Monitoring Task Force of the Arbeitsgemeinschaft für Neuropsychopharmakologie und Pharmakopsychiatrie. <i>Journal of Clinical Psychiatry</i> , 2020 , 81,	4.6	39
79	Guía de consenso de expertos para la monitorización terapéutica de drogas en neuropsicofarmacología. <i>Psiquiatría Biológica</i> , 2020 , 27, 83-95	0.2	0
78	Non-adherence to psychotropic medication assessed by plasma level in newly admitted psychiatric patients: Prevalence before acute admission. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 175-178	6.2	5
77	TDM in psychiatry and neurology: A comprehensive summary of the consensus guidelines for therapeutic drug monitoring in neuropsychopharmacology, update 2017; a tool for clinicians. <i>World Journal of Biological Psychiatry</i> , 2018 , 19, 162-174	3.8	64
76	Making the Case for Power Abuse Disorder as a Nosologic Entity. <i>Pharmacology</i> , 2017 , 100, 50-63	2.3	2
75	Preventive Strength of Dyadic Social Interaction against Reacquisition/Reexpression of Cocaine Conditioned Place Preference. <i>Frontiers in Behavioral Neuroscience</i> , 2017 , 11, 225	3.5	6
74	Changes in psychopathology in schizophrenia patients starting treatment with new-generation antipsychotics: therapeutic drug monitoring in a naturalistic treatment setting. <i>European Neuropsychopharmacology</i> , 2016 , 26, 717-28	1.2	6
73	Dyadic social interaction of C57BL/6 mice versus interaction with a toy mouse: conditioned place preference/aversion, substrain differences, and no development of a hierarchy. <i>Behavioural Pharmacology</i> , 2016 , 27, 279-88	2.4	8
72	Dyadic social interaction inhibits cocaine-conditioned place preference and the associated activation of the accumbens corridor. <i>Behavioural Pharmacology</i> , 2015 , 26, 580-94	2.4	34
71	Social interaction and cocaine conditioning in mice increase spontaneous spike frequency in the nucleus accumbens or septal nuclei as revealed by multielectrode array recordings. <i>Pharmacology</i> , 2015 , 95, 42-9	2.3	12
70	Reacquisition of cocaine conditioned place preference and its inhibition by previous social interaction preferentially affect D1-medium spiny neurons in the accumbens corridor. <i>Frontiers in Behavioral Neuroscience</i> , 2014 , 8, 317	3.5	16
69	Differences in social interaction- vs. cocaine reward in mouse vs. rat. <i>Frontiers in Behavioral Neuroscience</i> , 2014 , 8, 363	3.5	33

68	Increased conditioned place preference for cocaine in high anxiety related behavior (HAB) mice is associated with an increased activation in the accumbens corridor. <i>Frontiers in Behavioral Neuroscience</i> , 2014 , 8, 441	3.5	13
67	Effects of antipsychotic treatment on cognition in healthy subjects. <i>Journal of Psychopharmacology</i> , 2013 , 27, 374-85	4.6	23
66	Dyadic social interaction as an alternative reward to cocaine. <i>Frontiers in Psychiatry</i> , 2013 , 4, 100	5	44
65	Preventive role of social interaction for cocaine conditioned place preference: correlation with FosB/DeltaFosB and pCREB expression in rat mesocorticolimbic areas. <i>Frontiers in Behavioral Neuroscience</i> , 2012 , 6, 8	3.5	25
64	Brain regions associated with the acquisition of conditioned place preference for cocaine vs. social interaction. <i>Frontiers in Behavioral Neuroscience</i> , 2012 , 6, 63	3.5	34
63	Acetylcholine, drug reward and substance use disorder treatment: intra- and interindividual striatal and accumbal neuron ensemble heterogeneity may explain apparent discrepant findings. <i>Pharmacology</i> , 2012 , 90, 264-73	2.3	7
62	Activation of PKCzeta and PKMzeta in the nucleus accumbens core is necessary for the retrieval, consolidation and reconsolidation of drug memory. <i>PLoS ONE</i> , 2012 , 7, e30502	3.7	31
61	Conditioned place preference for social interaction in rats: contribution of sensory components. <i>Frontiers in Behavioral Neuroscience</i> , 2011 , 5, 80	3.5	47
60	Differential effects of accumbens core vs. shell lesions in a rat concurrent conditioned place preference paradigm for cocaine vs. social interaction. <i>PLoS ONE</i> , 2011 , 6, e26761	3.7	35
59	Reversal of cocaine-conditioned place preference and mesocorticolimbic Zif268 expression by social interaction in rats. <i>Addiction Biology</i> , 2011 , 16, 273-84	4.6	83
58	"There is no dose-response relationship in psychopharmacotherapy" vs "pharmacotherapy in psychiatry is based on ligand-receptor interaction": a unifying hypothesis and the need for plasma concentration based clinical trials. <i>Psychopharmacology</i> , 2011 , 217, 297-300	4.7	10
57	Effects of antipsychotic treatment on psychopathology and motor symptoms. A placebo-controlled study in healthy volunteers. <i>Psychopharmacology</i> , 2011 , 218, 733-48	4.7	11
56	Sigma1 receptor antagonist BD1047 enhances reversal of conditioned place preference from cocaine to social interaction. <i>Pharmacology</i> , 2011 , 87, 45-8	2.3	25
55	Comparable sensitivities of urine cotinine and breath carbon monoxide at follow-up time points of three months or more in a smoking cessation trial. <i>Pharmacology</i> , 2010 , 85, 234-40	2.3	13
54	Serum concentrations of paliperidone versus risperidone and clinical effects. <i>European Journal of Clinical Pharmacology</i> , 2010 , 66, 797-803	2.8	33
53	Withania somnifera prevents morphine withdrawal-induced decrease in spine density in nucleus accumbens shell of rats: a confocal laser scanning microscopy study. <i>Neurotoxicity Research</i> , 2009 , 16, 343-55	4.3	32
52	Haloperidol and risperidone have specific effects on altered pain sensitivity in the ketamine model of schizophrenia. <i>Psychopharmacology</i> , 2009 , 202, 579-87	4.7	15
51	The impact of early environmental rearing condition on the discriminative stimulus effects and Fos expression induced by cocaine in adult male and female rats. <i>Psychopharmacology</i> , 2009 , 203, 383-97	4.7	13

50	A randomized trial of short psychotherapy versus sustained-release bupropion for smoking cessation. <i>Addiction</i> , 2008 , 103, 2024-31	4.6	27
49	[Commentary] REPLICATION AND FURTHER SCIENTIFIC INVESTIGATIONS WILL TELL THE TRUTH. <i>Addiction</i> , 2008 , 103, 2033-2034	4.6	2
48	Nucleus accumbens core acetylcholine is preferentially activated during acquisition of drug- vs food-reinforced behavior. <i>Neuropsychopharmacology</i> , 2008 , 33, 3213-20	8.7	24
47	Pharmacological validation of a chronic social stress model of depression in rats: effects of reboxetine, haloperidol and diazepam. <i>Behavioural Pharmacology</i> , 2008 , 19, 183-96	2.4	84
46	Phase 1A safety assessment of intravenous amitriptyline. <i>Journal of Pain</i> , 2007 , 8, 549-55	5.2	15
45	Differential effects of intravenous R,S-(+/-)-3,4-methylenedioxymethamphetamine (MDMA, Ecstasy) and its S(+)- and R(-)-enantiomers on dopamine transmission and extracellular signal regulated kinase phosphorylation (pERK) in the rat nucleus accumbens shell and core. <i>Journal of Neurochemistry</i> , 2007 , 102, 181-92	6	47
44	Explaining the escalation of drug use in substance dependence: models and appropriate animal laboratory tests. <i>Pharmacology</i> , 2007 , 80, 65-119	2.3	112
43	Age and gender effects on olanzapine and risperidone plasma concentrations in children and adolescents. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2007 , 17, 665-74	2.9	36
42	Therapeutic drug monitoring-based clozapine dosing recommendations. <i>Therapeutic Drug Monitoring</i> , 2007 , 29, 130-1	3.2	4
41	Mirtazapine and breastfeeding: maternal and infant plasma levels. <i>American Journal of Psychiatry</i> , 2007 , 164, 348-9	11.9	17
40	Pharmacokinetics and elimination of quetiapine, venlafaxine, and trazodone during pregnancy and postpartum. <i>Journal of Clinical Psychopharmacology</i> , 2007 , 27, 720-2	1.7	22
39	Quetiapine cross reactivity with urine methadone immunoassays. <i>American Journal of Psychiatry</i> , 2007 , 164, 172	11.9	23
38	Methadone doses upon multiple readmissions to inpatient detoxification: Clinical evidence for very moderate opioid tolerance. <i>Pharmacology</i> , 2006 , 78, 38-43	2.3	3
37	Activation of muscarinic and nicotinic acetylcholine receptors in the nucleus accumbens core is necessary for the acquisition of drug reinforcement. <i>Journal of Neuroscience</i> , 2006 , 26, 6004-10	6.6	63
36	Influence of age, gender, body weight and valproate comedication on quetiapine plasma concentrations. <i>International Clinical Psychopharmacology</i> , 2006 , 21, 81-5	2.2	62
35	Peri-response pharmacokinetics of remifentanyl during a self-administration session indicates that neither blood nor brain levels are titrated. <i>Annals of the New York Academy of Sciences</i> , 2006 , 1074, 497-504	6.5	6
34	St. John's wort (<i>Hypericum perforatum</i>) and breastfeeding: plasma and breast milk concentrations of hyperforin for 5 mothers and 2 infants. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 305-9	4.6	35
33	Effect of 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") on dopamine transmission in the nucleus accumbens shell and core. <i>Brain Research</i> , 2005 , 1055, 143-8	3.7	39

32	Simultaneous intra-accumbens remifentanyl and dopamine kinetics suggest that neither determines within-session operant responding. <i>Psychopharmacology</i> , 2005 , 183, 201-9	4.7	18
31	Influence of age and gender on risperidone plasma concentrations. <i>Journal of Psychopharmacology</i> , 2005 , 19, 395-401	4.6	81
30	Therapeutic monitoring of aripiprazole by HPLC with column-switching and spectrophotometric detection. <i>Clinical Chemistry</i> , 2005 , 51, 1718-21	5.5	64
29	Subjective effects of slow-release bupropion versus caffeine as determined in a quasi-naturalistic setting. <i>Pharmacology</i> , 2004 , 70, 206-15	2.3	25
28	Drug history overrides opioid reinforcement in a rat runway procedure. <i>Pharmacology</i> , 2004 , 72, 225-30	2.3	5
27	Do vertical shifts in dose-response rate-relationships in operant conditioning procedures indicate "sensitization" to "drug wanting"? <i>Psychopharmacology</i> , 2004 , 171, 349-51; author reply 352-63	4.7	41
26	What the clinician still has to be reminded of. <i>Therapeutic Drug Monitoring</i> , 2004 , 26, 582	3.2	11
25	Methylenedioxymethamphetamine (MDMA, Ecstasy) serves as a robust positive reinforcer in a rat runway procedure. <i>Pharmacology</i> , 2003 , 69, 180-2	2.3	14
24	Intravenous drug injection habits: drug users' self-reports versus researchers' perception. <i>Pharmacology</i> , 2003 , 68, 49-56	2.3	37
23	Opioids, cocaine, and food change runtime distribution in a rat runway procedure. <i>Psychopharmacology</i> , 2003 , 169, 52-9	4.7	20
22	Neutral endopeptidase knockout induces hyperalgesia in a model of visceral pain, an effect related to bradykinin and nitric oxide. <i>Journal of Molecular Neuroscience</i> , 2002 , 18, 129-34	3.3	25
21	Reinforcing effect of subcutaneous morphine in a modified Ettenberg runway. <i>Journal of Molecular Neuroscience</i> , 2002 , 18, 135-42	3.3	6
20	Modeling addiction: trusted experimental approaches are tried in new applications. <i>Trends in Pharmacological Sciences</i> , 2002 , 23, 399-400	13.2	2
19	Lack of a distinctive behavioural effect of chromogranin-derived peptides in rodents. <i>Regulatory Peptides</i> , 2002 , 103, 85-91		1
18	Reinforcing effects of MDMA ("ecstasy") in drug-naive and cocaine-trained rats. <i>Pharmacology</i> , 2001 , 62, 138-44	2.3	76
17	Lack of reinforcing effect of the benzodiazepine and tricyclic antidepressant combination of diazepam and dothiepin. <i>Pharmacology</i> , 2001 , 62, 124-8	2.3	5
16	Alterations within the endogenous opioid system in mice with targeted deletion of the neutral endopeptidase (nekephalinase) gene. <i>Regulatory Peptides</i> , 2000 , 96, 53-8		23
15	Signal transduction efficacy of the highly potent mu opioid agonist 14-methoxymetopon. <i>Life Sciences</i> , 2000 , 66, 1871-7	6.8	18

14	Opioid receptor modulation of feeding-evoked dopamine release in the rat nucleus accumbens. <i>Brain Research</i> , 1998 , 785, 24-30	3.7	30
13	Pharmacotherapy of alcohol dependence. <i>Trends in Pharmacological Sciences</i> , 1997 , 18, 229-231	13.2	17
12	Calculation of agonist efficacy, apparent affinity, and receptor population changes after administration of insurmountable antagonists: comparison of different analytical approaches. <i>Journal of Pharmacological and Toxicological Methods</i> , 1996 , 35, 223-37	1.7	27
11	Receptor reserve and affinity of mu opioid agonists in mouse antinociception: correlation with receptor binding. <i>Life Sciences</i> , 1995 , 57, 2113-25	6.8	36
10	Opioid agonist effects on mouse writhing after irreversible mu receptor blockade with clocinnamox.. <i>Experimental and Clinical Psychopharmacology</i> , 1995 , 3, 323-329	3.2	6
9	Influence of milk on haloperidol pharmacokinetics.. <i>Experimental and Clinical Psychopharmacology</i> , 1994 , 2, 107-109	3.2	
8	Different behavioral profiles of the non-peptide substance P (NK1) antagonists CP-96,345 and RP 67580 in Swiss albino mice in the black-and-white box. <i>Neuroscience Letters</i> , 1993 , 151, 64-6	3.3	18
7	Different behavioral profiles of the non-peptide substance P (NK-1) antagonists CP-96,345 and RP 67580. <i>Regulatory Peptides</i> , 1993 , 46, 346-8		3
6	Stereoselective features of (R)- and (S)-atenolol: clinical pharmacological, pharmacokinetic, and radioligand binding studies. <i>Chirality</i> , 1993 , 5, 15-9	2.1	47
5	The substance P (NK1) receptor antagonist (+/-)-CP-96,345 causes sedation and motor impairment in Swiss albino mice in the black-and-white box behavioral paradigm. <i>Neuroscience Letters</i> , 1992 , 143, 169-72	3.3	22
4	Different stereoselective effects of (R)- and (S)-propafenone: clinical pharmacologic, electrophysiologic, and radioligand binding studies. <i>Clinical Pharmacology and Therapeutics</i> , 1990 , 47, 740-6	6.1	35
3	Widening potential for Ca ²⁺ antagonists: non-L-type Ca ²⁺ channel interaction. <i>Trends in Pharmacological Sciences</i> , 1990 , 11, 38-44	13.2	118
2	Stereoselective binding of niguldipine enantiomers to alpha 1A-adrenoceptors labeled with [3H]5-methyl-urapidil. <i>European Journal of Pharmacology</i> , 1989 , 172, 329-37		33
1	Human red-blood-cell Ca ²⁺ -antagonist binding sites. Evidence for an unusual receptor coupled to the nucleoside transporter. <i>FEBS Journal</i> , 1985 , 150, 67-77		64