

Stephen Eyre

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2994413/stephen-eyre-publications-by-year.pdf>

Version: 2024-04-18

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

133
papers

22,288
citations

49
h-index

144
g-index

144
ext. papers

24,960
ext. citations

10.7
avg, IF

7.36
L-index

#	Paper	IF	Citations
133	Uncovering genetic mechanisms of hypertension through multi-omic analysis of the kidney. <i>Nature Genetics</i> , 2021 , 53, 630-637	36.3	5
132	Persistent inflammatory and non-inflammatory mechanisms in refractory rheumatoid arthritis. <i>Nature Reviews Rheumatology</i> , 2021 , 17, 17-33	8.1	29
131	Functional genomics atlas of synovial fibroblasts defining rheumatoid arthritis heritability. <i>Genome Biology</i> , 2021 , 22, 247	18.3	6
130	Exploring the overlap between rheumatoid arthritis susceptibility loci and long non-coding RNA annotations. <i>PLoS ONE</i> , 2020 , 15, e0223939	3.7	1
129	Lymphocyte DNA methylation mediates genetic risk at shared immune-mediated disease loci. <i>Journal of Allergy and Clinical Immunology</i> , 2020 , 145, 1438-1451	11.5	10
128	Analysis of chromatin organization and gene expression in T cells identifies functional genes for rheumatoid arthritis. <i>Nature Communications</i> , 2020 , 11, 4402	17.4	12
127	Combined genetic analysis of juvenile idiopathic arthritis clinical subtypes identifies novel risk loci, target genes and key regulatory mechanisms. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	10
126	Exploring the overlap between rheumatoid arthritis susceptibility loci and long non-coding RNA annotations 2020 , 15, e0223939		
125	Exploring the overlap between rheumatoid arthritis susceptibility loci and long non-coding RNA annotations 2020 , 15, e0223939		
124	Exploring the overlap between rheumatoid arthritis susceptibility loci and long non-coding RNA annotations 2020 , 15, e0223939		
123	Exploring the overlap between rheumatoid arthritis susceptibility loci and long non-coding RNA annotations 2020 , 15, e0223939		
122	Genetics of rheumatoid arthritis: 2018 status. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 446-453	2.4	80
121	CD4+ and B Lymphocyte Expression Quantitative Traits at Rheumatoid Arthritis Risk Loci in Patients With Untreated Early Arthritis: Implications for Causal Gene Identification. <i>Arthritis and Rheumatology</i> , 2018 , 70, 361-370	9.5	26
120	Optimisation of methods for bacterial skin microbiome investigation: primer selection and comparison of the 454 versus MiSeq platform. <i>BMC Microbiology</i> , 2017 , 17, 23	4.5	71
119	The genetics revolution in rheumatology: large scale genomic arrays and genetic mapping. <i>Nature Reviews Rheumatology</i> , 2017 , 13, 421-432	8.1	18
118	The predictive value of serum S100A9 and response to etanercept is not confirmed in a large UK rheumatoid arthritis cohort. <i>Rheumatology</i> , 2017 , 56, 1019-1024	3.9	7
117	Genetics of Rheumatic Diseases 2017 , 327-343		

116	Genetic susceptibility to rheumatoid arthritis and its implications for novel drug discovery. <i>Expert Opinion on Drug Discovery</i> , 2016 , 11, 805-13	6.2	8
115	Brief Report: IRF4 Newly Identified as a Common Susceptibility Locus for Systemic Sclerosis and Rheumatoid Arthritis in a Cross-Disease Meta-Analysis of Genome-Wide Association Studies. <i>Arthritis and Rheumatology</i> , 2016 , 68, 2338-44	9.5	35
114	ImmunoChip Analyses of Epistasis in Rheumatoid Arthritis Confirm Multiple Interactions within MHC and Suggest Novel Non-MHC Epistatic Signals. <i>Journal of Rheumatology</i> , 2016 , 43, 839-45	4.1	6
113	One SNP at a Time: Moving beyond GWAS in Psoriasis. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 567-573	4.3	30
112	Loci associated with N-glycosylation of human IgG are not associated with rheumatoid arthritis: a Mendelian randomisation study. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 317-20	2.4	10
111	Identifying Causal Genes at the Multiple Sclerosis Associated Region 6q23 Using Capture Hi-C. <i>PLoS ONE</i> , 2016 , 11, e0166923	3.7	20
110	Capture Hi-C identifies a novel causal gene, IL20RA, in the pan-autoimmune genetic susceptibility region 6q23. <i>Genome Biology</i> , 2016 , 17, 212	18.3	62
109	Major histocompatibility complex harbors widespread genotypic variability of non-additive risk of rheumatoid arthritis including epistasis. <i>Scientific Reports</i> , 2016 , 6, 25014	4.9	11
108	A method to decipher pleiotropy by detecting underlying heterogeneity driven by hidden subgroups applied to autoimmune and neuropsychiatric diseases. <i>Nature Genetics</i> , 2016 , 48, 803-10	36.3	45
107	Differential Methylation as a Biomarker of Response to Etanercept in Patients With Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1353-60	9.5	46
106	Replication of Associations of Genetic Loci Outside the HLA Region With Susceptibility to Anti-Cyclic Citrullinated Peptide-Negative Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1603-13	9.5	24
105	High-density genotyping of immune loci in Koreans and Europeans identifies eight new rheumatoid arthritis risk loci. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, e13	2.4	76
104	The skin microbiome in psoriatic arthritis: methodology development and pilot data. <i>Lancet, The</i> , 2015 , 385 Suppl 1, S27	40	6
103	Genetics of RA susceptibility, what comes next?. <i>RMD Open</i> , 2015 , 1, e000028	5.9	9
102	A weighted genetic risk score using all known susceptibility variants to estimate rheumatoid arthritis risk. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 170-6	2.4	43
101	A large-scale genetic analysis reveals a strong contribution of the HLA class II region to giant cell arteritis susceptibility. <i>American Journal of Human Genetics</i> , 2015 , 96, 565-80	11	96
100	Widespread non-additive and interaction effects within HLA loci modulate the risk of autoimmune diseases. <i>Nature Genetics</i> , 2015 , 47, 1085-90	36.3	112
99	Investigating CD11c expression as a potential genomic biomarker of response to TNF inhibitor biologics in whole blood rheumatoid arthritis samples. <i>Arthritis Research and Therapy</i> , 2015 , 17, 359	5.7	5

98	Statistical colocalization of genetic risk variants for related autoimmune diseases in the context of common controls. <i>Nature Genetics</i> , 2015 , 47, 839-46	36.3	97
97	Novel rheumatoid arthritis susceptibility locus at 22q12 identified in an extended UK genome-wide association study. <i>Arthritis and Rheumatology</i> , 2014 , 66, 24-30	9.5	36
96	Genetics of rheumatoid arthritis contributes to biology and drug discovery. <i>Nature</i> , 2014 , 506, 376-81	50.4	1426
95	Investigation of an interleukin-6 receptor gene polymorphism (rs2228145) as a predictor of cardiovascular mortality in inflammatory polyarthritis: results from the Norfolk Arthritis Register. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 787-8	2.4	7
94	Fine mapping seronegative and seropositive rheumatoid arthritis to shared and distinct HLA alleles by adjusting for the effects of heterogeneity. <i>American Journal of Human Genetics</i> , 2014 , 94, 522-32	11	132
93	The bacterial skin microbiome in psoriatic arthritis, an unexplored link in pathogenesis: challenges and opportunities offered by recent technological advances. <i>Rheumatology</i> , 2014 , 53, 777-84	3.9	29
92	Take your PICS: moving from GWAS to immune function. <i>Immunity</i> , 2014 , 41, 883-5	32.3	5
91	Genetic analysis of the <i>Trichuris muris</i> -induced model of colitis reveals QTL overlap and a novel gene cluster for establishing colonic inflammation. <i>BMC Genomics</i> , 2013 , 14, 127	4.5	17
90	Enrichment of vitamin D response elements in RA-associated loci supports a role for vitamin D in the pathogenesis of RA. <i>Genes and Immunity</i> , 2013 , 14, 325-9	4.4	13
89	Common genetic variants associated with disease from genome-wide association studies are mutually exclusive in prostate cancer and rheumatoid arthritis. <i>BJU International</i> , 2013 , 111, 1148-55	5.6	7
88	Rare, low-frequency, and common variants in the protein-coding sequence of biological candidate genes from GWASs contribute to risk of rheumatoid arthritis. <i>American Journal of Human Genetics</i> , 2013 , 92, 15-27	11	72
87	Dense genotyping of immune-related disease regions identifies 14 new susceptibility loci for juvenile idiopathic arthritis. <i>Nature Genetics</i> , 2013 , 45, 664-9	36.3	256
86	Human genetics in rheumatoid arthritis guides a high-throughput drug screen of the CD40 signaling pathway. <i>PLoS Genetics</i> , 2013 , 9, e1003487	6	45
85	Predicting the risk of rheumatoid arthritis and its age of onset through modelling genetic risk variants with smoking. <i>PLoS Genetics</i> , 2013 , 9, e1003808	6	40
84	Identification of BACH2 and RAD51B as rheumatoid arthritis susceptibility loci in a meta-analysis of genome-wide data. <i>Arthritis and Rheumatism</i> , 2013 , 65, 3058-62		35
83	The potential use of expression profiling: implications for predicting treatment response in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1118-24	2.4	32
82	Identification of the tyrosine-protein phosphatase non-receptor type 2 as a rheumatoid arthritis susceptibility locus in europeans. <i>PLoS ONE</i> , 2013 , 8, e66456	3.7	24
81	High-density genetic mapping identifies new susceptibility loci for rheumatoid arthritis. <i>Nature Genetics</i> , 2012 , 44, 1336-40	36.3	436

80	Correlation of C-reactive protein haplotypes with serum C-reactive protein level and response to anti-tumor necrosis factor therapy in UK rheumatoid arthritis patients: results from the Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate cohort. <i>Arthritis Research and Therapy</i> , 2012 , 14, R214	5.7	16
79	A spectrum of susceptibility to rheumatoid arthritis within HLA-DRB1: stratification by autoantibody status in a large UK population. <i>Genes and Immunity</i> , 2012 , 13, 120-8	4.4	46
78	Genetic markers of rheumatoid arthritis susceptibility in anti-citrullinated peptide antibody negative patients. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1984-90	2.4	78
77	Informed conditioning on clinical covariates increases power in case-control association studies. <i>PLoS Genetics</i> , 2012 , 8, e1003032	6	58
76	Investigation of rheumatoid arthritis susceptibility loci in juvenile idiopathic arthritis confirms high degree of overlap. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1117-21	2.4	36
75	EULAR recommendations for terminology and research in individuals at risk of rheumatoid arthritis: report from the Study Group for Risk Factors for Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 638-41	2.4	263
74	Bayesian refinement of association signals for 14 loci in 3 common diseases. <i>Nature Genetics</i> , 2012 , 44, 1294-301	36.3	347
73	A genetic marker at the OLIG3/TNFAIP3 locus associates with methotrexate continuation in early inflammatory polyarthritis: results from the Norfolk Arthritis Register. <i>Pharmacogenomics Journal</i> , 2012 , 12, 128-33	3.5	12
72	The role of rheumatoid arthritis genetic susceptibility markers in the prediction of erosive disease in patients with early inflammatory polyarthritis: results from the Norfolk Arthritis Register. <i>Rheumatology</i> , 2011 , 50, 78-84	3.9	29
71	Subtype specific genetic associations for juvenile idiopathic arthritis: ERAP1 with the enthesitis related arthritis subtype and IL23R with juvenile psoriatic arthritis. <i>Arthritis Research and Therapy</i> , 2011 , 13, R12	5.7	52
70	Genetics of rheumatoid arthritis: GWAS and beyond. <i>Open Access Rheumatology: Research and Reviews</i> , 2011 , 3, 31-46	2.4	15
69	Case Study on Rheumatoid Arthritis 2011 , 307-323		
68	A systematic investigation of confirmed autoimmune loci in early-onset psoriasis reveals an association with IL2/IL21. <i>British Journal of Dermatology</i> , 2011 , 164, 660-4	4	20
67	HLA-DPB1-COL11A2 and three additional xMHC loci are independently associated with RA in a UK cohort. <i>Genes and Immunity</i> , 2011 , 12, 169-75	4.4	13
66	Examining the overlap between genome-wide rare variant association signals and linkage peaks in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1522-6		7
65	Evidence to support IL-13 as a risk locus for psoriatic arthritis but not psoriasis vulgaris. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1016-9	2.4	60
64	Genetic variants within the MAP kinase signalling network and anti-TNF treatment response in rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 98-103	2.4	48
63	The rheumatoid arthritis and juvenile idiopathic arthritis associated major (A) allele of rs2104286 is a loss of expression variant of IL2RA. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, A6-A6	2.4	

62	Study of the common genetic background for rheumatoid arthritis and systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 463-8	2.4	107
61	ASSIMILATOR: a new tool to inform selection of associated genetic variants for functional studies. <i>Bioinformatics</i> , 2011 , 27, 144-6	7.2	7
60	Association of the AFF3 gene and IL2/IL21 gene region with juvenile idiopathic arthritis. <i>Genes and Immunity</i> , 2010 , 11, 194-8	4.4	48
59	Association of the CCR5 gene with juvenile idiopathic arthritis. <i>Genes and Immunity</i> , 2010 , 11, 584-9	4.4	22
58	Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. <i>Nature</i> , 2010 , 464, 713-20	50.4	639
57	Meta-analysis and imputation refines the association of 15q25 with smoking quantity. <i>Nature Genetics</i> , 2010 , 42, 436-40	36.3	521
56	Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. <i>Nature Genetics</i> , 2010 , 42, 508-14	36.3	969
55	Combined effects of three independent SNPs greatly increase the risk estimate for RA at 6q23. <i>Human Molecular Genetics</i> , 2010 , 19, 4544-4544	5.6	78
54	PADI4 genotype is not associated with rheumatoid arthritis in a large UK Caucasian population. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 666-70	2.4	57
53	No evidence for association of the KLF12 gene with rheumatoid arthritis in a large UK cohort. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1407-8	2.4	4
52	Overlap of disease susceptibility loci for rheumatoid arthritis and juvenile idiopathic arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1049-53	2.4	53
51	Identification of AF4/FMR2 family, member 3 (AFF3) as a novel rheumatoid arthritis susceptibility locus and confirmation of two further pan-autoimmune susceptibility genes. <i>Human Molecular Genetics</i> , 2010 , 19, 4543-4543	5.6	78
50	Investigation of type 1 diabetes and coeliac disease susceptibility loci for association with juvenile idiopathic arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 2169-72	2.4	29
49	Association of CD40 with rheumatoid arthritis confirmed in a large UK case-control study. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 813-6	2.4	49
48	Overlapping genetic susceptibility variants between three autoimmune disorders: rheumatoid arthritis, type 1 diabetes and coeliac disease. <i>Arthritis Research and Therapy</i> , 2010 , 12, R175	5.7	79
47	Confirmation of association of the REL locus with rheumatoid arthritis susceptibility in the UK population. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1572-3	2.4	23
46	Rare variation at the TNFAIP3 locus and susceptibility to rheumatoid arthritis. <i>Human Genetics</i> , 2010 , 128, 627-33	6.3	27
45	Association of a rheumatoid arthritis susceptibility variant at the CCL21 locus with premature mortality in inflammatory polyarthritis patients. <i>Arthritis Care and Research</i> , 2010 , 62, 676-82	4.7	10

44	Identification of AF4/FMR2 family, member 3 (AFF3) as a novel rheumatoid arthritis susceptibility locus and confirmation of two further pan-autoimmune susceptibility genes. <i>Human Molecular Genetics</i> , 2009 , 18, 2518-22	5.6	70
43	Combined effects of three independent SNPs greatly increase the risk estimate for RA at 6q23. <i>Human Molecular Genetics</i> , 2009 , 18, 2693-9	5.6	77
42	Identification of a novel susceptibility locus for juvenile idiopathic arthritis by genome-wide association analysis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 258-63		60
41	Association of the IL2RA/CD25 gene with juvenile idiopathic arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 251-7		73
40	Reevaluation of the interaction between HLA-DRB1 shared epitope alleles, PTPN22, and smoking in determining susceptibility to autoantibody-positive and autoantibody-negative rheumatoid arthritis in a large UK Caucasian population. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2565-76		79
39	Outcomes of methotrexate therapy for psoriasis and relationship to genetic polymorphisms. <i>British Journal of Dermatology</i> , 2009 , 160, 438-41	4	55
38	Genome-wide association study identifies eight loci associated with blood pressure. <i>Nature Genetics</i> , 2009 , 41, 666-76	36.3	970
37	Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk. <i>Nature Genetics</i> , 2009 , 41, 1313-8	36.3	272
36	A re-evaluation of three putative functional single nucleotide polymorphisms in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1373-5	2.4	13
35	No association between polymorphisms in the interleukin-15 gene and early-onset psoriasis in a UK cohort suggests heterogeneity for this susceptibility locus identified in Chinese psoriasis patients. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 2904-5	4.3	5
34	Genetic variation in efflux transporters influences outcome to methotrexate therapy in patients with psoriasis. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 1925-9	4.3	88
33	Rheumatoid arthritis susceptibility loci at chromosomes 10p15, 12q13 and 22q13. <i>Nature Genetics</i> , 2008 , 40, 1156-9	36.3	125
32	Polymorphisms in the IL-12beta and IL-23R genes are associated with psoriasis of early onset in a UK cohort. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 1325-7	4.3	66
31	Polymorphisms in the PTPN22 region are associated with psoriasis of early onset. <i>British Journal of Dermatology</i> , 2008 , 158, 962-8	4	33
30	Association of the FCRL3 gene with rheumatoid arthritis: a further example of population specificity?. <i>Arthritis Research and Therapy</i> , 2008 , 10, 405	5.7	78
29	Re-evaluation of putative rheumatoid arthritis susceptibility genes in the post-genome wide association study era and hypothesis of a key pathway underlying susceptibility. <i>Human Molecular Genetics</i> , 2008 , 17, 2274-9	5.6	121
28	Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. <i>Nature Genetics</i> , 2007 , 39, 1329-37	36.3	1130
27	Rheumatoid arthritis association at 6q23. <i>Nature Genetics</i> , 2007 , 39, 1431-3	36.3	328

26	Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. <i>Nature</i> , 2007 , 447, 661-78	50.4	7801
25	Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. <i>Nature</i> , 2007 , 450, 887-92	50.4	421
24	Investigation of association between the TRAF family genes and RA susceptibility. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 1322-6	2.4	35
23	Investigation of genetic variation across the protein tyrosine phosphatase gene in patients with rheumatoid arthritis in the UK. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 683-6	2.4	26
22	Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. <i>Science</i> , 2007 , 316, 1336-41	33.3	1823
21	Investigation of the MHC2TA gene, associated with rheumatoid arthritis in a Swedish population, in a UK rheumatoid arthritis cohort. <i>Arthritis and Rheumatism</i> , 2006 , 54, 3417-22		20
20	Association of the FCRL3 gene with rheumatoid arthritis: a further example of population specificity?. <i>Arthritis Research and Therapy</i> , 2006 , 8, R117	5.7	27
19	Investigation of the SLC22A4 gene (associated with rheumatoid arthritis in a Japanese population) in a United Kingdom population of rheumatoid arthritis patients. <i>Arthritis and Rheumatism</i> , 2005 , 52, 752-8		31
18	Characterisation of the genomic architecture of human chromosome 17q and evaluation of different methods for haplotype block definition. <i>BMC Genetics</i> , 2005 , 6, 21	2.6	8
17	The type 1 diabetes susceptibility gene SUMO4 at IDDM5 is not associated with susceptibility to rheumatoid arthritis or juvenile idiopathic arthritis. <i>Rheumatology</i> , 2005 , 44, 1390-3	3.9	13
16	Investigation of polymorphisms in the PADI4 gene in determining severity of inflammatory polyarthritis. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64, 1311-5	2.4	28
15	Polymorphisms in the tumour necrosis factor gene are not associated with severity of inflammatory polyarthritis. <i>Annals of the Rheumatic Diseases</i> , 2004 , 63, 280-4	2.4	21
14	Association of protein kinase C alpha (PRKCA) gene with multiple sclerosis in a UK population. <i>Brain</i> , 2004 , 127, 1717-22	11.2	28
13	Polymorphisms of the equine major histocompatibility complex class II DRA locus. <i>Tissue Antigens</i> , 2004 , 64, 173-9		16
12	Investigation of susceptibility loci identified in the UK rheumatoid arthritis whole-genome scan in a further series of 217 UK affected sibling pairs. <i>Arthritis and Rheumatism</i> , 2004 , 50, 729-35		37
11	Haplotype analysis in simplex families and novel analytic approaches in a case-control cohort reveal no evidence of association of the CTLA-4 gene with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 748-52		48
10	A functional haplotype of the PADI4 gene associated with rheumatoid arthritis in a Japanese population is not associated in a United Kingdom population. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1117-21		170
9	Evidence for a novel rheumatoid arthritis susceptibility locus on chromosome 6p. <i>Arthritis and Rheumatism</i> , 2004 , 50, 3823-30		17

8	Whole-genome scan, in a complex disease, using 11,245 single-nucleotide polymorphisms: comparison with microsatellites. <i>American Journal of Human Genetics</i> , 2004 , 75, 54-64	11	198
7	Whole-genome linkage analysis of rheumatoid arthritis susceptibility loci in 252 affected sibling pairs in the United Kingdom. <i>Arthritis and Rheumatism</i> , 2002 , 46, 632-9		171
6	Two novel polymorphisms in the human transforming growth factor beta 2 gene. <i>Genes and Immunity</i> , 2001 , 2, 295-6	4.4	8
5	High resolution linkage and association mapping identifies a novel rheumatoid arthritis susceptibility locus homologous to one linked to two rat models of inflammatory arthritis. <i>Human Molecular Genetics</i> , 2001 , 10, 1901-6	5.6	47
4	Linkage of a marker in intron D of the estrogen synthase locus to rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1999 , 42, 1617-20		15
3	Autosomal dominant (Beukes) premature degenerative osteoarthropathy of the hip joint maps to an 11-cM region on chromosome 4q35. <i>American Journal of Human Genetics</i> , 1999 , 64, 904-8	11	44
2	Risk-taking behavior in adolescents: the paradigm. <i>Annals of the New York Academy of Sciences</i> , 1997 , 817, 1-35	6.5	55
1	Functional genomics atlas of synovial fibroblasts defining rheumatoid arthritis heritability		1