

Guowei Xia

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2997165/publications.pdf>

Version: 2024-02-01

20
papers

386
citations

840776

11
h-index

752698

20
g-index

23
all docs

23
docs citations

23
times ranked

683
citing authors

#	ARTICLE	IF	CITATIONS
1	The prognostic value of C-reactive protein in renal cell carcinoma: A systematic review and meta-analysis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 50.e1-50.e8.	1.6	58
2	Ki-67 is an independent indicator in non-muscle invasive bladder cancer (NMIBC); Combination of EORTC risk scores and Ki-67 expression could improve the risk stratification of NMIBC. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 42.e13-42.e19.	1.6	46
3	Genome-Wide Association Study of Bladder Cancer in a Chinese Cohort Reveals a New Susceptibility Locus at 5q12.3. <i>Cancer Research</i> , 2016, 76, 3277-3284.	0.9	46
4	Tumor-Associated Macrophages: A Potential Target for Cancer Therapy. <i>Frontiers in Oncology</i> , 2021, 11, 693517.	2.8	46
5	Periostin identified as a potential biomarker of prostate cancer by iTRAQ-proteomics analysis of prostate biopsy. <i>Proteome Science</i> , 2011, 9, 22.	1.7	37
6	COPB2 Is Upregulated in Prostate Cancer and Regulates PC-3 Cell Proliferation, Cell Cycle, and Apoptosis. <i>Archives of Medical Research</i> , 2016, 47, 411-418.	3.3	30
7	A functional variant in <i>TP63</i> at 3q28 associated with bladder cancer risk by creating an miR-140 binding site. <i>International Journal of Cancer</i> , 2016, 139, 65-74.	5.1	27
8	Human epidermal growth factor receptor 2: a significant indicator for predicting progression in non-muscle-invasive bladder cancer especially in high-risk groups. <i>World Journal of Urology</i> , 2015, 33, 1951-1957.	2.2	20
9	Epigallocatechin gallate inhibits the growth and promotes the apoptosis of bladder cancer cells. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 3513-3518.	1.8	15
10	Coatomer subunit beta 2 (COPB2), identified by label-free quantitative proteomics, regulates cell proliferation and apoptosis in human prostate carcinoma cells. <i>Biochemical and Biophysical Research Communications</i> , 2018, 495, 473-480.	2.1	15
11	Are EORTC risk tables suitable for Chinese patients with non-muscle-invasive bladder cancer?. <i>Cancer Epidemiology</i> , 2014, 38, 157-161.	1.9	13
12	Characterization and validation of long noncoding RNAs as new candidates in prostate cancer. <i>Cancer Cell International</i> , 2020, 20, 531.	4.1	7
13	Potential Alterations of Functional Connectivity Analysis in the Patients with Chronic Prostatitis/Chronic Pelvic Pain Syndrome. <i>Neural Plasticity</i> , 2021, 2021, 1-9.	2.2	5
14	Differential expression profiles of circRNAs in human prostate cancer based on chip and bioinformatic analysis. <i>International Journal of Clinical and Experimental Pathology</i> , 2020, 13, 1045-1052.	0.5	5
15	Retroperitoneal Laparoscopic Ureterolithotomy for Proximal Ureteral Calculi in Selected Patients. <i>Scientific World Journal</i> , The, 2014, 2014, 1-5.	2.1	4
16	Postoperative renormalization of C-reactive protein with adjuvant lienal polypeptide and its association with tumour recurrence in T1 clear cell renal cell carcinoma. <i>Journal of International Medical Research</i> , 2016, 44, 620-626.	1.0	4
17	NOS3 895G>T and CBR3 730G>A Are Associated with Recurrence Risk in Non-Muscle-Invasive Bladder Cancer with Intravesical Instillations of THP. <i>Chemotherapy</i> , 2018, 63, 191-197.	1.6	3
18	High-level expression of periostin is significantly correlated with tumor angiogenesis in prostate cancer. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 1569-1574.	0.5	2

#	ARTICLE	IF	CITATIONS
19	Lysine demethylase 5A promotes prostate adenocarcinoma progression by suppressing microRNA-330-3p expression and activating the COPB2/PI3K/AKT axis in an ETS1-dependent manner. <i>Journal of Cell Communication and Signaling</i> , 2022, 16, 579-599.	3.4	2
20	A Cumulative Analysis of Current Evidence for Association between Expression of Epithelial-Mesenchymal Transition Markers and Clinicopathological Outcomes in Patients after Radical Prostatectomy. <i>Annals of Clinical and Laboratory Science</i> , 2018, 48, 18-28.	0.2	1