

Per G Farup

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/299987/per-g-farup-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

83
papers

1,915
citations

25
h-index

40
g-index

92
ext. papers

2,278
ext. citations

3.2
avg, IF

5
L-index

#	Paper	IF	Citations
83	The Peripheral Kynurenine Pathway and Psychosomatic Comorbidity in Subjects with Morbid Obesity Undergoing Bariatric Surgery. <i>Applied Sciences (Switzerland)</i> , 2022 , 12, 2648	2.6	1
82	Changes in Faecal Short-Chain Fatty Acids after Weight-Loss Interventions in Subjects with Morbid Obesity. <i>Nutrients</i> , 2020 , 12,	6.7	13
81	Are the Results of a Combined Behavioural and Surgical Treatment of Morbid Obesity Satisfactory and Predictable?. <i>Nutrients</i> , 2020 , 12,	6.7	2
80	APOE - a genetic marker of comorbidity in subjects with morbid obesity. <i>BMC Medical Genetics</i> , 2020 , 21, 146	2.1	4
79	Changes in serum zonulin in individuals with morbid obesity after weight-loss interventions: a prospective cohort study. <i>BMC Endocrine Disorders</i> , 2020 , 20, 108	3.3	9
78	Effects of a fall prevention program in elderly: a pragmatic observational study in two orthopedic departments. <i>Clinical Interventions in Aging</i> , 2019 , 14, 145-154	4	9
77	Are Nonnutritive Sweeteners Obesogenic? Associations between Diet, Faecal Microbiota, and Short-Chain Fatty Acids in Morbidly Obese Subjects. <i>Journal of Obesity</i> , 2019 , 2019, 4608315	3.7	8
76	Irritable bowel syndrome-like symptoms and health related quality of life two years after Roux-en-Y gastric bypass - a prospective cohort study. <i>BMC Gastroenterology</i> , 2019 , 19, 204	3	5
75	Changes in C-reactive protein, neopterin and lactoferrin differ after conservative and surgical weight loss in individuals with morbid obesity. <i>Scientific Reports</i> , 2019 , 9, 17695	4.9	8
74	Evaluation of a faecal dysbiosis test for irritable bowel syndrome in subjects with and without obesity. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018 , 78, 109-113	2	9
73	Is perceived intolerance to milk and wheat associated with the corresponding IgG and IgA food antibodies? A cross sectional study in subjects with morbid obesity and gastrointestinal symptoms. <i>BMC Gastroenterology</i> , 2018 , 18, 22	3	2
72	Separating "good" from "bad" faecal dysbiosis - evidence from two cross-sectional studies. <i>BMC Obesity</i> , 2018 , 5, 30	3.6	5
71	Changes in gastrointestinal symptoms and food tolerance 6 months following weight loss surgery: associations with dietary changes, weight loss and the surgical procedure. <i>BMC Obesity</i> , 2018 , 5, 29	3.6	5
70	Faecal Microbial Markers and Psychobiological Disorders in Subjects with Morbid Obesity. A Cross-Sectional Study. <i>Behavioral Sciences (Basel, Switzerland)</i> , 2018 , 8,	2.3	2
69	A Conservative Weight Loss Intervention Relieves Bowel Symptoms in Morbidly Obese Subjects with Irritable Bowel Syndrome: A Prospective Cohort Study. <i>Journal of Obesity</i> , 2018 , 2018, 3732753	3.7	12
68	Functional Bowel Disorders Are Associated with a Central Immune Activation. <i>Gastroenterology Research and Practice</i> , 2017 , 2017, 1642912	2	8
67	Anti- antibodies (ASCA) are associated with body fat mass and systemic inflammation, but not with dietary yeast consumption: a cross-sectional study. <i>BMC Obesity</i> , 2017 , 4, 28	3.6	7

66	Intake of non-nutritive sweeteners is associated with an unhealthy lifestyle: a cross-sectional study in subjects with morbid obesity. <i>BMC Obesity</i> , 2017 , 4, 41	3.6	9
65	Prevalence and predictors of irritable bowel syndrome in patients with morbid obesity: a cross-sectional study. <i>BMC Obesity</i> , 2017 , 4, 22	3.6	15
64	The Relationships among Tryptophan, Kynurenine, Indoleamine 2,3-Dioxygenase, Depression, and Neuropsychological Performance. <i>Frontiers in Psychology</i> , 2017 , 8, 1561	3.4	37
63	Changes in patient safety culture after restructuring of intensive care units: Two cross-sectional studies. <i>Intensive and Critical Care Nursing</i> , 2016 , 32, 58-65	3.1	11
62	The effect of ApoE e4 on blood pressure in patients with and without depression. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 1365-70	3.1	4
61	Patients with depression display cytokine levels in serum and cerebrospinal fluid similar to patients with diffuse neurological symptoms without a defined diagnosis. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 817-22	3.1	20
60	Faecal short-chain fatty acids - a diagnostic biomarker for irritable bowel syndrome?. <i>BMC Gastroenterology</i> , 2016 , 16, 51	3	51
59	The association between patient safety culture and burnout and sense of coherence: A cross-sectional study in restructured and not restructured intensive care units. <i>Intensive and Critical Care Nursing</i> , 2016 , 36, 26-34	3.1	34
58	Cognitive Functions and Depression in Patients with Irritable Bowel Syndrome. <i>Gastroenterology Research and Practice</i> , 2015 , 2015, 438329	2	8
57	Are measurements of patient safety culture and adverse events valid and reliable? Results from a cross sectional study. <i>BMC Health Services Research</i> , 2015 , 15, 186	2.9	22
56	Predictors of health-related quality of life in patients with irritable bowel syndrome. A cross-sectional study in Norway. <i>Health and Quality of Life Outcomes</i> , 2015 , 13, 113	3	16
55	The Culture of Incident Reporting and Feedback: A Cross-Sectional Study in a Hospital Setting. <i>Open Journal of Nursing</i> , 2015 , 05, 1042-1052	0.3	1
54	Differences in medication knowledge and risk of errors between graduating nursing students and working registered nurses: comparative study. <i>BMC Health Services Research</i> , 2014 , 14, 580	2.9	38
53	Improvement of drug dose calculations by classroom teaching or e-learning: a randomised controlled trial in nurses. <i>BMJ Open</i> , 2014 , 4, e006025	3	16
52	Vitamin B6 deficiency and diseases in elderly people--a study in nursing homes. <i>BMC Geriatrics</i> , 2013 , 13, 13	4.1	28
51	Vitamin B6 in plasma - sample stability and the reference limits. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2013 , 73, 476-9	2	11
50	IgG and IgG4 antibodies in subjects with irritable bowel syndrome: a case control study in the general population. <i>BMC Gastroenterology</i> , 2012 , 12, 166	3	25
49	Diet in subjects with irritable bowel syndrome: a cross-sectional study in the general population. <i>BMC Gastroenterology</i> , 2012 , 12, 61	3	26

48	Probiotics, symptoms, and gut microbiota: what are the relations? A randomized controlled trial in subjects with irritable bowel syndrome. <i>Gastroenterology Research and Practice</i> , 2012 , 2012, 214102	2	22
47	Drugs and constipation in elderly in nursing homes: what is the relation?. <i>Gastroenterology Research and Practice</i> , 2012 , 2012, 290231	2	33
46	Irritable bowel syndrome. <i>Gastroenterology Research and Practice</i> , 2012 , 2012, 612479	2	2
45	Low intake of vitamin B6 is associated with irritable bowel syndrome symptoms. <i>Nutrition Research</i> , 2011 , 31, 356-61	4	28
44	Questions about mesalazine and the irritable bowel syndrome. <i>Alimentary Pharmacology and Therapeutics</i> , 2011 , 34, 1036-7; author reply 1037-8	6.1	1
43	Effectiveness of laxatives in elderly--a cross sectional study in nursing homes. <i>BMC Geriatrics</i> , 2011 , 11, 76	4.1	11
42	Constipation and diarrhoea - common adverse drug reactions? A cross sectional study in the general population. <i>BMC Clinical Pharmacology</i> , 2011 , 11, 2		39
41	What causes treatment failure - the patient, primary care, secondary care or inadequate interaction in the health services?. <i>BMC Health Services Research</i> , 2011 , 11, 111	2.9	8
40	Medication knowledge, certainty, and risk of errors in health care: a cross-sectional study. <i>BMC Health Services Research</i> , 2011 , 11, 175	2.9	37
39	A candidate probiotic with unfavourable effects in subjects with irritable bowel syndrome: a randomised controlled trial. <i>BMC Gastroenterology</i> , 2010 , 10, 16	3	65
38	How does comorbidity affect cost of health care in patients with irritable bowel syndrome? A cohort study in general practice. <i>BMC Gastroenterology</i> , 2010 , 10, 31	3	28
37	Burnout in health-care professionals during reorganizations and downsizing. A cohort study in nurses. <i>BMC Nursing</i> , 2010 , 9, 8	3.2	27
36	Alternative vs. conventional treatment given on-demand for gastroesophageal reflux disease: a randomised controlled trial. <i>BMC Complementary and Alternative Medicine</i> , 2009 , 9, 3	4.7	3
35	A comparison of diagnostic tests for lactose malabsorption--which one is the best?. <i>BMC Gastroenterology</i> , 2009 , 9, 82	3	39
34	Disease knowledge after an educational program in patients with GERD--a randomized controlled trial. <i>BMC Health Services Research</i> , 2008 , 8, 236	2.9	8
33	Patient education in gastro-oesophageal reflux disease: a randomized controlled trial. <i>European Journal of Gastroenterology and Hepatology</i> , 2007 , 19, 1104-10	2.2	11
32	Prevalence, comorbidity and impact of irritable bowel syndrome in Norway. <i>Scandinavian Journal of Gastroenterology</i> , 2006 , 41, 650-6	2.4	102
31	Digestive symptoms and their psychosocial impact: validation of a questionnaire. <i>Scandinavian Journal of Gastroenterology</i> , 2006 , 41, 1019-27	2.4	5

30	Perceived food intolerance in subjects with irritable bowel syndrome-- etiology, prevalence and consequences. <i>European Journal of Clinical Nutrition</i> , 2006 , 60, 667-72	5.2	249
29	How useful are the Rome II criteria for identification of upper gastrointestinal disorders in general practice?. <i>Scandinavian Journal of Gastroenterology</i> , 2005 , 40, 1284-9	2.4	9
28	The value of a general therapeutic approach in subjects with irritable bowel syndrome. <i>Alimentary Pharmacology and Therapeutics</i> , 2005 , 21, 21-7	6.1	11
27	Abdominal complaints in general practice: diagnoses and characteristics of patients. <i>Scandinavian Journal of Primary Health Care</i> , 2005 , 23, 126; author reply 126-7	2.7	
26	In response to Montes and Monge. <i>Scandinavian Journal of Gastroenterology</i> , 2004 , 39, 1033-1033	2.4	
25	Comorbidity of irritable bowel syndrome in general practice: a striking feature with clinical implications. <i>Alimentary Pharmacology and Therapeutics</i> , 2004 , 20, 1195-203	6.1	108
24	Tegaserod--merely a laxative?. <i>Scandinavian Journal of Gastroenterology</i> , 2004 , 39, 404-5	2.4	1
23	Abdominal complaints in general practice. <i>Scandinavian Journal of Primary Health Care</i> , 2004 , 22, 157-62	2.7	16
22	Diagnosing irritable bowel syndrome: poor agreement between general practitioners and the Rome II criteria. <i>Scandinavian Journal of Gastroenterology</i> , 2004 , 39, 448-53	2.4	47
21	Lactose malabsorption in a population with irritable bowel syndrome: prevalence and symptoms. A case-control study. <i>Scandinavian Journal of Gastroenterology</i> , 2004 , 39, 645-9	2.4	87
20	Collaboration by use of the Internet yields data of high quality and detects non-uniform management of patients with <i>Helicobacter pylori</i> infection. <i>Scandinavian Journal of Gastroenterology</i> , 2002 , 37, 1466-70	2.4	2
19	The effect of <i>Helicobacter pylori</i> retreatment with ranitidine bismuth citrate, clarithromycin, and metronidazole depends on the first-line therapy. <i>Journal of Clinical Gastroenterology</i> , 2002 , 35, 379-82	3	6
18	Comparison of three triple regimens with omeprazole or ranitidine bismuth citrate for <i>Helicobacter pylori</i> eradication. <i>Scandinavian Journal of Gastroenterology</i> , 2002 , 37, 1374-9	2.4	21
17	Mesalazine 4 g daily given as prolonged-release granules twice daily and four times daily is at least as effective as prolonged-release tablets four times daily in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2001 , 7, 237-42	4.5	50
16	Does short-term treatment with proton pump inhibitors cause rebound aggravation of symptoms?. <i>Journal of Clinical Gastroenterology</i> , 2001 , 33, 206-9	3	14
15	Patients with functional dyspepsia responding to omeprazole have a characteristic gastro-oesophageal reflux pattern. <i>Scandinavian Journal of Gastroenterology</i> , 1999 , 34, 575-9	2.4	46
14	The symptomatic effect of cisapride in patients with irritable bowel syndrome and constipation. <i>Scandinavian Journal of Gastroenterology</i> , 1998 , 33, 128-31	2.4	43
13	Mesalazine suppositories versus hydrocortisone foam in patients with distal ulcerative colitis. A comparison of the efficacy and practicality of two topical treatment regimens. <i>Scandinavian Journal of Gastroenterology</i> , 1995 , 30, 164-70	2.4	70

12	Are frequent short gastro-oesophageal reflux episodes the cause of symptoms in patients with non-ulcer dyspepsia responding to treatment with ranitidine?. <i>Scandinavian Journal of Gastroenterology</i> , 1995 , 30, 829-32	2.4	20
11	Long-term treatment with 300 mg ranitidine once daily after dilatation of peptic oesophageal strictures. <i>Scandinavian Journal of Gastroenterology</i> , 1992 , 27, 594-8	2.4	19
10	The multi-crossover model for classifying patients as responders to a given treatment. <i>Scandinavian Journal of Gastroenterology</i> , 1991 , 26, 763-70	2.4	17
9	Effect of cisapride on symptoms and biliary drainage in patients with postcholecystectomy syndrome. <i>Scandinavian Journal of Gastroenterology</i> , 1991 , 26, 945-50	2.4	8
8	Ranitidine for non-ulcer dyspepsia. A clinical study of the symptomatic effect of ranitidine and a classification and characterization of the responders to treatment. <i>Scandinavian Journal of Gastroenterology</i> , 1991 , 26, 1209-16	2.4	27
7	Duodenal ulcer treated with omeprazole: healing and relapse rates. Does treatment duration influence subsequent remission?. <i>Scandinavian Journal of Gastroenterology</i> , 1989 , 24, 1107-12	2.4	11
6	Sphincter of Oddi dysfunction. Dynamic cholescintigraphy and endoscopic retrograde cholangiopancreatography with papillotomy in diagnosis, treatment, and follow-up study. <i>Scandinavian Journal of Gastroenterology</i> , 1989 , 24, 956-60	2.4	15
5	Zaltidine: an effective but hepatotoxic H ₂ -receptor antagonist. <i>Scandinavian Journal of Gastroenterology</i> , 1988 , 23, 655-8	2.4	11
4	Low-dose antacids or cimetidine for duodenal ulcer?. <i>Gastroenterology</i> , 1988 , 95, 1465-9	13.3	25
3	Rate of Healing of Benign Gastric Ulcer During Treatment with Omeprazole. <i>Scandinavian Journal of Gastroenterology</i> , 1986 , 21, 180-180	2.4	1
2	The rate of healing of duodenal ulcers during omeprazole treatment. <i>Scandinavian Journal of Gastroenterology</i> , 1985 , 20, 691-5	2.4	20
1	Idiopathic Hypoparathyroidism and Hyperthyroidism. <i>Acta Medica Scandinavica</i> , 1977 , 202, 261-4		4