

# Kozo Tanno

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3000405/publications.pdf>

Version: 2024-02-01

124  
papers

3,324  
citations

201385

27  
h-index

189595

50  
g-index

130  
all docs

130  
docs citations

130  
times ranked

6081  
citing authors

#	ARTICLE	IF	CITATIONS
1	Long-term Response of <i>Helicobacter pylori</i> Antibody Titer After Eradication Treatment in Middle-aged Japanese: JPHC-NEXT Study. <i>Journal of Epidemiology</i> , 2023, 33, 1-7.	1.1	3
2	Association Between Birth Weight and Risk of Pregnancy-Induced Hypertension and Gestational Diabetes in Japanese Women: JPHC-NEXT Study. <i>Journal of Epidemiology</i> , 2022, 32, 168-173.	1.1	6
3	Psychological Distress in Responders and Nonresponders in a 5-year Follow-up Health Survey: The RIAS Study. <i>Journal of Epidemiology</i> , 2022, 32, 527-534.	1.1	3
4	Combined Associations of Liver Enzymes and Obesity With Diabetes Mellitus Prevalence: The Tohoku Medical Megabank Community-based Cohort Study. <i>Journal of Epidemiology</i> , 2022, 32, 221-227.	1.1	9
5	The Potential of a Stroke Registry Using Diagnosis Procedure Combination Data from All Hospitals in a Japanese Prefecture. <i>Cerebrovascular Diseases</i> , 2022, 51, 447-452.	0.8	4
6	Risk Factors of Long-Term Care Insurance Certification in Japan: A Scoping Review. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 2162.	1.2	5
7	Association between Social Isolation and Total Mortality after the Great East Japan Earthquake in Iwate Prefecture: Findings from the TMM CommCohort Study. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 4343.	1.2	1
8	Association between total type I collagen N-terminal propeptide and coronary artery disease risk score in the general Japanese population. <i>IJC Heart and Vasculature</i> , 2022, 41, 101056.	0.6	2
9	Slightly Elevated Serum Ionic Fluoride Levels Inhibit Insulin Secretion and Increase Glucose Levels in a General Japanese Population: a Cross-sectional Study. <i>Biological Trace Element Research</i> , 2021, 199, 2819-2825.	1.9	5
10	Study Profile of the Tohoku Medical Megabank Community-Based Cohort Study. <i>Journal of Epidemiology</i> , 2021, 31, 65-76.	1.1	81
11	Telomere shortening as a stress-related biomarker in children exposed to maternal chronic stress in utero measured 7 years after the Great East Japan Earthquake. <i>Psychiatry Research</i> , 2021, 295, 113565.	1.7	4
12	Five-year blood pressure trajectories of survivors of the tsunami following the Great East Japan Earthquake in Iwate. <i>Hypertension Research</i> , 2021, 44, 581-590.	1.5	2
13	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. <i>Journal of Cancer Survivorship</i> , 2021, 15, 912-921.	1.5	9
14	Body mass index and colorectal cancer risk: A Mendelian randomization study. <i>Cancer Science</i> , 2021, 112, 1579-1588.	1.7	25
15	Plasma Xanthine Oxidoreductase Activity Is Associated with a High Risk of Cardiovascular Disease in a General Japanese Population. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 1894.	1.2	9
16	GWAS Identified IL4R and the Major Histocompatibility Complex Region as the Associated Loci of Total Serum IgE Levels in 9,260 Japanese Individuals. <i>Journal of Investigative Dermatology</i> , 2021, 141, 2749-2752.	0.3	4
17	Weight Gain After 20 Years of Age is Associated with Unfavorable Lifestyle and Increased Prevalence of Metabolic Disorders. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2021, Volume 14, 2065-2075.	1.1	7
18	Increase in Body Weight Following Residential Displacement: 5-year Follow-up After the 2011 Great East Japan Earthquake and Tsunami. <i>Journal of Epidemiology</i> , 2021, 31, 328-334.	1.1	4

#	ARTICLE	IF	CITATIONS
19	Association between the social isolation and depressive symptoms after the great East Japan earthquake: findings from the baseline survey of the TMM CommCohort study. BMC Public Health, 2021, 21, 925.	1.2	14
20	Low educational level increases functional disability risk subsequent to heart failure in Japan: On behalf of the Iwate KENCO study group. PLoS ONE, 2021, 16, e0253017.	1.1	0
21	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. Ocular Surface, 2021, 21, 306-312.	2.2	14
22	Utility of urinary albumin excretion as an index for stratifying the residual cardiovascular risk in patients undergoing antihypertensive agents treatment. Journal of Hypertension, 2021, Publish Ahead of Print, 2431-2438.	0.3	0
23	Relation Between Body Mass Index and Dry Eye Disease: The Japan Public Health Center-Based Prospective Study for the Next Generation. Eye and Contact Lens, 2021, 47, 449-455.	0.8	8
24	Age-Specific Incidence Rates of Acute Coronary Syndrome in Japan—A Special Gift From a Collaboration of Cardiologists and Epidemiologists. Circulation Journal, 2021, 85, 1731-1734.	0.7	1
25	Association between Milk Intake and Incident Stroke among Japanese Community Dwellers: The Iwate-KENCO Study. Nutrients, 2021, 13, 3781.	1.7	2
26	The prevalence of current smokers and alcohol drinkers among cancer survivors and subjects with no history of cancer among participants in a community-based cardiometabolic screening program in Miyagi prefecture, Japan: a comparison with nationally representative surveys in other countries. Cancer Medicine, 2021, , .	1.3	5
27	How do patients with chronic illnesses respond to a public health crisis? Evidence from diabetic patients in Japan during the COVID-19 pandemic. SSM - Population Health, 2021, 16, 100961.	1.3	2
28	Public access to summary statistics for genome-wide association studies of body mass index, weight, and height among healthy Japanese individuals: the Japanese Consortium of Genetic Epidemiology studies. Journal of Epidemiology, 2021, , .	1.1	0
29	Longitudinal DNA methylation dynamics as a practical indicator in clinical epigenetics. Clinical Epigenetics, 2021, 13, 219.	1.8	10
30	dbTMM: an integrated database of large-scale cohort, genome and clinical data for the Tohoku Medical Megabank Project. Human Genome Variation, 2021, 8, 44.	0.4	7
31	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. Journal of Epidemiology, 2020, 30, 46-54.	1.1	30
32	The Value of a Cystatin C-based Estimated Glomerular Filtration Rate for Cardiovascular Assessment in a General Japanese Population: Results From the Iwate Tohoku Medical Megabank Project. Journal of Epidemiology, 2020, 30, 260-267.	1.1	5
33	Cohort Profile: Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study (TMM) Tj ETQq1 1 0.784314 rgBT /Overl... 2020, 49, 18-19m.	0.9	107
34	Physical inactivity, prolonged sedentary behaviors, and use of visual display terminals as potential risk factors for dry eye disease: JPHC-NEXT study. Ocular Surface, 2020, 18, 56-63.	2.2	42
35	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. Journal of Epidemiology, 2020, 30, 396-403.	1.1	11
36	Ten-Year Cerebrovascular Disease Trend Occurrence by Population-based Stroke Registry in an Aging Japan Local Prefecture. Journal of Stroke and Cerebrovascular Diseases, 2020, 29, 104580.	0.7	6

#	ARTICLE	IF	CITATIONS
37	Increased incidence of metabolic syndrome among older survivors relocated to temporary housing after the 2011 Great East Japan earthquake & tsunami. <i>Metabolism Open</i> , 2020, 7, 100042.	1.4	7
38	Effect of temporary housing on incidence of diabetes mellitus in survivors of a tsunami-stricken area in 2011 Japan disaster: a serial cross-sectional RIAS study. <i>Scientific Reports</i> , 2020, 10, 15400.	1.6	3
39	Poor self-rated health predicts the incidence of functional disability in elderly community dwellers in Japan: a prospective cohort study. <i>BMC Geriatrics</i> , 2020, 20, 328.	1.1	12
40	Machine learning for effectively avoiding overfitting is a crucial strategy for the genetic prediction of polygenic psychiatric phenotypes. <i>Translational Psychiatry</i> , 2020, 10, 294.	2.4	11
41	Occurrence of Cerebrovascular Diseases Decreased after the Great East Japan Earthquake and Tsunami of 2011. <i>Cerebrovascular Diseases Extra</i> , 2020, 10, 105-115.	0.5	1
42	Hypertensive disorders of pregnancy, obesity, and hypertension in later life by age group: a cross-sectional analysis. <i>Hypertension Research</i> , 2020, 43, 1277-1283.	1.5	14
43	Additional prognostic value of electrocardiographic left ventricular hypertrophy in traditional cardiovascular risk assessments in chronic kidney disease. <i>Journal of Hypertension</i> , 2020, 38, 1149-1157.	0.3	4
44	Large-scale genome-wide association study in a Japanese population identifies novel susceptibility loci across different diseases. <i>Nature Genetics</i> , 2020, 52, 669-679.	9.4	304
45	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. <i>Nature Communications</i> , 2020, 11, 3175.	5.8	34
46	Genome-wide association study identifies new loci for albuminuria in the Japanese population. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 1-9.	0.7	9
47	Abstract P539: The Association Between Social Isolation and Metabolic Syndrome in Japanese Adults: The Tohoku Medical Megabank Project. <i>Circulation</i> , 2020, 141, .	1.6	0
48	Abstract 3486: Body mass index and colorectal cancer risk in Japanese populations: a Mendelian randomization study. , 2020, , .		0
49	Abstract P483: Living Arrangement Affects Urine Na/k Ratio Among 44,085 Residents: Tohoku Medical Megabank Project Community-based Cohort Study. <i>Circulation</i> , 2020, 141, .	1.6	0
50	12 new susceptibility loci for prostate cancer identified by genome-wide association study in Japanese population. <i>Nature Communications</i> , 2019, 10, 4422.	5.8	49
51	Characterizing rare and low-frequency height-associated variants in the Japanese population. <i>Nature Communications</i> , 2019, 10, 4393.	5.8	123
52	Lack of social support and social trust as potential risk factors for dry eye disease: JPHC-NEXT study. <i>Ocular Surface</i> , 2019, 17, 278-284.	2.2	3
53	Susceptibility Loci for Tanning Ability in the Japanese Population Identified by a Genome-Wide Association Study from the Tohoku Medical Megabank Project Cohort Study. <i>Journal of Investigative Dermatology</i> , 2019, 139, 1605-1608.e13.	0.3	14
54	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep duration in the Japanese population. <i>Sleep</i> , 2019, 42, .	0.6	16

#	ARTICLE	IF	CITATIONS
55	Association between high-sensitivity cardiac troponin T and future cardiovascular incidence in a general Japanese population: results from the Tohoku medical megabank project. <i>Biomarkers</i> , 2019, 24, 566-573.	0.9	10
56	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. <i>Nature Genetics</i> , 2019, 51, 379-386.	9.4	164
57	Abstract P031: Plasma Xanthine Oxidoreductase Activity is Associated With Endothelial Dysfunction and Increased Arterial Stiffness in a General Japanese Population: the Iwate Tohoku Medical Megabank Project. <i>Circulation</i> , 2019, 139, .	1.6	0
58	Abstract P331: Unfavorable Social Factors Increased the Risk of Future Disability Subsequent to Heart Failure in Elderly Japanese Community Dwellers: The Iwate KENCO Study Group. <i>Circulation</i> , 2019, 139, .	1.6	0
59	Abstract P086: Significant Association of House Collapse With High Urine Na/K Ratios Among 24,580 Coastal Residents Who Survived the Great East Japan Earthquake. <i>Circulation</i> , 2019, 139, .	1.6	0
60	Abstract P401: Female Gender is a Significant Risk Factor for Functional Disability After Hemorrhagic Stroke but Not Ischemic Stroke Among Japanese Community-Dwelling Elderly Individuals. <i>Circulation</i> , 2019, 139, .	1.6	0
61	Burden of high blood pressure as a contributing factor to stroke in the Japanese community-based diabetic population. <i>Hypertension Research</i> , 2018, 41, 531-538.	1.5	9
62	Prognostic Value of Electrocardiographic Left Ventricular Hypertrophy on Cardiovascular Risk in a Non-Hypertensive Community-Based Population. <i>American Journal of Hypertension</i> , 2018, 31, 895-901.	1.0	12
63	Genome-wide analysis of polymorphism-sodium interaction effect on blood pressure identifies a novel 3' BCL11B gene desert locus. <i>Scientific Reports</i> , 2018, 8, 14162.	1.6	10
64	Ability of B-Type Natriuretic Peptide Testing to Predict Cardioembolic Stroke in the General Population—Comparisons With C-Reactive Protein and Urinary Albumin. <i>Circulation Journal</i> , 2018, 82, 1017-1025.	0.7	13
65	The urine albumin-creatinine ratio is a predictor for incident long-term care in a general population. <i>PLoS ONE</i> , 2018, 13, e0195013.	1.1	8
66	Risk of stroke and heart failure attributable to atrial fibrillation in middle-aged and elderly people: Results from a five-year prospective cohort study of Japanese community dwellers. <i>Journal of Epidemiology</i> , 2017, 27, 360-367.	1.1	11
67	The Impact and Effectivity of an Inventory Survey for a Stroke Registry in Iwate Prefecture. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017, 26, 2160-2166.	0.7	3
68	Genetic Predisposition to Ischemic Stroke. <i>Stroke</i> , 2017, 48, 253-258.	1.0	64
69	Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. <i>Nature Genetics</i> , 2017, 49, 1458-1467.	9.4	380
70	Genome-wide identification of inter-individually variable DNA methylation sites improves the efficacy of epigenetic association studies. <i>Npj Genomic Medicine</i> , 2017, 2, 11.	1.7	59
71	Population-based biobank participants' preferences for receiving genetic test results. <i>Journal of Human Genetics</i> , 2017, 62, 1037-1048.	1.1	24
72	Genome-wide meta-analysis in Japanese populations identifies novel variants at the TMC6 and SIX3/SIX2 loci associated with HbA1c. <i>Scientific Reports</i> , 2017, 7, 16147.	1.6	28

#	ARTICLE	IF	CITATIONS
73	Comparison between urine albumin-to-creatinine ratio and urine protein dipstick testing for prevalence and ability to predict the risk for chronic kidney disease in the general population (Iwate-KENCO study): a prospective community-based cohort study. <i>BMC Nephrology</i> , 2016, 17, 46.	0.8	11
74	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. <i>PLoS ONE</i> , 2016, 11, e0147519.	1.1	21
75	Low-grade albuminuria and incidence of cardiovascular disease and all-cause mortality in nondiabetic and normotensive individuals. <i>Journal of Hypertension</i> , 2016, 34, 506-512.	0.3	53
76	Sustained Increase in the Incidence of Acute Decompensated Heart Failure After the 2011 Japan Earthquake and Tsunami. <i>American Journal of Cardiology</i> , 2016, 118, 1374-1379.	0.7	12
77	Standardized Prevalence Ratios for Atrial Fibrillation in Adult Dialysis Patients in Japan. <i>Journal of Epidemiology</i> , 2016, 26, 272-276.	1.1	9
78	The Tohoku Medical Megabank Project: Design and Mission. <i>Journal of Epidemiology</i> , 2016, 26, 493-511.	1.1	236
79	The influence of personality and perceived stress on the development of breast cancer: 20-year follow-up of 29,098 Japanese women. <i>Scientific Reports</i> , 2016, 6, 32559.	1.6	16
80	Association between relocation and changes in cardiometabolic risk factors: a longitudinal study in tsunami survivors of the 2011 Great East Japan Earthquake. <i>BMJ Open</i> , 2016, 6, e011291.	0.8	26
81	Weight Gain in Survivors Living in Temporary Housing in the Tsunami-Stricken Area during the Recovery Phase following the Great East Japan Earthquake and Tsunami. <i>PLoS ONE</i> , 2016, 11, e0166817.	1.1	18
82	Relative and absolute risks of all-cause and cause-specific deaths attributable to atrial fibrillation in middle-aged and elderly community dwellers. <i>International Journal of Cardiology</i> , 2015, 184, 692-698.	0.8	13
83	Elevated blood pressure at the first measurement predicts cardiovascular disease independently from the subsequent second reading in men, but not in women. <i>Clinical and Experimental Hypertension</i> , 2015, 37, 39-44.	0.5	2
84	Associations of number of teeth with risks for all-cause mortality and cause-specific mortality in middle-aged and elderly men in the northern part of Japan: the Iwate-KENCO study. <i>Community Dentistry and Oral Epidemiology</i> , 2014, 42, 358-365.	0.9	22
85	Extent of Flood Damage Increased Cerebrovascular Disease Incidences in Iwate Prefecture after the Great East Japan Earthquake and Tsunami of 2011. <i>Cerebrovascular Diseases</i> , 2014, 37, 451-459.	0.8	7
86	Corrigendum to "High risks of all-cause and cardiovascular deaths in apparently healthy middle-aged people with preserved glomerular filtration rate and albuminuria: A prospective cohort study" [Int J Cardiol 170 (2013) 167-172]. <i>International Journal of Cardiology</i> , 2014, 175, 595.	0.8	0
87	Bradycardia is associated with future cardiovascular diseases and death in men from the general population. <i>Atherosclerosis</i> , 2014, 236, 116-120.	0.4	9
88	Cardiovascular Risk Stratification With Plasma B-Type Natriuretic Peptide Levels in a Community-Based Hypertensive Cohort. <i>American Journal of Cardiology</i> , 2014, 113, 682-686.	0.7	5
89	Usefulness of risk grading system using albuminuria for predicting cardiovascular events and all-cause death in chronic kidney disease: A population-based prospective cohort study in Japan. <i>International Journal of Cardiology</i> , 2014, 175, 576-577.	0.8	0
90	High risks of all-cause and cardiovascular deaths in apparently healthy middle-aged people with preserved glomerular filtration rate and albuminuria: A prospective cohort study. <i>International Journal of Cardiology</i> , 2013, 170, 167-172.	0.8	16



#	ARTICLE	IF	CITATIONS
91	Concordance of CKD stages in estimation by the CKD-EPI equation and estimation by the MDRD equation in the Japanese general population: The Iwate KENCO Study. <i>International Journal of Cardiology</i> , 2013, 165, 377-379.	0.8	6
92	Incidence Rate of Cerebrovascular Diseases in Northern Japan Determined from the Iwate Stroke Registry with an Inventory Survey System. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013, 22, e317-e322.	0.7	19
93	Predictive Value of Lipoprotein Indices for Residual Risk of Acute Myocardial Infarction and Sudden Death in Men With Low-Density Lipoprotein Cholesterol Levels <math>\leq 120</math> mg/dl. <i>American Journal of Cardiology</i> , 2013, 112, 1063-1068.	0.7	13
94	Influence of the Great East Japan Earthquake and Tsunami 2011 on Occurrence of Cerebrovascular Diseases in Iwate, Japan. <i>Stroke</i> , 2013, 44, 1518-1524.	1.0	58
95	Associations of marital status with mortality from all causes and mortality from cardiovascular disease in Japanese haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 1013-1020.	0.4	15
96	Conflicting effect of alcohol on cardiovascular risk: a clue to understand the different etiologies of coronary artery disease, stroke and peripheral artery disease. <i>Hypertension Research</i> , 2013, 36, 16-18.	1.5	5
97	Comparison of Predictability of Future Cardiovascular Events Between Chronic Kidney Disease (CKD) Stage Based on CKD Epidemiology Collaboration Equation and That Based on Modification of Diet in Renal Disease Equation in the Japanese General Population. <i>Circulation Journal</i> , 2013, 77, 1315-1325.	0.7	25
98	Factors Related to Tooth Loss Among Community-Dwelling Middle-aged and Elderly Japanese Men. <i>Journal of Epidemiology</i> , 2013, 23, 301-306.	1.1	42
99	Plasma B-type Natriuretic Peptide as a Predictor of Cardiovascular Events in Subjects with Atrial Fibrillation: A Community-Based Study. <i>PLoS ONE</i> , 2013, 8, e81243.	1.1	16
100	Influence of mild-to-moderate alcohol consumption on cardiovascular diseases in men from the general population. <i>Atherosclerosis</i> , 2012, 224, 222-227.	0.4	11
101	Poor self-rated health is significantly associated with elevated C-reactive protein levels in women, but not in men, in the Japanese general population. <i>Journal of Psychosomatic Research</i> , 2012, 73, 225-231.	1.2	28
102	Plasma B-Type Natriuretic Peptide Is Useful for Cardiovascular Risk Assessment in Community-Based Diabetes Subjects. <i>International Heart Journal</i> , 2012, 53, 176-181.	0.5	22
103	Corrigendum to "Gender-specific risk stratification with plasma B-type natriuretic peptide for future onset of congestive heart failure and mortality in the Japanese general population" <i>Int J Cardiol</i> 143 (2010) 124-129. <i>International Journal of Cardiology</i> , 2011, 146, 485.	0.8	0
104	Serum ionic fluoride concentrations are significantly decreased after treatment with alendronate in patients with osteoporosis. <i>Clinica Chimica Acta</i> , 2011, 412, 2146-2149.	0.5	6
105	Serum Low-Density Lipoprotein to High-Density Lipoprotein Ratio as a Predictor of Future Acute Myocardial Infarction Among Men in a 2.7-Year Cohort Study of a Japanese Northern Rural Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011, 18, 89-98.	0.9	63
106	Serum C-Reactive Protein Levels and Death and Cardiovascular Events in Mild to Moderate Chronic Kidney Disease. <i>International Heart Journal</i> , 2011, 52, 180-184.	0.5	7
107	Seropositivity for Anti-HCV Core Antigen is Independently Associated With Increased All-Cause, Cardiovascular, and Liver Disease-Related Mortality in Hemodialysis Patients. <i>Journal of Epidemiology</i> , 2011, 21, 491-499.	1.1	24
108	Survey of early preterm singleton live births in Iwate Prefecture as a provincial model of the medical situation in Japan. <i>Journal of Obstetrics and Gynaecology Research</i> , 2011, 37, 867-873.	0.6	0

#	ARTICLE	IF	CITATIONS
109	Sex-Specific Threshold Levels of Plasma B-Type Natriuretic Peptide for Prediction of Cardiovascular Event Risk in a Japanese Population Initially Free of Cardiovascular Disease. <i>American Journal of Cardiology</i> , 2011, 108, 1564-1569.	0.7	12
110	Serum selenium levels are inversely associated with death risk among hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3331-3338.	0.4	56
111	Serum Selenium Levels in Hemodialysis Patients Are Significantly Lower than Those in Healthy Controls. <i>Blood Purification</i> , 2011, 32, 43-47.	0.9	24
112	Plasma B-Type Natriuretic Peptide Level and Cardiovascular Events in Chronic Kidney Disease in a Community-Based Population. <i>Circulation Journal</i> , 2010, 74, 792-797.	0.7	24
113	Standardized Prevalence Ratios for Chronic Hepatitis C Virus Infection Among Adult Japanese Hemodialysis Patients. <i>Journal of Epidemiology</i> , 2010, 20, 30-39.	1.1	18
114	Prehypertension Subtype With Elevated C-Reactive Protein: Risk of Ischemic Stroke in a General Japanese Population. <i>American Journal of Hypertension</i> , 2010, 23, 1108-1113.	1.0	24
115	Gender-specific risk stratification with plasma B-type natriuretic peptide for future onset of congestive heart failure and mortality in the Japanese general population. <i>International Journal of Cardiology</i> , 2010, 143, 124-129.	0.8	11
116	Serum ionic fluoride concentrations are related to renal function and menopause status but not to age in a Japanese general population. <i>Clinica Chimica Acta</i> , 2010, 411, 263-266.	0.5	16
117	Comparison of low-density lipoprotein cholesterol concentrations measured by a direct homogeneous assay and by the Friedewald formula in a large community population. <i>Clinica Chimica Acta</i> , 2010, 411, 1774-1780.	0.5	42
118	Associations of ikigai as a positive psychological factor with all-cause mortality and cause-specific mortality among middle-aged and elderly Japanese people: Findings from the Japan Collaborative Cohort Study. <i>Journal of Psychosomatic Research</i> , 2009, 67, 67-75.	1.2	89
119	Cardiovascular risk factors in the Japanese northeastern rural population. <i>International Journal of Cardiology</i> , 2009, 137, 226-235.	0.8	40
120	Serum C-reactive protein levels can be used to predict future ischemic stroke and mortality in Japanese men from the general population. <i>Atherosclerosis</i> , 2009, 204, 234-238.	0.4	65
121	Predictive value of plasma B-type natriuretic peptide for ischemic stroke: A community-based longitudinal study. <i>Atherosclerosis</i> , 2009, 207, 298-303.	0.4	36
122	Association of hip fracture incidence and intake of calcium, magnesium, vitamin D, and vitamin K. <i>European Journal of Epidemiology</i> , 2008, 23, 219-225.	2.5	76
123	Dietary intake of n-3 polyunsaturated fatty acids is inversely associated with CRP levels, especially among male smokers. <i>Atherosclerosis</i> , 2008, 201, 184-191.	0.4	29
124	Mortality Risk Attributable to Atrial Fibrillation in Middle-Aged and Elderly People in the Japanese General Population Nineteen-Year Follow-up in NIPPON DATA80. <i>Circulation Journal</i> , 2007, 71, 814-819.	0.7	61