
Frédéric Courbon

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3007676/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Efficacy and safety of 177Lu‑DOTATATE in patients with advanced pancreatic neuroendocrine tumours: data from the NETTER-R international, retrospective study. European Journal of Nuclear Medicine and Molecular Imaging, 2022, 49, 3529-3537.	3.3	6
2	Gated ¹⁸ F-FDG PET/CT of the Lung Using a Respiratory Spirometric Gating Device: A Feasibility Study. Journal of Nuclear Medicine Technology, 2019, 47, 227-232.	0.4	4
3	Incidental Finding of Intrathyroid Metastases of Prostatic Cancer on 18F-Choline PET/CT. Clinical Nuclear Medicine, 2019, 44, e101-e103.	0.7	3
4	Evaluation of 2 diuretic 18fluorine-fluorodeoxyglucose positron emission tomography/computed tomography imaging protocols for intrapelvic cancer. Quarterly Journal of Nuclear Medicine and Molecular Imaging, 2019, 63, 284-291.	0.4	4
5	Controversies and consensus in the innovation access for cancer therapy in the European countries: on the subject of metastatic prostate cancer. Annals of Oncology, 2017, 28, 421-426.	0.6	4
6	Phase II Study of a Radiotherapy Total Dose Increase in Hypoxic Lesions Identified by ¹⁸ F-Misonidazole PET/CT in Patients with Non–Small Cell Lung Carcinoma (RTEP5 Study). Journal of Nuclear Medicine, 2017, 58, 1045-1053.	2.8	70
7	Clinical outcomes 1 year after empiric 1311 therapy for hyperthyroid disorders. Nuclear Medicine Communications, 2017, 38, 756-763.	0.5	18
8	¹⁸ Fâ€Fluorineâ€18â€ <scp>l</scp> â€dihydroxyphenylalanine (¹⁸ Fâ€DOPA) positive iso peritoneal carcinomatosis from a MENIIâ€related medullary thyroid carcinoma. About an atypical metastatic site and utility of ¹⁸ Fâ€FDOPA. Clinical Case Reports (discontinued), 2015, 3, 81-83.	lated 0.2	1
9	Nonsurgical giant cell tumour of the tendon sheath or of the diffuse type: Are MRI or 18F-FDG PET/CT able to provide an accurate prediction of long-term outcome?. European Journal of Nuclear Medicine and Molecular Imaging, 2015, 42, 397-408.	3.3	22
10	Oncocytic Adenoma of Thyroid Incidentally Detected by 18F-Fluorocholine PET/CT. Journal of Nuclear Medicine Technology, 2015, 43, 133-134.	0.4	18
11	Comparison of an alternative and existing binning methods to reduce the acquisition duration of 4D PET/CT. Medical Physics, 2014, 41, 112503.	1.6	3
12	Evaluating digestive neuroendocrine tumor progression and therapeutic responses in the era of targeted therapies: state of the art. Endocrine-Related Cancer, 2014, 21, R105-R120.	1.6	59
13	Phase II study of temozolomide in combination with topotecan (TOTEM) in relapsed or refractory neuroblastoma: A European Innovative Therapies for Children with Cancer-SIOP-European Neuroblastoma study. European Journal of Cancer, 2014, 50, 170-177.	1.3	47
14	A new respiratory gating device to improve 4D PET/CT. Medical Physics, 2013, 40, 032501.	1.6	13
15	Interobserver Agreement of Qualitative Analysis and Tumor Delineation of ¹⁸ F-Fluoromisonidazole and 3â€2-Deoxy-3â€2- ¹⁸ F-Fluorothymidine PET Images in Lung Cancer. Journal of Nuclear Medicine, 2013, 54, 1543-1550.	2.8	44
16	Breast Imaging: Understanding How Accuracy is Measured When Lesion are the Unit of Analysis. Breast Journal, 2013, 19, 225-226.	0.4	0
17	The retrospective binning method improves the consistency of phase binning in respiratory-gated PET/CT. Physics in Medicine and Biology, 2012, 57, 7829-7841.	1.6	22
18	Routine FDG PETâ€CT in patients with a highâ€risk localized melanoma has a high predictive positive value for nodal disease and high negative predictive value for the presence of distant metastases. Journal of the European Academy of Dermatology and Venereology, 2012, 26, 1431-1435.	1.3	15

Frédéric Courbon

#	Article	IF	CITATIONS
19	Functional testicular evaluation using PET/CT with 18F-fluorodeoxyglucose. European Journal of Nuclear Medicine and Molecular Imaging, 2012, 39, 129-137.	3.3	14
20	Fluorodeoxyglucose positron emission tomography fails to detect distant metastases at initial staging of melanoma patients with metastatic involvement of sentinel lymph node. British Journal of Dermatology, 2011, 164, 1235-1240.	1.4	31
21	Evaluation of a Trainer Phantom in the Learning Phase of Sentinel Lymph Node Identification in Breast Cancer. World Journal of Surgery, 2011, 35, 995-1001.	0.8	4
22	Coregistration of Prechemotherapy PET-CT for Planning Pediatric Hodgkin's Disease Radiotherapy Significantly Diminishes Interobserver Variability of Clinical Target Volume Definition. International Journal of Radiation Oncology Biology Physics, 2011, 80, 793-799.	0.4	31
23	Importance of cholesterol and oxysterols metabolism in the pharmacology of tamoxifen and other AEBS ligands. Chemistry and Physics of Lipids, 2011, 164, 432-437.	1.5	51
24	Quality controls and delineation protocol of PET/CT gated acquisition in function of the movement amplitude, size of spheres and signal over background ratio. Proceedings of SPIE, 2011, , .	0.8	4
25	Development of a new radioligand for cholecystokinin receptor subtype 2 scintigraphy: From molecular modeling to in vivo evaluation. Bioorganic and Medicinal Chemistry, 2010, 18, 5400-5412.	1.4	12
26	Trastuzumab induced in vivo tissue remodelling associated in vitro with inhibition of the active forms of AKT and PTEN and RhoB induction in an ovarian carcinoma model. British Journal of Cancer, 2010, 103, 61-72.	2.9	12
27	Preclinical and Clinical Evidence that Deoxy-2-[18F]fluoro- <scp>D</scp> -glucose Positron Emission Tomography with Computed Tomography Is a Reliable Tool for the Detection of Early Molecular Responses to Erlotinib in Head and Neck Cancer. Clinical Cancer Research, 2010, 16, 4434-4445.	3.2	27
28	Short―and longâ€ŧerm somatostatin analogue treatment in patients with hypoglycaemia related to endogenous hyperinsulinism. Clinical Endocrinology, 2008, 68, 904-911.	1.2	50
29	Adenoid cystic carcinoma of the lung: Interest of 18FDG PET/CT in the management of an atypical presentation. Lung Cancer, 2008, 59, 133-136.	0.9	18
30	Indium-111–Pentetreotide Scintigraphy and Somatostatin Receptor Subtype 2 Expression: New Prognostic Factors for Malignant Well-Differentiated Endocrine Tumors. Journal of Clinical Oncology, 2008, 26, 963-970.	0.8	99
31	Pilot Study of Neoadjuvant Treatment with Erlotinib in Nonmetastatic Head and Neck Squamous Cell Carcinoma. Clinical Cancer Research, 2007, 13, 7086-7092.	3.2	68
32	Preparation and Use of 1311 Magic Gel as a Dosimeter for Targeted Radionuclide Therapy. Cancer Biotherapy and Radiopharmaceuticals, 2006, 21, 427-436.	0.7	10
33	Defective Efficacy of Retinoic Acid Treatment in Patients with Metastatic Thyroid Carcinoma. Thyroid, 2006, 16, 1025-1031.	2.4	44
34	V-Q SPECT in Prone Position Is a Suitable Way to Improve Irradiation of Inoperable Lung Cancer. Clinical Nuclear Medicine, 2006, 31, 284-288.	0.7	0
35	Octreotide in insulinoma patients: efficacy on hypoglycemia, relationships with Octreoscan scintigraphy and immunostaining with anti-sst2A and anti-sst5 antibodies. European Journal of Endocrinology, 2005, 152, 757-767.	1.9	160
36	The incorporation of SPECT functional lung imaging into inverse radiotherapy planning for non-small cell lung cancer. Radiotherapy and Oncology, 2005, 77, 271-277.	0.3	100

#	Article	IF	CITATIONS
37	Cardiac MIBG scintigraphy is a sensitive tool for detecting cardiac sympathetic denervation in Parkinson's disease. Movement Disorders, 2003, 18, 890-897.	2.2	202
38	Summary of the Standards, Options and Recommendations for the use of positron emission tomography with 2-[18F]fluoro-2-deoxy-D-glucose (FDP-PET scanning) in oncology (2002). British Journal of Cancer, 2003, 89, S84-S91.	2.9	36
39	Granulocyte-macrophage colony-stimulating factor alone or with dacarbazine in metastatic melanoma: a randomized phase II trial. British Journal of Cancer, 2001, 85, 1467-1471.	2.9	14
40	A pharmacokinetic interpretation of increasing concentrations of DEHP in haemodialysed patients. Medical Engineering and Physics, 2000, 22, 157-165.	0.8	15