

# Erwin London

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/3015296/erwin-london-publications-by-citations.pdf>  
**Version:** 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140 papers	12,243 citations	54 h-index	110 g-index
149 ext. papers	13,088 ext. citations	4.3 avg, IF	6.69 L-index

#	Paper	IF	Citations
140	Structure and function of sphingolipid- and cholesterol-rich membrane rafts. <i>Journal of Biological Chemistry</i> , <b>2000</b> , 275, 17221-4	5.4	1868
139	On the origin of sphingolipid/cholesterol-rich detergent-insoluble cell membranes: physiological concentrations of cholesterol and sphingolipid induce formation of a detergent-insoluble, liquid-ordered lipid phase in model membranes. <i>Biochemistry</i> , <b>1997</b> , 36, 10944-53	3.2	607
138	Parallax method for direct measurement of membrane penetration depth utilizing fluorescence quenching by spin-labeled phospholipids. <i>Biochemistry</i> , <b>1987</b> , 26, 39-45	3.2	599
137	Insolubility of lipids in triton X-100: physical origin and relationship to sphingolipid/cholesterol membrane domains (rafts). <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2000</b> , 1508, 182-95	3.8	520
136	Structure of detergent-resistant membrane domains: does phase separation occur in biological membranes?. <i>Biochemical and Biophysical Research Communications</i> , <b>1997</b> , 240, 1-7	3.4	456
135	The effect of sterol structure on membrane lipid domains reveals how cholesterol can induce lipid domain formation. <i>Biochemistry</i> , <b>2000</b> , 39, 843-9	3.2	439
134	Effect of the structure of natural sterols and sphingolipids on the formation of ordered sphingolipid/sterol domains (rafts). Comparison of cholesterol to plant, fungal, and disease-associated sterols and comparison of sphingomyelin, cerebroside, and ceramide. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 33540-6	5.4	435
133	Fluorimetric determination of critical micelle concentration avoiding interference from detergent charge. <i>Analytical Biochemistry</i> , <b>1984</b> , 139, 408-12	3.1	353
132	Cholesterol and sphingolipid enhance the Triton X-100 insolubility of glycosylphosphatidylinositol-anchored proteins by promoting the formation of detergent-insoluble ordered membrane domains. <i>Journal of Biological Chemistry</i> , <b>1998</b> , 273, 1150-7	5.4	346
131	Ceramide selectively displaces cholesterol from ordered lipid domains (rafts): implications for lipid raft structure and function. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 9997-10004	5.4	325
130	Location of diphenylhexatriene (DPH) and its derivatives within membranes: comparison of different fluorescence quenching analyses of membrane depth. <i>Biochemistry</i> , <b>1998</b> , 37, 8180-90	3.2	294
129	Insights into lipid raft structure and formation from experiments in model membranes. <i>Current Opinion in Structural Biology</i> , <b>2002</b> , 12, 480-6	8.1	231
128	How principles of domain formation in model membranes may explain ambiguities concerning lipid raft formation in cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , <b>2005</b> , 1746, 203-20	4.9	197
127	Extension of the parallax analysis of membrane penetration depth to the polar region of model membranes: use of fluorescence quenching by a spin-label attached to the phospholipid polar headgroup. <i>Biochemistry</i> , <b>1993</b> , 32, 10826-31	3.2	192
126	Transmembrane orientation of hydrophobic alpha-helices is regulated both by the relationship of helix length to bilayer thickness and by the cholesterol concentration. <i>Biochemistry</i> , <b>1997</b> , 36, 10213-20	3.2	191
125	Effect of pH on the conformation of diphtheria toxin and its implications for membrane penetration. <i>Biochemistry</i> , <b>1985</b> , 24, 5458-64	3.2	182
124	Preparation and properties of asymmetric vesicles that mimic cell membranes: effect upon lipid raft formation and transmembrane helix orientation. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 6079-92	5.4	142

123	Relationship between sterol/steroid structure and participation in ordered lipid domains (lipid rafts): implications for lipid raft structure and function. <i>Biochemistry</i> , <b>2004</b> , 43, 1010-8	3.2	137
122	Diphtheria toxin: membrane interaction and membrane translocation. <i>BBA - Biomembranes</i> , <b>1992</b> , 1113, 25-51		130
121	Control of the transmembrane orientation and interhelical interactions within membranes by hydrophobic helix length. <i>Biochemistry</i> , <b>1999</b> , 38, 5905-12	3.2	122
120	Fluorescence quenching in model membranes. 1. Characterization of quenching caused by a spin-labeled phospholipid. <i>Biochemistry</i> , <b>1981</b> , 20, 1932-8	3.2	120
119	Anchoring of tryptophan and tyrosine analogs at the hydrocarbon-polar boundary in model membrane vesicles: parallax analysis of fluorescence quenching induced by nitroxide-labeled phospholipids. <i>Biochemistry</i> , <b>1995</b> , 34, 15475-9	3.2	118
118	Cholesterol precursors stabilize ordinary and ceramide-rich ordered lipid domains (lipid rafts) to different degrees. Implications for the Bloch hypothesis and sterol biosynthesis disorders. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 21903-21913	5.4	116
117	Interaction of diphtheria toxin T domain with molten globule-like proteins and its implications for translocation. <i>Science</i> , <b>1999</b> , 284, 955-7	33.3	114
116	Calibration of the parallax fluorescence quenching method for determination of membrane penetration depth: refinement and comparison of quenching by spin-labeled and brominated lipids. <i>Biochemistry</i> , <b>1992</b> , 31, 5312-22	3.2	114
115	Effect of the structure of lipids favoring disordered domain formation on the stability of cholesterol-containing ordered domains (lipid rafts): identification of multiple raft-stabilization mechanisms. <i>Biophysical Journal</i> , <b>2007</b> , 93, 4307-18	2.9	105
114	Palmitoylation and intracellular domain interactions both contribute to raft targeting of linker for activation of T cells. <i>Journal of Biological Chemistry</i> , <b>2005</b> , 280, 18931-42	5.4	102
113	Measurement of lipid nanodomain (raft) formation and size in sphingomyelin/POPC/cholesterol vesicles shows TX-100 and transmembrane helices increase domain size by coalescing preexisting nanodomains but do not induce domain formation. <i>Biophysical Journal</i> , <b>2011</b> , 101, 2417-25	2.9	101
112	Asymmetric GUVs prepared by MCD-mediated lipid exchange: an FCS study. <i>Biophysical Journal</i> , <b>2011</b> , 100, L1-3	2.9	98
111	How interaction of perfringolysin O with membranes is controlled by sterol structure, lipid structure, and physiological low pH: insights into the origin of perfringolysin O-lipid raft interaction. <i>Journal of Biological Chemistry</i> , <b>2008</b> , 283, 4632-42	5.4	93
110	Determination of the location of fluorescent probes attached to fatty acids using parallax analysis of fluorescence quenching: effect of carboxyl ionization state and environment on depth. <i>Biochemistry</i> , <b>1992</b> , 31, 5322-7	3.2	87
109	Using a novel dual fluorescence quenching assay for measurement of tryptophan depth within lipid bilayers to determine hydrophobic alpha-helix locations within membranes. <i>Biochemistry</i> , <b>2003</b> , 42, 3265-74	3.2	86
108	Acyl chain length and saturation modulate interleaflet coupling in asymmetric bilayers: effects on dynamics and structural order. <i>Biophysical Journal</i> , <b>2012</b> , 103, 2311-9	2.9	85
107	Proving lipid rafts exist: membrane domains in the prokaryote <i>Borrelia burgdorferi</i> have the same properties as eukaryotic lipid rafts. <i>PLoS Pathogens</i> , <b>2013</b> , 9, e1003353	7.6	84
106	An amino acid "transmembrane tendency" scale that approaches the theoretical limit to accuracy for prediction of transmembrane helices: relationship to biological hydrophobicity. <i>Protein Science</i> , <b>2006</b> , 15, 1987-2001	6.3	84

105	Cholesterol lipids of <i>Borrelia burgdorferi</i> form lipid rafts and are required for the bactericidal activity of a complement-independent antibody. <i>Cell Host and Microbe</i> , <b>2010</b> , 8, 331-42	23.4	83
104	Cumulative effects of amino acid substitutions and hydrophobic mismatch upon the transmembrane stability and conformation of hydrophobic alpha-helices. <i>Biochemistry</i> , <b>2003</b> , 42, 3275-83 <sup>2</sup>	3.2	83
103	Lipid exchange between <i>Borrelia burgdorferi</i> and host cells. <i>PLoS Pathogens</i> , <b>2013</b> , 9, e1003109	7.6	81
102	The location of fluorescence probes with charged groups in model membranes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>1998</b> , 1374, 63-76	3.8	79
101	Subnanometer Structure of an Asymmetric Model Membrane: Interleaflet Coupling Influences Domain Properties. <i>Langmuir</i> , <b>2016</b> , 32, 5195-200	4	79
100	Preparation of asymmetric phospholipid vesicles for use as cell membrane models. <i>Nature Protocols</i> , <b>2018</b> , 13, 2086-2101	18.8	79
99	Preparation and properties of asymmetric large unilamellar vesicles: interleaflet coupling in asymmetric vesicles is dependent on temperature but not curvature. <i>Biophysical Journal</i> , <b>2011</b> , 100, 2671-8	7.8	76
98	Exclusion of a transmembrane-type peptide from ordered-lipid domains (rafts) detected by fluorescence quenching: extension of quenching analysis to account for the effects of domain size and domain boundaries. <i>Biochemistry</i> , <b>2003</b> , 42, 12376-90	3.2	76
97	Effect of ceramide N-acyl chain and polar headgroup structure on the properties of ordered lipid domains (lipid rafts). <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2007</b> , 1768, 2205-12	3.8	73
96	Islet Amyloid Polypeptide Membrane Interactions: Effects of Membrane Composition. <i>Biochemistry</i> , <b>2017</b> , 56, 376-390	3.2	72
95	Ordered raft domains induced by outer leaflet sphingomyelin in cholesterol-rich asymmetric vesicles. <i>Biophysical Journal</i> , <b>2015</b> , 108, 2212-22	2.9	68
94	H NMR Shows Slow Phospholipid Flip-Flop in Gel and Fluid Bilayers. <i>Langmuir</i> , <b>2017</b> , 33, 3731-3741	4	65
93	Preparation of artificial plasma membrane mimicking vesicles with lipid asymmetry. <i>PLoS ONE</i> , <b>2014</b> , 9, e87903	3.7	60
92	Membrane topography of the hydrophobic anchor sequence of poliovirus 3A and 3AB proteins and the functional effect of 3A/3AB membrane association upon RNA replication. <i>Biochemistry</i> , <b>2007</b> , 46, 5185-99	3.2	59
91	Identification of shallow and deep membrane-penetrating forms of diphtheria toxin T domain that are regulated by protein concentration and bilayer width. <i>Journal of Biological Chemistry</i> , <b>1997</b> , 272, 25091-8	5.4	57
90	Position and ionization state of Asp in the core of membrane-inserted alpha helices control both the equilibrium between transmembrane and nontransmembrane helix topography and transmembrane helix positioning. <i>Biochemistry</i> , <b>2004</b> , 43, 8794-806	3.2	57
89	Fluorescence quenching in model membranes An analysis of the local phospholipid environments of diphenylhexatriene and gramicidin A?. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>1981</b> , 649, 89-97 <sup>3</sup>	3.8	57
88	Effect of cyclodextrin and membrane lipid structure upon cyclodextrin-lipid interaction. <i>Langmuir</i> , <b>2013</b> , 29, 14631-8	4	55

87	The Effect of Membrane Lipid Composition on the Formation of Lipid Ultrananodomains. <i>Biophysical Journal</i> , <b>2015</b> , 109, 1630-8	2.9	54
86	Altering hydrophobic sequence lengths shows that hydrophobic mismatch controls affinity for ordered lipid domains (rafts) in the multitransmembrane strand protein perfringolysin O. <i>Journal of Biological Chemistry</i> , <b>2013</b> , 288, 1340-52	5.4	54
85	How bacterial protein toxins enter cells; the role of partial unfolding in membrane translocation. <i>Molecular Microbiology</i> , <b>1992</b> , 6, 3277-82	4.1	53
84	Groups with polar characteristics can locate at both shallow and deep locations in membranes: the behavior of dansyl and related probes. <i>Biochemistry</i> , <b>1998</b> , 37, 4603-11	3.2	52
83	Investigation of membrane structure using fluorescence quenching by spin-labels. A review of recent studies. <i>Molecular and Cellular Biochemistry</i> , <b>1982</b> , 45, 181-8	4.2	51
82	Measuring the depth of amino acid residues in membrane-inserted peptides by fluorescence quenching. <i>Current Topics in Membranes</i> , <b>2002</b> , 52, 89-115	2.2	50
81	Notch-modifying xylosyltransferase structures support an SNI-like retaining mechanism. <i>Nature Chemical Biology</i> , <b>2015</b> , 11, 847-54	11.7	49
80	Analyzing topography of membrane-inserted diphtheria toxin T domain using BODIPY-streptavidin: at low pH, helices 8 and 9 form a transmembrane hairpin but helices 5-7 form stable nonclassical inserted segments on the cis side of the bilayer. <i>Biochemistry</i> , <b>2004</b> , 43, 9127-39	3.2	47
79	Identifying transmembrane states and defining the membrane insertion boundaries of hydrophobic helices in membrane-inserted diphtheria toxin T domain. <i>Journal of Biological Chemistry</i> , <b>1998</b> , 273, 22950-6	5.4	47
78	Efficient replacement of plasma membrane outer leaflet phospholipids and sphingolipids in cells with exogenous lipids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, 14025-14030	11.5	45
77	Control of the depth of molecules within membranes by polar groups: determination of the location of anthracene-labeled probes in model membranes by parallax analysis of nitroxide-labeled phospholipid induced fluorescence quenching. <i>Biochemistry</i> , <b>1995</b> , 34, 11460-6	3.2	41
76	Effect of sequence hydrophobicity and bilayer width upon the minimum length required for the formation of transmembrane helices in membranes. <i>Journal of Molecular Biology</i> , <b>2007</b> , 374, 671-87	6.5	40
75	The effects of polar and/or ionizable residues in the core and flanking regions of hydrophobic helices on transmembrane conformation and oligomerization. <i>Biochemistry</i> , <b>2000</b> , 39, 9632-40	3.2	40
74	Membrane topography of the T domain of diphtheria toxin probed with single tryptophan mutants. <i>Biochemistry</i> , <b>1998</b> , 37, 17915-22	3.2	40
73	Raft-like membrane domains in pathogenic microorganisms. <i>Current Topics in Membranes</i> , <b>2015</b> , 75, 233-68	6.8	35
72	Perfringolysin O association with ordered lipid domains: implications for transmembrane protein raft affinity. <i>Biophysical Journal</i> , <b>2010</b> , 99, 3255-63	2.9	35
71	The dependence of lipid asymmetry upon phosphatidylcholine acyl chain structure. <i>Journal of Lipid Research</i> , <b>2013</b> , 54, 223-31	6.3	33
70	The control of transmembrane helix transverse position in membranes by hydrophilic residues. <i>Journal of Molecular Biology</i> , <b>2007</b> , 374, 1251-69	6.5	33

69	Behavior of diphtheria toxin T domain containing substitutions that block normal membrane insertion at Pro345 and Leu307: control of deep membrane insertion and coupling between deep insertion of hydrophobic subdomains. <i>Biochemistry</i> , <b>2005</b> , 44, 4488-98	3.2	33
68	Role of predicted transmembrane domains for type III translocation, pore formation, and signaling by the <i>Yersinia pseudotuberculosis</i> YopB protein. <i>Infection and Immunity</i> , <b>2005</b> , 73, 2433-43	3.7	32
67	Simple procedure for reversed-phase high-performance liquid chromatographic purification of long hydrophobic peptides that form transmembrane helices. <i>Analytical Biochemistry</i> , <b>1997</b> , 251, 113-6	3.1	31
66	The effect of sterol structure upon clathrin-mediated and clathrin-independent endocytosis. <i>Journal of Cell Science</i> , <b>2017</b> , 130, 2682-2695	5.3	30
65	Scanning the membrane-bound conformation of helix 1 in the colicin E1 channel domain by site-directed fluorescence labeling. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 885-95	5.4	30
64	Cholesterol lipids and cholesterol-containing lipid rafts in bacteria. <i>Chemistry and Physics of Lipids</i> , <b>2016</b> , 199, 11-16	3.7	29
63	Changes in glucosylceramide structure affect virulence and membrane biophysical properties of <i>Cryptococcus neoformans</i> . <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2017</b> , 1859, 2224-2233	3.8	29
62	The effect of interactions involving ionizable residues flanking membrane-inserted hydrophobic helices upon helix-helix interaction. <i>Biochemistry</i> , <b>2003</b> , 42, 10833-42	3.2	28
61	Lipid rafts can form in the inner and outer membranes of <i>Borrelia burgdorferi</i> and have different properties and associated proteins. <i>Molecular Microbiology</i> , <b>2018</b> , 108, 63-76	4.1	27
60	The dependence of lipid asymmetry upon polar headgroup structure. <i>Journal of Lipid Research</i> , <b>2013</b> , 54, 3385-93	6.3	27
59	Membrane Structure-Function Insights from Asymmetric Lipid Vesicles. <i>Accounts of Chemical Research</i> , <b>2019</b> , 52, 2382-2391	24.3	26
58	The influence of natural lipid asymmetry upon the conformation of a membrane-inserted protein (perfringolysin O). <i>Journal of Biological Chemistry</i> , <b>2014</b> , 289, 5467-78	5.4	26
57	Selective association of outer surface lipoproteins with the lipid rafts of <i>Borrelia burgdorferi</i> . <i>MBio</i> , <b>2014</b> , 5, e00899-14	7.8	26
56	Lipid Structure and Composition Control Consequences of Interleaflet Coupling in Asymmetric Vesicles. <i>Biophysical Journal</i> , <b>2018</b> , 115, 664-678	2.9	25
55	Effect of lipid composition on the topography of membrane-associated hydrophobic helices: stabilization of transmembrane topography by anionic lipids. <i>Journal of Molecular Biology</i> , <b>2008</b> , 379, 704-18	6.5	25
54	Topography of helices 5-7 in membrane-inserted diphtheria toxin T domain: identification and insertion boundaries of two hydrophobic sequences that do not form a stable transmembrane hairpin. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 16517-27	5.4	25
53	Sphingolipids and membrane domains: recent advances. <i>Handbook of Experimental Pharmacology</i> , <b>2013</b> , 33-55	3.2	23
52	Interaction of the membrane-inserted diphtheria toxin T domain with peptides and its possible implications for chaperone-like T domain behavior. <i>Biochemistry</i> , <b>2002</b> , 41, 3243-53	3.2	23



51	Transmembrane vs. non-transmembrane hydrophobic helix topography in model and natural membranes. <i>Current Opinion in Structural Biology</i> , <b>2009</b> , 19, 464-72	8.1	22
50	Topography of diphtheria toxin A chain inserted into lipid vesicles. <i>Biochemistry</i> , <b>2005</b> , 44, 2183-96	3.2	21
49	Effect of sterol structure on ordered membrane domain (raft) stability in symmetric and asymmetric vesicles. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2019</b> , 1861, 1112-1122	3.8	20
48	A novel leaflet-selective fluorescence labeling technique reveals differences between inner and outer leaflets at high bilayer curvature. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2012</b> , 1818, 1284-90	3.8	20
47	Topography of the hydrophilic helices of membrane-inserted diphtheria toxin T domain: TH1-TH3 as a hydrophilic tether. <i>Biochemistry</i> , <b>2006</b> , 45, 8124-34	3.2	20
46	The membrane topography of the diphtheria toxin T domain linked to the a chain reveals a transient transmembrane hairpin and potential translocation mechanisms. <i>Biochemistry</i> , <b>2009</b> , 48, 10446-56	3.2	18
45	Effect of lipid composition and amino acid sequence upon transmembrane peptide-accelerated lipid transleaflet diffusion (Flip-Flop). <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2016</b> , 1858, 1812-20	3.8	18
44	The effect of hydrophilic substitutions and anionic lipids upon the transverse positioning of the transmembrane helix of the ErbB2 (neu) protein incorporated into model membrane vesicles. <i>Journal of Molecular Biology</i> , <b>2010</b> , 396, 209-20	6.5	17
43	Use of Trp mutations to evaluate the conformational behavior and membrane insertion of A and B chains in whole diphtheria toxin. <i>Biochemistry</i> , <b>1997</b> , 36, 16300-8	3.2	17
42	Nanodomains can persist at physiologic temperature in plasma membrane vesicles and be modulated by altering cell lipids. <i>Journal of Lipid Research</i> , <b>2020</b> , 61, 758-766	6.3	16
41	Behavior of the deeply inserted helices in diphtheria toxin T domain: helices 5, 8, and 9 interact strongly and promote pore formation, while helices 6/7 limit pore formation. <i>Biochemistry</i> , <b>2008</b> , 47, 4565-74	3.2	15
40	Fluorescence quenching assay of sphingolipid/phospholipid phase separation in model membranes. <i>Methods in Enzymology</i> , <b>2000</b> , 312, 272-90	1.7	15
39	Transmembrane protein (perfringolysin o) association with ordered membrane domains (rafts) depends upon the raft-associating properties of protein-bound sterol. <i>Biophysical Journal</i> , <b>2013</b> , 105, 2733-42	2.9	14
38	Sterol structure dependence of insulin receptor and insulin-like growth factor 1 receptor activation. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2019</b> , 1861, 819-826	3.8	13
37	Using Sterol Substitution to Probe the Role of Membrane Domains in Membrane Functions. <i>Lipids</i> , <b>2015</b> , 50, 721-34	1.6	13
36	Induction of Ordered Lipid Raft Domain Formation by Loss of Lipid Asymmetry. <i>Biophysical Journal</i> , <b>2020</b> , 119, 483-492	2.9	10
35	Toward elucidating the membrane topology of helix two of the colicin E1 channel domain. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 32375-84	5.4	10
34	Location of diphenylhexatriene (DPH) and its derivatives within membranes: comparison of different fluorescence quenching analyses of membrane depth. <i>Biochemistry</i> , <b>1999</b> , 38, 2610	3.2	10

33	Ordered Membrane Domain-Forming Properties of the Lipids of <i>Borrelia burgdorferi</i> . <i>Biophysical Journal</i> , <b>2016</b> , 111, 2666-2675	2.9	9
32	Analyzing transmembrane protein and hydrophobic helix topography by dual fluorescence quenching. <i>Methods in Molecular Biology</i> , <b>2013</b> , 974, 279-95	1.4	8
31	Sphingomyelins and ent-Sphingomyelins Form Homophilic Nano-Subdomains within Liquid Ordered Domains. <i>Biophysical Journal</i> , <b>2020</b> , 119, 539-552	2.9	8
30	Sterol Structure Strongly Modulates Membrane-Islet Amyloid Polypeptide Interactions. <i>Biochemistry</i> , <b>2018</b> , 57, 1868-1879	3.2	7
29	Replacing plasma membrane outer leaflet lipids with exogenous lipid without damaging membrane integrity. <i>PLoS ONE</i> , <b>2019</b> , 14, e0223572	3.7	7
28	Strong correlation between statistical transmembrane tendency and experimental hydrophobicity scales for identification of transmembrane helices. <i>Journal of Membrane Biology</i> , <b>2009</b> , 229, 165-8	2.3	7
27	Low pH-induced pore formation by the T domain of botulinum toxin type A is dependent upon NaCl concentration. <i>Journal of Membrane Biology</i> , <b>2010</b> , 236, 191-201	2.3	7
26	Detecting ordered domain formation (lipid rafts) in model membranes using Tempo. <i>Methods in Molecular Biology</i> , <b>2007</b> , 398, 29-40	1.4	7
25	Highly Hydrophilic Segments Attached to Hydrophobic Peptides Translocate Rapidly across Membranes. <i>Langmuir</i> , <b>2016</b> , 32, 10752-10760	4	6
24	Analyzing Transmembrane Protein and Hydrophobic Helix Topography by Dual Fluorescence Quenching. <i>Methods in Molecular Biology</i> , <b>2019</b> , 2003, 351-368	1.4	6
23	Preparation and Drug Entrapment Properties of Asymmetric Liposomes Containing Cationic and Anionic Lipids. <i>Langmuir</i> , <b>2020</b> , 36, 12521-12531	4	6
22	Effects of host cell sterol composition upon internalization of and clustered $\beta$ integrin. <i>Journal of Biological Chemistry</i> , <b>2018</b> , 293, 1466-1479	5.4	6
21	Membrane fusion: A new role for lipid domains?. <i>Nature Chemical Biology</i> , <b>2015</b> , 11, 383-4	11.7	5
20	Mapping peptide thiol accessibility in membranes using a quaternary ammonium isotope-coded mass tag (ICMT). <i>Bioconjugate Chemistry</i> , <b>2013</b> , 24, 1235-47	6.3	5
19	Using model membrane-inserted hydrophobic helices to study the equilibrium between transmembrane and nontransmembrane states. <i>Journal of General Physiology</i> , <b>2007</b> , 130, 229-32	3.4	5
18	Decreasing Transmembrane Segment Length Greatly Decreases Perfringolysin O Pore Size. <i>Journal of Membrane Biology</i> , <b>2015</b> , 248, 517-27	2.3	4
17	The phenyltetraene lysophospholipid analog PTE-ET-18-OMe as a fluorescent anisotropy probe of liquid ordered membrane domains (lipid rafts) and ceramide-rich membrane domains. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2007</b> , 1768, 2213-21	3.8	4
16	<i>Helicobacter pylori</i> lipids can form ordered membrane domains (rafts). <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2019</b> , 1861, 183050	3.8	3



15	Analysis of Lipids and Lipid Rafts in <i>Borrelia</i> . <i>Methods in Molecular Biology</i> , <b>2018</b> , 1690, 69-82	1.4	3
14	Preparation and Physical Properties of Asymmetric Model Membrane Vesicles. <i>Springer Series in Biophysics</i> , <b>2017</b> , 1-27		3
13	Phospholipid exchange shows insulin receptor activity is supported by both the propensity to form wide bilayers and ordered raft domains. <i>Journal of Biological Chemistry</i> , <b>2021</b> , 297, 101010	5.4	3
12	Fluorescence Quenching by a Brominated Detergent: Application to Diphtheria Toxin Structure. <i>Annals of the New York Academy of Sciences</i> , <b>1984</b> , 435, 558-559	6.5	2
11	New Insights into How Cholesterol and Unsaturation Control Lipid Domain Formation. <i>Biophysical Journal</i> , <b>2016</b> , 111, 465-466	2.9	2
10	14. Formation and properties of asymmetric lipid vesicles prepared using cyclodextrin-catalyzed lipid exchange <b>2019</b> , 441-464		1
9	Preparation and utility of asymmetric lipid vesicles for studies of perfringolysin O-lipid interactions. <i>Methods in Enzymology</i> , <b>2021</b> , 649, 253-276	1.7	1
8	Preparation of Asymmetric Vesicles with Trapped CsCl Avoids Osmotic Imbalance, Non-Physiological External Solutions, and Minimizes Leakage. <i>Langmuir</i> , <b>2021</b> , 37, 11611-11617	4	1
7	Using cyclodextrin-induced lipid substitution to study membrane lipid and ordered membrane domain (raft) function in cells. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2022</b> , 1864, 183774	3.8	1
6	Kiss and Run Asymmetric Vesicles to Investigate Coupling. <i>Biophysical Journal</i> , <b>2019</b> , 117, 1009-1011	2.9	0
5	Cholesterol and sphingomyelin are critical for FcγR-mediated phagocytosis of <i>Cryptococcus neoformans</i> by macrophages. <i>Journal of Biological Chemistry</i> , <b>2021</b> , 297, 101411	5.4	0
4	LOSS OF PLASMA MEMBRANE LIPID ASYMMETRY CAN INDUCE ORDERED DOMAIN (RAFT) FORMATION. <i>Journal of Lipid Research</i> , <b>2021</b> , 100155	6.3	0
3	The Fluorescent Dye 1,6-Diphenyl-1,3,5-hexatriene Binds to Amyloid Fibrils Formed by Human Amylin and Provides a New Probe of Amylin Amyloid Kinetics. <i>Biochemistry</i> , <b>2021</b> , 60, 1964-1970	3.2	0
2	Replacing plasma membrane outer leaflet lipids with exogenous lipid without damaging membrane integrity <b>2019</b> , 14, e0223572		
1	Replacing plasma membrane outer leaflet lipids with exogenous lipid without damaging membrane integrity <b>2019</b> , 14, e0223572		