

# Ralph Huckelhoven

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/302540/publications.pdf>

Version: 2024-02-01

129  
papers

10,339  
citations

43973

48  
h-index

35952

97  
g-index

151  
all docs

151  
docs citations

151  
times ranked

10903  
citing authors

#	ARTICLE	IF	CITATIONS
1	The endophytic fungus <i>Piriformospora indica</i> reprograms barley to salt-stress tolerance, disease resistance, and higher yield. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 13386-13391.	3.3	1,153
2	SNARE-protein-mediated disease resistance at the plant cell wall. <i>Nature</i> , 2003, 425, 973-977.	13.7	904
3	Cell Wall-Associated Mechanisms of Disease Resistance and Susceptibility. <i>Annual Review of Phytopathology</i> , 2007, 45, 101-127.	3.5	475
4	The root endophytic fungus <i>Piriformospora indica</i> requires host cell death for proliferation during mutualistic symbiosis with barley. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 18450-18457.	3.3	372
5	Hypersensitive Cell Death and Papilla Formation in Barley Attacked by the Powdery Mildew Fungus Are Associated with Hydrogen Peroxide but Not with Salicylic Acid Accumulation. <i>Plant Physiology</i> , 1999, 119, 1251-1260.	2.3	353
6	Endophyte or parasite – what decides?. <i>Current Opinion in Plant Biology</i> , 2006, 9, 358-363.	3.5	317
7	<i>Bipolaris sorokiniana</i> , a cereal pathogen of global concern: cytological and molecular approaches towards better control. <i>Molecular Plant Pathology</i> , 2002, 3, 185-195.	2.0	310
8	A lectin S-domain receptor kinase mediates lipopolysaccharide sensing in <i>Arabidopsis thaliana</i> . <i>Nature Immunology</i> , 2015, 16, 426-433.	7.0	286
9	Multivesicular bodies participate in a cell wall-associated defence response in barley leaves attacked by the pathogenic powdery mildew fungus. <i>Cellular Microbiology</i> , 2006, 8, 1009-1019.	1.1	274
10	Reactive oxygen intermediates in plant-microbe interactions: Who is who in powdery mildew resistance?. <i>Planta</i> , 2003, 216, 891-902.	1.6	213
11	A Compromised Mlo Pathway Affects the Response of Barley to the Necrotrophic Fungus <i>Bipolaris sorokiniana</i> (Teleomorph: <i>Cochliobolus sativus</i> ) and Its Toxins. <i>Phytopathology</i> , 2001, 91, 127-133.	1.1	184
12	The receptor-like MLO protein and the RAC/ROP family G-protein RACB modulate actin reorganization in barley attacked by the biotrophic powdery mildew fungus <i>Blumeria graminis</i> f.sp. <i>hordei</i> . <i>Plant Journal</i> , 2004, 41, 291-303.	2.8	172
13	Multivesicular compartments proliferate in susceptible and resistant MLA12 barley leaves in response to infection by the biotrophic powdery mildew fungus. <i>New Phytologist</i> , 2006, 172, 563-576.	3.5	172
14	Overexpression of barley BAX inhibitor 1 induces breakdown of mlo-mediated penetration resistance to <i>Blumeria graminis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 5555-5560.	3.3	171
15	BAX Inhibitor-1, an ancient cell death suppressor in animals and plants with prokaryotic relatives. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2004, 9, 299-307.	2.2	169
16	Do Plant Cells Secrete Exosomes Derived from Multivesicular Bodies?. <i>Plant Signaling and Behavior</i> , 2007, 2, 4-7.	1.2	152
17	Cell biology of the plant-powdery mildew interaction. <i>Current Opinion in Plant Biology</i> , 2011, 14, 738-746.	3.5	148
18	A Small GTP-Binding Host Protein Is Required for Entry of Powdery Mildew Fungus into Epidermal Cells of Barley. <i>Plant Physiology</i> , 2002, 128, 1447-1454.	2.3	147

#	ARTICLE	IF	CITATIONS
19	Bacterial medium-chain 3-hydroxy fatty acid metabolites trigger immunity in <i>Arabidopsis</i> plants. <i>Science</i> , 2019, 364, 178-181.	6.0	145
20	A Barley ROP GTPase ACTIVATING PROTEIN Associates with Microtubules and Regulates Entry of the Barley Powdery Mildew Fungus into Leaf Epidermal Cells. <i>Plant Cell</i> , 2011, 23, 2422-2439.	3.1	127
21	Tissue-Specific Superoxide Generation at Interaction Sites in Resistant and Susceptible Near-Isogenic Barley Lines Attacked by the Powdery Mildew Fungus ( <i>Erysiphe graminis</i> f. sp. <i>hordei</i> ). <i>Molecular Plant-Microbe Interactions</i> , 1998, 11, 292-300.	1.4	123
22	Functional analysis of barley RAC/ROP G-protein family members in susceptibility to the powdery mildew fungus. <i>Plant Journal</i> , 2003, 36, 589-601.	2.8	123
23	Biotic and Abiotic Stress Responses in Crop Plants. <i>Agronomy</i> , 2018, 8, 267.	1.3	117
24	Cell wall invertases, key enzymes in the modulation of plant metabolism during defence responses. <i>Molecular Plant Pathology</i> , 2014, 15, 858-864.	2.0	110
25	The Deubiquitinating Enzyme AMSH1 and the ESCRT-III Subunit VPS2.1 Are Required for Autophagic Degradation in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013, 25, 2236-2252.	3.1	107
26	The fungal core effector <i>Pep1</i> is conserved across smuts of dicots and monocots. <i>New Phytologist</i> , 2015, 206, 1116-1126.	3.5	100
27	Powdery mildew susceptibility and biotrophic infection strategies. <i>FEMS Microbiology Letters</i> , 2005, 245, 9-17.	0.7	95
28	The Barley Apoptosis Suppressor Homologue Bax Inhibitor-1 Compromises Nonhost Penetration Resistance of Barley to the Inappropriate Pathogen <i>Blumeria graminis</i> f. sp. <i>tritici</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 484-490.	1.4	90
29	Oomycete small RNAs bind to the plant RNA-induced silencing complex for virulence. <i>ELife</i> , 2020, 9, .	2.8	89
30	BAX INHIBITOR-1 Is Required for Full Susceptibility of Barley to Powdery Mildew. <i>Molecular Plant-Microbe Interactions</i> , 2010, 23, 1217-1227.	1.4	84
31	Apoplastic pH Signaling in Barley Leaves Attacked by the Powdery Mildew Fungus <i>Blumeria graminis</i> f. sp. <i>hordei</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 118-123.	1.4	83
32	Over-expression of the cell death regulator BAX inhibitor-1 in barley confers reduced or enhanced susceptibility to distinct fungal pathogens. <i>Theoretical and Applied Genetics</i> , 2009, 118, 455-463.	1.8	83
33	Ectopic Expression of Constitutively Activated RACB in Barley Enhances Susceptibility to Powdery Mildew and Abiotic Stress. <i>Plant Physiology</i> , 2005, 139, 353-362.	2.3	80
34	Differential expression of putative cell death regulator genes in near-isogenic, resistant and susceptible barley lines during interaction with the powdery mildew fungus. <i>Plant Molecular Biology</i> , 2001, 47, 739-748.	2.0	78
35	Bacteria-Triggered Systemic Immunity in Barley Is Associated with WRKY and ETHYLENE RESPONSIVE FACTORS But Not with Salicylic Acid. <i>Plant Physiology</i> , 2014, 166, 2133-2151.	2.3	76
36	Barley <i>Mla</i> and <i>Rar</i> mutants compromised in the hypersensitive cell death response against <i>Blumeria graminis</i> f.sp. <i>hordei</i> are modified in their ability to accumulate reactive oxygen intermediates at sites of fungal invasion. <i>Planta</i> , 2000, 212, 16-24.	1.6	68

#	ARTICLE	IF	CITATIONS
37	Root-to-shoot signalling: apoplastic alkalization, a general stress response and defence factor in barley ( <i>Hordeum vulgare</i> ). <i>Protoplasma</i> , 2005, 227, 17-24.	1.0	68
38	Polarized Defense Against Fungal Pathogens Is Mediated by the Jacalin-Related Lectin Domain of Modular Poaceae -Specific Proteins. <i>Molecular Plant</i> , 2016, 9, 514-527.	3.9	67
39	The White Barley Mutant Albostrians Shows a Supersusceptible but Symptomless Interaction Phenotype with the Hemibiotrophic Fungus <i>Bipolaris sorokiniana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 366-373.	1.4	66
40	Respiratory Burst Oxidase Homologue A of barley contributes to penetration by the powdery mildew fungus <i>Blumeria graminis</i> f. sp. <i>hordei</i> . <i>Journal of Experimental Botany</i> , 2006, 57, 3781-3791.	2.4	65
41	Constitutively activated barley ROPs modulate epidermal cell size, defense reactions and interactions with fungal leaf pathogens. <i>Plant Cell Reports</i> , 2008, 27, 1877-1887.	2.8	65
42	Accommodation of powdery mildew fungi in intact plant cells. <i>Journal of Plant Physiology</i> , 2008, 165, 5-18.	1.6	65
43	Non-host resistance of barley is associated with a hydrogen peroxide burst at sites of attempted penetration by wheat powdery mildew fungus. <i>Molecular Plant Pathology</i> , 2001, 2, 199-205.	2.0	63
44	RBOHF2 of Barley Is Required for Normal Development of Penetration Resistance to the Parasitic Fungus <i>Blumeria graminis</i> f. sp. <i>hordei</i> . <i>Molecular Plant-Microbe Interactions</i> , 2010, 23, 1143-1150.	1.4	60
45	Barley ROP Binding Kinase1 Is Involved in Microtubule Organization and in Basal Penetration Resistance to the Barley Powdery Mildew Fungus. <i>Plant Physiology</i> , 2012, 159, 311-320.	2.3	56
46	The conserved oligomeric G <sub>o</sub> complex is involved in penetration resistance of barley to the barley powdery mildew fungus. <i>Molecular Plant Pathology</i> , 2013, 14, 230-240.	2.0	54
47	Mechanistic and genetic overlap of barley host and non-host resistance to <i>Blumeria graminis</i> . <i>Molecular Plant Pathology</i> , 2004, 5, 389-396.	2.0	52
48	Reduced Carbohydrate Availability Enhances the Susceptibility of Arabidopsis toward <i>Colletotrichum higginsianum</i> . <i>Plant Physiology</i> , 2013, 162, 225-238.	2.3	51
49	Endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 of Arabidopsis (AtCEP1) is involved in pathogen defense. <i>Frontiers in Plant Science</i> , 2014, 5, 58.	1.7	51
50	Transport and secretion in plant-microbe interactions. <i>Current Opinion in Plant Biology</i> , 2007, 10, 573-579.	3.5	49
51	Barley RIC171 interacts with RACB in planta and supports entry of the powdery mildew fungus. <i>Cellular Microbiology</i> , 2008, 10, 1815-1826.	1.1	49
52	Cell-autonomous defense, reorganization and trafficking of membranes in plant-microbe interactions. <i>New Phytologist</i> , 2014, 204, 815-822.	3.5	47
53	Barley disease susceptibility factor RACB acts in epidermal cell polarity and positioning of the nucleus. <i>Journal of Experimental Botany</i> , 2016, 67, 3263-3275.	2.4	47
54	Superoxide and Hydrogen Peroxide Play Different Roles in the Nonhost Interaction of Barley and Wheat with Inappropriate formae speciales of <i>Blumeria graminis</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 304-312.	1.4	45

#	ARTICLE	IF	CITATIONS
55	Alcohol dehydrogenase 1 of barley modulates susceptibility to the parasitic fungus <i>Blumeria graminis</i> f.sp. <i>hordei</i> . <i>Journal of Experimental Botany</i> , 2011, 62, 3449-3457.	2.4	45
56	RIP3 and AtKinesin-13A – A novel interaction linking Rho proteins of plants to microtubules. <i>European Journal of Cell Biology</i> , 2010, 89, 906-916.	1.6	43
57	Co-immunoprecipitation-based identification of putative BAX INHIBITOR-1 interacting proteins involved in cell death regulation and plant-powdery mildew interactions. <i>Molecular Plant Pathology</i> , 2013, 14, 791-802.	2.0	42
58	Fate of <i>Fusarium</i> Toxins during the Malting Process. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 1377-1384.	2.4	41
59	Metrafenone: studies on the mode of action of a novel cereal powdery mildew fungicide. <i>Pest Management Science</i> , 2006, 62, 393-401.	1.7	40
60	Expression of barley BAX Inhibitor-1 in carrots confers resistance to <i>Botrytis cinerea</i> . <i>Molecular Plant Pathology</i> , 2006, 7, 279-284.	2.0	39
61	Good Riddance? Breaking Disease Susceptibility in the Era of New Breeding Technologies. <i>Agronomy</i> , 2018, 8, 114.	1.3	39
62	Genetic loss of susceptibility: a costly route to disease resistance?. <i>Plant Pathology</i> , 2013, 62, 56-62.	1.2	38
63	The formation of a camalexin-biosynthetic metabolon. <i>Plant Cell</i> , 2019, 31, tpc.00403.2019.	3.1	38
64	The Arabidopsis leucine-rich repeat receptor-like kinase MIK2 is a crucial component of early immune responses to a fungal-derived elicitor. <i>New Phytologist</i> , 2021, 229, 3453-3466.	3.5	38
65	A set of Arabidopsis genes involved in the accommodation of the downy mildew pathogen <i>Hyaloperonospora arabidopsidis</i> . <i>PLoS Pathogens</i> , 2019, 15, e1007747.	2.1	37
66	Mutations in <i>Ror1</i> and <i>Ror2</i> genes cause modification of hydrogen peroxide accumulation in <i>mlo</i> -barley under attack from the powdery mildew fungus. <i>Molecular Plant Pathology</i> , 2000, 1, 287-292.	2.0	36
67	Transient over-expression of barley BAX Inhibitor-1 weakens oxidative defence and MLA12-mediated resistance to <i>Blumeria graminis</i> f.sp. <i>hordei</i> . <i>Molecular Plant Pathology</i> , 2006, 7, 543-552.	2.0	36
68	CYP83A1 is required for metabolic compatibility of Arabidopsis with the adapted powdery mildew fungus <i>Erysiphe cruciferarum</i> . <i>New Phytologist</i> , 2014, 202, 1310-1319.	3.5	35
69	The Arabidopsis microtubule-associated protein MAP65-3 supports infection by filamentous biotrophic pathogens by down-regulating salicylic acid-dependent defenses. <i>Journal of Experimental Botany</i> , 2016, 67, 1731-1743.	2.4	35
70	A barley powdery mildew fungus non-autonomous retrotransposon encodes a peptide that supports penetration success on barley. <i>Journal of Experimental Botany</i> , 2018, 69, 3745-3758.	2.4	35
71	The Subcellular Localization of Tubby-Like Proteins and Participation in Stress Signaling and Root Colonization by the Mutualist <i>Piriformospora indica</i> . <i>Plant Physiology</i> , 2012, 160, 349-364.	2.3	34
72	The Receptor Kinase IMPAIRED OOMYCETE SUSCEPTIBILITY1 Attenuates Abscisic Acid Responses in Arabidopsis. <i>Plant Physiology</i> , 2014, 166, 1506-1518.	2.3	32

#	ARTICLE	IF	CITATIONS
73	A New Reference Genome Shows the One-Speed Genome Structure of the Barley Pathogen <i>Ramularia collo-cygni</i> . <i>Genome Biology and Evolution</i> , 2018, 10, 3243-3249.	1.1	30
74	Functional assessment of the pathogenesis-related protein PR-1b in barley. <i>Plant Science</i> , 2003, 165, 1275-1280.	1.7	29
75	LIFEGUARD proteins support plant colonization by biotrophic powdery mildew fungi. <i>Journal of Experimental Botany</i> , 2013, 64, 3855-3867.	2.4	29
76	Enemy at the gates: traffic at the plant cell pathogen interface. <i>Cellular Microbiology</i> , 2008, 10, 2400-2407.	1.1	28
77	Transgenic Suppression of Cell Death Limits Penetration Success of the Soybean Rust Fungus <i>Phakopsora pachyrhizi</i> into Epidermal Cells of Barley. <i>Phytopathology</i> , 2009, 99, 220-226.	1.1	27
78	Biotransformation of Pyrene by Cell Cultures of Soybean ( <i>Glycine max</i> L.), Wheat ( <i>Triticum aestivum</i> L.), Jimsonweed ( <i>Datura stramonium</i> L.), and Purple Foxglove ( <i>Digitalis purpurea</i> L.). <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 263-269.	2.4	26
79	Alternative Cell Death Mechanisms Determine Epidermal Resistance in Incompatible Barley- <i>Ustilago</i> Interactions. <i>Molecular Plant-Microbe Interactions</i> , 2014, 27, 403-414.	1.4	26
80	Macroarray expression analysis of barley susceptibility and nonhost resistance to <i>Blumeria graminis</i> . <i>Journal of Plant Physiology</i> , 2006, 163, 657-670.	1.6	24
81	Assessment and Introduction of Quantitative Resistance to Fusarium Head Blight in Elite Spring Barley. <i>Phytopathology</i> , 2013, 103, 1252-1259.	1.1	24
82	Cyanobacterial Phytochrome2 Regulates the Heterotrophic Metabolism and Has a Function in the Heat and High-Light Stress Response. <i>Plant Physiology</i> , 2014, 164, 2157-2166.	2.3	24
83	Silencing of <i>RBOHF2</i> Causes Leaf Age-Dependent Accelerated Senescence, Salicylic Acid Accumulation, and Powdery Mildew Resistance in Barley. <i>Molecular Plant-Microbe Interactions</i> , 2017, 30, 906-918.	1.4	24
84	The Arabidopsis ROP-activated receptor-like cytoplasmic kinase RLCK VI <sub>A3</sub> is involved in control of basal resistance to powdery mildew and trichome branching. <i>Plant Cell Reports</i> , 2015, 34, 457-468.	2.8	23
85	Recognition and defence of plant-infecting fungal pathogens. <i>Journal of Plant Physiology</i> , 2021, 256, 153324.	1.6	23
86	Pathogenesis-associated transcriptional patterns in Triticeae. <i>Journal of Plant Physiology</i> , 2011, 168, 9-19.	1.6	22
87	The effective papilla hypothesis. <i>New Phytologist</i> , 2014, 204, 438-440.	3.5	22
88	Effect of nitrogen fertilization on Fusarium head blight in spring barley. <i>Crop Protection</i> , 2016, 88, 18-27.	1.0	22
89	A barley SKP1-like protein controls abundance of the susceptibility factor RACB and influences the interaction of barley with the barley powdery mildew fungus. <i>Molecular Plant Pathology</i> , 2016, 17, 184-195.	2.0	20
90	RGL1-GOLVEN signaling promotes cell surface immune receptor abundance to regulate plant immunity. <i>EMBO Reports</i> , 2022, 23, e53281.	2.0	20

#	ARTICLE	IF	CITATIONS
91	Involvement of <i>Arabidopsis thaliana</i> endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 (AtCEP1) in powdery mildew-induced and AtCPR5-controlled cell death. <i>PLoS ONE</i> , 2017, 12, e0183870.	1.1	19
92	Ramularia leaf spot disease of barley is highly host genotype-dependent and suppressed by continuous drought stress in the field. <i>Journal of Plant Diseases and Protection</i> , 2021, 128, 749-767.	1.6	19
93	Interactive signal transfer between host and pathogen during successful infection of barley leaves by <i>Blumeria graminis</i> and <i>Bipolaris sorokiniana</i> . <i>Journal of Plant Physiology</i> , 2008, 165, 52-59.	1.6	18
94	The White Barley Mutant <i>Albostrians</i> Shows Enhanced Resistance to the Biotroph <i>Blumeria graminis</i> f. sp. <i>hordei</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 374-382.	1.4	17
95	Model Wheat Genotypes as Tools to Uncover Effective Defense Mechanisms Against the Hemibiotrophic Fungus <i>Bipolaris sorokiniana</i> . <i>Phytopathology</i> , 2005, 95, 528-532.	1.1	17
96	The Current Epidemic of the Barley Pathogen <i>Ramularia collo-cygni</i> Derives from a Population Expansion and Shows Global Admixture. <i>Phytopathology</i> , 2019, 109, 2161-2168.	1.1	17
97	PAMP-triggered immune responses in barley and susceptibility to powdery mildew. <i>Plant Signaling and Behavior</i> , 2016, 11, e1197465.	1.2	16
98	Influence of inoculum and climatic factors on the severity of <i>Fusarium</i> head blight in German spring and winter barley. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2016, 33, 489-499.	1.1	15
99	Barley ROP-Interactive Partner-a organizes into RAC1- and MICROTUBULE-ASSOCIATED ROP-GTPASE ACTIVATING PROTEIN 1-dependent membrane domains. <i>BMC Plant Biology</i> , 2020, 20, 94.	1.6	15
100	Barley ADH-1 modulates susceptibility to Bgh and is involved in chitin-induced systemic resistance. <i>Plant Physiology and Biochemistry</i> , 2018, 123, 281-287.	2.8	14
101	ROPGAPs of <i>Arabidopsis</i> limit susceptibility to powdery mildew. <i>Plant Signaling and Behavior</i> , 2011, 6, 1691-1694.	1.2	13
102	Influence of <i>Fusarium</i> isolates on the expression of barley genes related to plant defense and malting quality. <i>Journal of Cereal Science</i> , 2016, 69, 17-24.	1.8	13
103	Regulation and Functions of ROP GTPases in Plant-Microbe Interactions. <i>Cells</i> , 2020, 9, 2016.	1.8	13
104	Population studies of the wild tomato species <i>Solanum chilense</i> reveal geographically structured major gene-mediated pathogen resistance. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020, 287, 20202723.	1.2	13
105	MILDEW LOCUS O Mutation Does Not Affect Resistance to Grain Infections with <i>Fusarium</i> spp. and <i>Ramularia collo-cygni</i> . <i>Phytopathology</i> , 2015, 105, 1214-1219.	1.1	12
106	Occurrence of <i>sdh</i> Mutations in German <i>Alternaria solani</i> Isolates and Potential Impact on Boscalid Sensitivity In Vitro, in the Greenhouse, and in the Field. <i>Plant Disease</i> , 2019, 103, 3065-3071.	0.7	12
107	Superoxide Generation in Chemically Activated Resistance of Barley in Response to Inoculation with the Powdery Mildew Fungus. <i>Journal of Phytopathology</i> , 1999, 147, 1-4.	0.5	11
108	Infection of barley with the parasitic fungus <i>Blumeria graminis</i> f.sp. <i>hordei</i> results in the induction of <i>HvADH1</i> and <i>HvADH2</i> . <i>Plant Signaling and Behavior</i> , 2011, 6, 1584-1587.	1.2	11

#	ARTICLE	IF	CITATIONS
109	Barley susceptibility factor RACB modulates transcript levels of signalling protein genes in compatible interaction with <i>Blumeria graminis</i> f.sp. <i>hordei</i> . <i>Molecular Plant Pathology</i> , 2018, 19, 393-404.	2.0	11
110	ROP INTERACTIVE PARTNER b Interacts with RACB and Supports Fungal Penetration into Barley Epidermal Cells. <i>Plant Physiology</i> , 2020, 184, 823-836.	2.3	11
111	Steroidal Saponins – New Sources to Develop Potato ( <i>Solanum tuberosum</i> L.) Genotypes Resistant against Certain <i>Phytophthora infestans</i> Strains. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 7447-7459.	2.4	11
112	Fusarium infection of malting barley has to be managed over the entire value chain. <i>Journal of Plant Diseases and Protection</i> , 2018, 125, 1-4.	1.6	10
113	Superoxide Generation in Chemically Activated Resistance of Barley in Response to Inoculation with the Powdery Mildew Fungus. Superoxid Bildung während Chemisch Aktivierter Resistenz in Gerste als Reaktion auf Inokulation mit Echtem Gerstenmehltau. <i>Journal of Phytopathology</i> , 1999, 147, 1-4.	0.5	10
114	A barley Engulfment and Motility domain containing protein modulates Rho GTPase activating protein HvMAGAP1 function in the barley powdery mildew interaction. <i>Plant Molecular Biology</i> , 2014, 84, 469-478.	2.0	9
115	Evolutionarily conserved partial gene duplication in the Triticeae tribe of grasses confers pathogen resistance. <i>Genome Biology</i> , 2018, 19, 116.	3.8	9
116	Analysis of archive samples of spring and winter barley support an increase in individual Fusarium species in Bavarian barley grain over the last decades. <i>Journal of Plant Diseases and Protection</i> , 2019, 126, 247-254.	1.6	9
117	Biosynthesis of $\pm$ -solanine and $\pm$ -chaconine in potato leaves ( <i>Solanum tuberosum</i> L.) – A $^{13}\text{C}$ study. <i>Food Chemistry</i> , 2021, 365, 130461.	4.2	9
118	<i>Fusarium</i> Species on Barley Malt: Is Visual Assessment an Appropriate Tool for Detection?. <i>Cereal Chemistry</i> , 2017, 94, 659-669.	1.1	8
119	Re-evaluation of superoxide scavenging capacity of xanthohumol. <i>Free Radical Research</i> , 2010, 44, 1435-1444.	1.5	7
120	The Role of Seven-Transmembrane Domain MLO Proteins, Heterotrimeric G-Proteins, and Monomeric RAC/ROPs in Plant Defense. <i>Signaling and Communication in Plants</i> , 2010, , 197-220.	0.5	6
121	Whole-genome sequencing elucidates the species-wide diversity and evolution of fungicide resistance in the early blight pathogen <i>Alternaria solani</i> . <i>Evolutionary Applications</i> , 2022, 15, 1605-1620.	1.5	6
122	Ectopic expression of barley constitutively activated ROPs supports susceptibility to powdery mildew and bacterial wildfire in tobacco. <i>European Journal of Plant Pathology</i> , 2009, 125, 317-327.	0.8	5
123	Quantitative resistance differences between and within natural populations of <i>Solanum chilense</i> against the oomycete pathogen <i>Phytophthora infestans</i> . <i>Ecology and Evolution</i> , 2021, 11, 7768-7778.	0.8	5
124	Host Genotype and Weather Effects on Fusarium Head Blight Severity and Mycotoxin Load in Spring Barley. <i>Toxins</i> , 2022, 14, 125.	1.5	5
125	Plant small monomeric G-proteins (RAC/ROPs) of barley are common elements of susceptibility to fungal leaf pathogens, cell expansion and stomata development. <i>Plant Signaling and Behavior</i> , 2009, 4, 109-110.	1.2	4
126	Posttranslational modification of the RHO of plants protein RACB by phosphorylation and cross-kingdom conserved ubiquitination. <i>PLoS ONE</i> , 2022, 17, e0258924.	1.1	4



#	ARTICLE	IF	CITATIONS
127	Quantitative disease resistance and fungal pathogenicity in Triticeae. <i>Journal of Plant Physiology</i> , 2011, 168, 1-2.	1.6	3
128	TECHNIQUE TO ASSESS GENE FUNCTION IN HOP ( <i>HUMULUS LUPULUS L.</i> ) - POWDERY MILDEW INTERACTIONS. <i>Acta Horticulturae</i> , 2013, , 59-66.	0.1	1
129	Special issue "Deepen knowledge in plant pathology for innovative agroecology". <i>European Journal of Plant Pathology</i> , 2018, 152, 853-854.	0.8	0