

Mayuresh S Korgaonkar

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3036464/publications.pdf>

Version: 2024-02-01

109
papers

5,649
citations

125106

35
h-index

100535

70
g-index

117
all docs

117
docs citations

117
times ranked

10074
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Negative association between anterior insula activation and resilience during sustained attention: an fMRI twin study. <i>Psychological Medicine</i> , 2023, 53, 3187-3199. | 2.7 | 2 |
| 2 | Brainmarker-I Differentially Predicts Remission to Various Attention-Deficit/Hyperactivity Disorder Treatments: A Discovery, Transfer, and Blinded Validation Study. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2023, 8, 52-60. | 1.1 | 11 |
| 3 | Cognitive and Executive Contributions to Trail-Making Task Performance on Adolescents With and Without Attention Deficit Hyperactivity Disorder. <i>Journal of Attention Disorders</i> , 2022, 26, 881-892. | 1.5 | 1 |
| 4 | Intrinsic Functional Connectomes Characterize Neuroticism in Major Depressive Disorder and Predict Antidepressant Treatment Outcomes. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2022, 7, 276-284. | 1.1 | 3 |
| 5 | Abnormal habenula functional connectivity characterizes treatment-resistant depression. <i>NeuroImage: Clinical</i> , 2022, 34, 102990. | 1.4 | 12 |
| 6 | Remodeling of the Cortical Structural Connectome in Posttraumatic Stress Disorder: Results From the ENIGMA-PGC Posttraumatic Stress Disorder Consortium. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2022, 7, 935-948. | 1.1 | 2 |
| 7 | Genetic variants associated with longitudinal changes in brain structure across the lifespan. <i>Nature Neuroscience</i> , 2022, 25, 421-432. | 7.1 | 75 |
| 8 | Neural processes during response inhibition in complex posttraumatic stress disorder. <i>Depression and Anxiety</i> , 2022, 39, 307-314. | 2.0 | 4 |
| 9 | Altered resting-state neural networks in children and adolescents with functional neurological disorder. <i>NeuroImage: Clinical</i> , 2022, 35, 103110. | 1.4 | 7 |
| 10 | Inhibition-related modulation of salience and frontoparietal networks predicts cognitive control ability and inattention symptoms in children with ADHD. <i>Molecular Psychiatry</i> , 2021, 26, 4016-4025. | 4.1 | 48 |
| 11 | Altered white matter microstructural organization in posttraumatic stress disorder across 3047 adults: results from the PGC-ENIGMA PTSD consortium. <i>Molecular Psychiatry</i> , 2021, 26, 4315-4330. | 4.1 | 69 |
| 12 | Cortical volume abnormalities in posttraumatic stress disorder: an ENIGMA-psychiatric genomics consortium PTSD workgroup mega-analysis. <i>Molecular Psychiatry</i> , 2021, 26, 4331-4343. | 4.1 | 52 |
| 13 | Distinct neural mechanisms of emotional processing in prolonged grief disorder. <i>Psychological Medicine</i> , 2021, 51, 587-595. | 2.7 | 14 |
| 14 | The distinctive neural circuitry of complex posttraumatic stress disorder during threat processing. <i>Psychological Medicine</i> , 2021, 51, 1121-1128. | 2.7 | 16 |
| 15 | White matter anisotropy and response to cognitive behavior therapy for posttraumatic stress disorder. <i>Translational Psychiatry</i> , 2021, 11, 14. | 2.4 | 3 |
| 16 | Investigating neural circuits of emotion regulation to distinguish euthymic patients with bipolar disorder and major depressive disorder. <i>Bipolar Disorders</i> , 2021, 23, 284-294. | 1.1 | 12 |
| 17 | Structural brain network topology underpinning ADHD and response to methylphenidate treatment. <i>Translational Psychiatry</i> , 2021, 11, 150. | 2.4 | 23 |
| 18 | Neural activity during response inhibition associated with improvement of dysphoric symptoms of PTSD after trauma-focused psychotherapy—an EEG-fMRI study. <i>Translational Psychiatry</i> , 2021, 11, 218. | 2.4 | 10 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | The functional connectome in posttraumatic stress disorder. <i>Neurobiology of Stress</i> , 2021, 14, 100321. | 1.9 | 15 |
| 20 | Neural activity during response inhibition in mild traumatic brain injury and posttraumatic stress disorder. <i>Neurobiology of Stress</i> , 2021, 14, 100308. | 1.9 | 4 |
| 21 | Abnormalities in Habenula Functional Connectivity Characterize Treatment-Resistant Depression. <i>Biological Psychiatry</i> , 2021, 89, S352. | 0.7 | 0 |
| 22 | No support for white matter connectivity differences in the combined and inattentive ADHD presentations. <i>PLoS ONE</i> , 2021, 16, e0245028. | 1.1 | 4 |
| 23 | Default-mode and fronto-parietal network connectivity during rest distinguishes asymptomatic patients with bipolar disorder and major depressive disorder. <i>Translational Psychiatry</i> , 2021, 11, 547. | 2.4 | 29 |
| 24 | Neural correlates of emotional processing in panic disorder. <i>NeuroImage: Clinical</i> , 2021, 32, 102902. | 1.4 | 4 |
| 25 | Trauma and posttraumatic stress disorder modulate polygenic predictors of hippocampal and amygdala volume. <i>Translational Psychiatry</i> , 2021, 11, 637. | 2.4 | 4 |
| 26 | Precision in psychiatryâ€”A roadmap to translate neurobiological measures to the clinic. <i>Bipolar Disorders</i> , 2021, 23, 747-750. | 1.1 | 2 |
| 27 | Connectivity of the Cognitive Control Network During Response Inhibition as a Predictive and Response Biomarker in Major Depression: Evidence From a Randomized Clinical Trial. <i>Biological Psychiatry</i> , 2020, 87, 462-472. | 0.7 | 42 |
| 28 | Intrinsic connectomes are a predictive biomarker of remission in major depressive disorder. <i>Molecular Psychiatry</i> , 2020, 25, 1537-1549. | 4.1 | 99 |
| 29 | Investigating the neural basis of cognitive control dysfunction in mood disorders. <i>Bipolar Disorders</i> , 2020, 22, 286-295. | 1.1 | 22 |
| 30 | Intrinsic functional connectivity of the default mode and cognitive control networks relate to change in behavioral performance over two years. <i>Cortex</i> , 2020, 132, 180-190. | 1.1 | 8 |
| 31 | Intrinsic connectomes underlying response to trauma-focused psychotherapy in post-traumatic stress disorder. <i>Translational Psychiatry</i> , 2020, 10, 270. | 2.4 | 15 |
| 32 | The role of progressive oral implant rehabilitation in mastication, cognition and oral healthâ€”related quality of life outcomesâ€”A pilot to define the protocol. <i>Journal of Oral Rehabilitation</i> , 2020, 47, 1368-1381. | 1.3 | 14 |
| 33 | Reappraisal-related neural predictors of treatment response to cognitive behavior therapy for post-traumatic stress disorder. <i>Psychological Medicine</i> , 2020, 51, 1-11. | 2.7 | 15 |
| 34 | A Systematic Review of Imaging Studies in the Combined and Inattentive Subtypes of Attention Deficit Hyperactivity Disorder. <i>Frontiers in Integrative Neuroscience</i> , 2020, 14, 31. | 1.0 | 46 |
| 35 | Differential neural predictors of treatment response for fear and dysphoric features of posttraumatic stress disorder. <i>Depression and Anxiety</i> , 2020, 37, 1026-1036. | 2.0 | 10 |
| 36 | Neural Circuits Underlying Treatment-Resistant Depression. <i>Biological Psychiatry</i> , 2020, 87, S311. | 0.7 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Neurophysiological markers of attention distinguish bipolar disorder and unipolar depression. <i>Journal of Affective Disorders</i> , 2020, 274, 411-419. | 2.0 | 9 |
| 38 | Understanding the neural mechanisms of lisdexamfetamine dimesylate (LDX) pharmacotherapy in Binge Eating Disorder (BED): a study protocol. <i>Journal of Eating Disorders</i> , 2019, 7, 23. | 1.3 | 15 |
| 39 | Profound and reproducible patterns of reduced regional gray matter characterize major depressive disorder. <i>Translational Psychiatry</i> , 2019, 9, 176. | 2.4 | 21 |
| 40 | Gender-specific structural abnormalities in major depressive disorder revealed by voxel-based analysis. <i>NeuroImage: Clinical</i> , 2019, 21, 101668. | 1.4 | 20 |
| 41 | The effects of bullying in depression on white matter integrity. <i>Behavioural Brain Research</i> , 2019, 363, 149-154. | 1.2 | 7 |
| 42 | Characterizing neurocognitive markers of familial risk for depression using multi-modal imaging, behavioral and self-report measures. <i>Journal of Affective Disorders</i> , 2019, 253, 336-342. | 2.0 | 18 |
| 43 | Reply to: Two Methodologies in "Amygdala Activation and Connectivity to Emotional Processing Distinguishes Asymptomatic Patients With Bipolar Disorders and Unipolar Depression" That Can Produce False-Positive Results and Some Statistical Recommendations. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 411-413. | 1.1 | 0 |
| 44 | A Neuroethics Framework for the Australian Brain Initiative. <i>Neuron</i> , 2019, 101, 365-369. | 3.8 | 11 |
| 45 | Understanding autism spectrum disorder and social functioning in children with neurofibromatosis type 1: protocol for a cross-sectional multimodal study. <i>BMJ Open</i> , 2019, 9, e030601. | 0.8 | 11 |
| 46 | Amygdala Activation and Connectivity to Emotional Processing Distinguishes Asymptomatic Patients With Bipolar Disorders and Unipolar Depression. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 361-370. | 1.1 | 30 |
| 47 | Diffusion Tensor Imaging Analysis of Mild Traumatic Brain Injury and Posttraumatic Stress Disorder. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 81-90. | 1.1 | 10 |
| 48 | "Motoring in idle" The default mode and somatomotor networks are overactive in children and adolescents with functional neurological symptoms. <i>NeuroImage: Clinical</i> , 2018, 18, 730-743. | 1.4 | 32 |
| 49 | Intrinsic functional connectivity predicts remission on antidepressants: a randomized controlled trial to identify clinically applicable imaging biomarkers. <i>Translational Psychiatry</i> , 2018, 8, 57. | 2.4 | 79 |
| 50 | Impaired engagement of the ventral attention system in neurofibromatosis type 1. <i>Brain Imaging and Behavior</i> , 2018, 12, 499-508. | 1.1 | 12 |
| 51 | Smaller Hippocampal Volume in Posttraumatic Stress Disorder: A Multisite ENIGMA-PGC Study: Subcortical Volumetry Results From Posttraumatic Stress Disorder Consortia. <i>Biological Psychiatry</i> , 2018, 83, 244-253. | 0.7 | 335 |
| 52 | Profiling risk for depressive disorder by circuit, behavior and self-report measures of emotion function. <i>Journal of Affective Disorders</i> , 2018, 227, 595-602. | 2.0 | 10 |
| 53 | A negative association between brainstem pontine grey-matter volume, well-being and resilience in healthy twins. <i>Journal of Psychiatry and Neuroscience</i> , 2018, 43, 386-395. | 1.4 | 15 |
| 54 | Effects of methylphenidate on cognition and behaviour in children with neurofibromatosis type 1: a study protocol for a randomised placebo-controlled crossover trial. <i>BMJ Open</i> , 2018, 8, e021800. | 0.8 | 12 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | T59. Does White Matter Microstructural Integrity Differ in the Combined and Inattentive Subtypes of ADHD? A Diffusion Tensor Imaging Study. <i>Biological Psychiatry</i> , 2018, 83, S151. | 0.7 | 2 |
| 56 | Cognitive ability is associated with changes in the functional organization of the cognitive control brain network. <i>Human Brain Mapping</i> , 2018, 39, 5028-5038. | 1.9 | 22 |
| 57 | Gray Matter Atrophy in the Cerebellum—Evidence of Increased Vulnerability of the Crus and Vermis with Advancing Age. <i>Cerebellum</i> , 2017, 16, 388-397. | 1.4 | 11 |
| 58 | EEG connectivity between the subgenual anterior cingulate and prefrontal cortices in response to antidepressant medication. <i>European Neuropsychopharmacology</i> , 2017, 27, 301-312. | 0.3 | 32 |
| 59 | 247. Structural Networks Characterise Methylphenidate Treatment Response in ADHD. <i>Biological Psychiatry</i> , 2017, 81, S101-S102. | 0.7 | 1 |
| 60 | 253. Functional Connectome Networks Underlying Outcomes of Antidepressant Medication in Major Depressive Disorders. <i>Biological Psychiatry</i> , 2017, 81, S104. | 0.7 | 1 |
| 61 | 326. Clustering by Salience Network Activation to Emotional Faces Identifies a Transdiagnostic Subtype that is Associated with Specific Interoceptive Related Symptoms. <i>Biological Psychiatry</i> , 2017, 81, S133-S134. | 0.7 | 0 |
| 62 | Regional brain network organization distinguishes the combined and inattentive subtypes of Attention Deficit Hyperactivity Disorder. <i>NeuroImage: Clinical</i> , 2017, 15, 383-390. | 1.4 | 54 |
| 63 | The neural basis of deficient response inhibition in children with neurofibromatosis type 1: Evidence from a functional MRI study. <i>Cortex</i> , 2017, 93, 1-11. | 1.1 | 14 |
| 64 | Grey matter abnormalities in children and adolescents with functional neurological symptom disorder. <i>NeuroImage: Clinical</i> , 2017, 15, 306-314. | 1.4 | 49 |
| 65 | Brain functional connectome abnormalities in amyotrophic lateral sclerosis are associated with disability and cortical hyperexcitability. <i>European Journal of Neurology</i> , 2017, 24, 1507-1517. | 1.7 | 23 |
| 66 | 87. Volume of Sub-Cortical Structures in Posttraumatic Stress Disorder from Multi-Site Investigation by ENIGMA and PGC Consortia. <i>Biological Psychiatry</i> , 2017, 81, S36-S37. | 0.7 | 2 |
| 67 | Cognitive control network anatomy correlates with neurocognitive behavior: A longitudinal study. <i>Human Brain Mapping</i> , 2017, 38, 631-643. | 1.9 | 73 |
| 68 | Functional and structural connectivity in als: insights from mri connectome analyses and tms. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2017, 88, e1.17-e1. | 0.9 | 0 |
| 69 | Human amygdala engagement moderated by early life stress exposure is a biobehavioral target for predicting recovery on antidepressants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 11955-11960. | 3.3 | 50 |
| 70 | Altered gray matter organization in children and adolescents with ADHD: a structural covariance connectome study. <i>Translational Psychiatry</i> , 2016, 6, e947-e947. | 2.4 | 32 |
| 71 | Potential structural and functional biomarkers of upper motor neuron dysfunction in ALS. <i>Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration</i> , 2016, 17, 85-92. | 1.1 | 32 |
| 72 | Is the Alzheimer's disease cortical thickness signature a biological marker for memory?. <i>Brain Imaging and Behavior</i> , 2016, 10, 517-523. | 1.1 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Frontoparietal Activation During Response Inhibition Predicts Remission to Antidepressants in Patients With Major Depression. <i>Biological Psychiatry</i> , 2016, 79, 274-281. | 0.7 | 116 |
| 74 | Effects of TORC1 Inhibition during the Early and Established Phases of Polycystic Kidney Disease. <i>PLoS ONE</i> , 2016, 11, e0164193. | 1.1 | 15 |
| 75 | Prediction of Nonremission to Antidepressant Therapy Using Diffusion Tensor Imaging. <i>Journal of Clinical Psychiatry</i> , 2016, 77, e436-e443. | 1.1 | 29 |
| 76 | Magnetic Resonance Imaging Measures of Brain Structure to Predict Antidepressant Treatment Outcome in Major Depressive Disorder. <i>EBioMedicine</i> , 2015, 2, 37-45. | 2.7 | 53 |
| 77 | Identification of a Common Neurobiological Substrate for Mental Illness. <i>JAMA Psychiatry</i> , 2015, 72, 305. | 6.0 | 1,050 |
| 78 | Imaging predictors of remission to anti-depressant medications in major depressive disorder. <i>Journal of Affective Disorders</i> , 2015, 186, 134-144. | 2.0 | 38 |
| 79 | Amygdala Reactivity to Emotional Faces in the Prediction of General and Medication-Specific Responses to Antidepressant Treatment in the Randomized iSPOT-D Trial. <i>Neuropsychopharmacology</i> , 2015, 40, 2398-2408. | 2.8 | 168 |
| 80 | COGNITION-CHILDHOOD MALTREATMENT INTERACTIONS IN THE PREDICTION OF ANTIDEPRESSANT OUTCOMES IN MAJOR DEPRESSIVE DISORDER PATIENTS: RESULTS FROM THE iSPOT-D TRIAL. <i>Depression and Anxiety</i> , 2015, 32, 594-604. | 2.0 | 64 |
| 81 | Emotion circuits differentiate symptoms of psychosis versus mania in adolescents. <i>Neurocase</i> , 2015, 21, 592-600. | 0.2 | 0 |
| 82 | Tractography of the Brainstem in Major Depressive Disorder Using Diffusion Tensor Imaging. <i>PLoS ONE</i> , 2014, 9, e84825. | 1.1 | 33 |
| 83 | Cerebral responses to innocuous somatic pressure stimulation following aerobic exercise rehabilitation in chronic pain patients: a functional magnetic resonance imaging study. <i>International Journal of General Medicine</i> , 2014, 7, 425. | 0.8 | 11 |
| 84 | Body mass index and brain structure in healthy children and adolescents. <i>International Journal of Neuroscience</i> , 2014, 124, 49-55. | 0.8 | 100 |
| 85 | Pyrrolidine dithiocarbamate reduces the progression of total kidney volume and cyst enlargement in experimental polycystic kidney disease. <i>Physiological Reports</i> , 2014, 2, e12196. | 0.7 | 13 |
| 86 | Establishing the resting state default mode network derived from functional magnetic resonance imaging tasks as an endophenotype: A twins study. <i>Human Brain Mapping</i> , 2014, 35, 3893-3902. | 1.9 | 56 |
| 87 | The genetic and neuroanatomical basis of social dysfunction: Lessons from neurofibromatosis type 1. <i>Human Brain Mapping</i> , 2014, 35, 2372-2382. | 1.9 | 30 |
| 88 | Thalamic volume and thalamo-cortical white matter tracts correlate with motor and verbal memory performance. <i>NeuroImage</i> , 2014, 91, 77-83. | 2.1 | 36 |
| 89 | Diffusion tensor imaging predictors of treatment outcomes in major depressive disorder. <i>British Journal of Psychiatry</i> , 2014, 205, 321-328. | 1.7 | 126 |
| 90 | Abnormal Structural Networks Characterize Major Depressive Disorder: A Connectome Analysis. <i>Biological Psychiatry</i> , 2014, 76, 567-574. | 0.7 | 293 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Impact of early vs. late childhood early life stress on brain morphometrics. <i>Brain Imaging and Behavior</i> , 2013, 7, 196-203. | 1.1 | 134 |
| 92 | Brain imaging predictors and the international study to predict optimized treatment for depression: study protocol for a randomized controlled trial. <i>Trials</i> , 2013, 14, 224. | 0.7 | 34 |
| 93 | Widespread reductions in gray matter volume in depression. <i>NeuroImage: Clinical</i> , 2013, 3, 332-339. | 1.4 | 301 |
| 94 | Using Standardized fMRI Protocols to Identify Patterns of Prefrontal Circuit Dysregulation that are Common and Specific to Cognitive and Emotional Tasks in Major Depressive Disorder: First Wave Results from the iSPOT-D Study. <i>Neuropsychopharmacology</i> , 2013, 38, 863-871. | 2.8 | 113 |
| 95 | Chronic effects of dietary vitamin D deficiency without increased calcium supplementation on the progression of experimental polycystic kidney disease. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 305, F574-F582. | 1.3 | 14 |
| 96 | Higher education is an age-independent predictor of white matter integrity and cognitive control in late adolescence. <i>Developmental Science</i> , 2013, 16, 653-664. | 1.3 | 88 |
| 97 | Neuroplasticity in the Adaptation to Prosthodontic Treatment. <i>Journal of Orofacial Pain</i> , 2013, 27, 206-216. | 1.7 | 58 |
| 98 | Early Exposure to Traumatic Stressors Impairs Emotional Brain Circuitry. <i>PLoS ONE</i> , 2013, 8, e75524. | 1.1 | 31 |
| 99 | GSK3B and MAPT Polymorphisms Are Associated with Grey Matter and Intracranial Volume in Healthy Individuals. <i>PLoS ONE</i> , 2013, 8, e71750. | 1.1 | 8 |
| 100 | The TWIN-E Project in Emotional Wellbeing: Study Protocol and Preliminary Heritability Results Across Four MRI and DTI Measures. <i>Twin Research and Human Genetics</i> , 2012, 15, 419-441. | 0.3 | 40 |
| 101 | Mapping inter-regional connectivity of the entire cortex to characterize major depressive disorder. <i>NeuroReport</i> , 2012, 23, 566-571. | 0.6 | 54 |
| 102 | P.2.c.014 Prediction of antidepressant response in the iSPOT-D trial from baseline fMRI – preliminary findings. <i>European Neuropsychopharmacology</i> , 2012, 22, S258. | 0.3 | 0 |
| 103 | Testing the white matter retrogenesis hypothesis of cognitive aging. <i>Neurobiology of Aging</i> , 2012, 33, 1699-1715. | 1.5 | 139 |
| 104 | Hippocampal volume varies with educational attainment across the life-span. <i>Frontiers in Human Neuroscience</i> , 2012, 6, 307. | 1.0 | 109 |
| 105 | Regional heterogeneity in limbic maturational changes: Evidence from integrating cortical thickness, volumetric and diffusion tensor imaging measures. <i>NeuroImage</i> , 2011, 55, 868-879. | 2.1 | 55 |
| 106 | Obesity Is Associated With Reduced White Matter Integrity in Otherwise Healthy Adults*. <i>Obesity</i> , 2011, 19, 500-504. | 1.5 | 204 |
| 107 | Loss of white matter integrity in major depressive disorder: Evidence using tract-based spatial statistical analysis of diffusion tensor imaging. <i>Human Brain Mapping</i> , 2011, 32, 2161-2171. | 1.9 | 180 |
| 108 | Limbic dysregulation is associated with lowered heart rate variability and increased trait anxiety in healthy adults. <i>Human Brain Mapping</i> , 2009, 30, 47-58. | 1.9 | 72 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Intrinsic Functional Connectivity in the Default Mode Network Differentiates the Combined and Inattentive Attention Deficit Hyperactivity Disorder Types. <i>Frontiers in Human Neuroscience</i> , 0, 16, . | 1.0 | 4 |