## Caroline A Antler

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/30402/publications.pdf

Version: 2024-02-01

1937685 1720034 18 71 4 7 citations h-index g-index papers 18 18 18 25 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Residual, differential neurobehavioral deficits linger after multiple recovery nights following chronic sleep restriction or acute total sleep deprivation. Sleep, 2021, 44, .	1.1	32
2	Cognitive throughput and working memory raw scores consistently differentiate resilient and vulnerable groups to sleep loss. Sleep, 2021, 44, .	1.1	12
3	Raw scores on subjective sleepiness, fatigue, and vigor metrics consistently define resilience and vulnerability to sleep loss. Sleep, 2022, 45, .	1.1	9
4	Concordance of multiple methods to define resiliency and vulnerability to sleep loss depends on Psychomotor Vigilance Test metric. Sleep, 2022, 45, .	1.1	7
5	Cortisol and C-Reactive Protein Vary During Sleep Loss and Recovery but Are Not Markers of Neurobehavioral Resilience. Frontiers in Physiology, 2021, 12, 782860.	2.8	6
6	The Symptom Structure of Seasonal Affective Disorder: Integrating Results from Factor and Network Analyses in the Context of the Dual Vulnerability Model. Journal of Psychopathology and Behavioral Assessment, 2021, 43, 95-107.	1.2	3
7	Acute sleep deprivation in humans. , 2023, , 217-229.		1
8	The 3-Minute Psychomotor Vigilance Test Demonstrates Inadequate Convergent Validity Relative to the 10-Minute Psychomotor Vigilance Test Across Sleep Loss and Recovery. Frontiers in Neuroscience, 2022, 16, 815697.	2.8	1
9	121 Behavioral Attention Relationships Vary Between Demographic Groups Across Sleep Loss and Recovery. Sleep, 2021, 44, A49-A50.	1.1	O
10	128 Raw Scores Best Differentiate Resilience and Vulnerability to Sleep Loss for Cognitive Throughput and Working Memory. Sleep, 2021, 44, A52-A52.	1.1	0
11	116 Behavioral Attention Raw Scores Best Differentiate Cognitive Resilience and Vulnerability to Sleep Loss. Sleep, 2021, 44, A47-A48.	1.1	O
12	127 Cognitive Throughput, Behavioral Attention, and Sleepiness Show Robust Relationships During Sleep Loss but Not During Recovery. Sleep, 2021, 44, A52-A52.	1.1	0
13	118 Relationships Between Perceptions of Subjective States Differ by Sleep Loss and During Recovery in Healthy Adults. Sleep, 2021, 44, A48-A48.	1.1	0
14	117 Comparison of Various Methods to Differentiate Resilience and Vulnerability to Sleep Loss Using Self-Rated Measures. Sleep, 2021, 44, A48-A48.	1.1	0
15	115 Age and Sex Differences in Behavioral Attention Across Baseline, Sleep Loss, and Recovery. Sleep, 2021, 44, A47-A47.	1.1	O
16	122 Different Duration Psychomotor Vigilance Tests Show Robust Stable Relationships Across Sleep Loss That Deteriorate in Recovery. Sleep, 2021, 44, A50-A50.	1.1	0
17	119 Behavioral Attention and Sleepiness Display Robust Stable Relationships Across Sleep Loss but not Across Recovery. Sleep, 2021, 44, A48-A49.	1.1	0
18	0283 Neurobehavioral Resilience and Vulnerability to Sleep Loss Differs Between Objective and Self-Rated Metrics Regardless of Categorization Method Utilized. Sleep, 2022, 45, A128-A128.	1.1	O