Luda Diatchenko

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3041993/luda-diatchenko-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118	10,195	47	100
papers	citations	h-index	g-index
127 ext. papers	11,926 ext. citations	8.6 avg, IF	5.54 L-index

#	Paper	IF	Citations
118	Single-cell RNA sequencing reveals time- and sex-specific responses of mouse spinal cord microglia to peripheral nerve injury and links ApoE to chronic pain <i>Nature Communications</i> , 2022 , 13, 843	17.4	5
117	Identifying genetic determinants of inflammatory pain in mice using a large-scale gene-targeted screen <i>Pain</i> , 2022 , 163, 1139-1157	8	0
116	Modeling Secondary Phenotypes Conditional on Genotypes in Casellontrol Studies. <i>Stats</i> , 2022 , 5, 203	-21549	
115	Acute inflammatory response via neutrophil activation protects against the development of chronic pain <i>Science Translational Medicine</i> , 2022 , 14, eabj9954	17.5	1
114	Sex- and age-specific genetic analysis of chronic back pain. <i>Pain</i> , 2021 , 162, 1176-1187	8	6
113	Phenotypic profile clustering pragmatically identifies diagnostically and mechanistically informative subgroups of chronic pain patients. <i>Pain</i> , 2021 , 162, 1528-1538	8	2
112	Single cell transcriptomics of primate sensory neurons identifies cell types associated with chronic pain. <i>Nature Communications</i> , 2021 , 12, 1510	17.4	35
111	Identification and characterization of novel candidate compounds targeting 6- and 7-transmembrane Eppioid receptor isoforms. <i>British Journal of Pharmacology</i> , 2021 , 178, 2709-2726	8.6	2
110	Mast cell stabilizer ketotifen fumarate reverses inflammatory but not neuropathic-induced mechanical pain in mice. <i>Pain Reports</i> , 2021 , 6, e902	3.5	O
109	Alternative Splicing of Opioid Receptor Genes Shows a Conserved Pattern for 6TM Receptor Variants. <i>Cellular and Molecular Neurobiology</i> , 2021 , 41, 1039-1055	4.6	1
108	Reversion mutation of cDNA CA8-204 minigene construct produces a truncated functional peptide that regulates calcium release in vitro and produces profound analgesia in vivo. <i>Mammalian Genome</i> , 2020 , 31, 287-294	3.2	1
107	The dichotomous role of epiregulin in pain. <i>Pain</i> , 2020 , 161, 1052-1064	8	7
106	The geriatric pain experience in mice: intact cutaneous thresholds but altered responses to tonic and chronic pain. <i>Neurobiology of Aging</i> , 2020 , 89, 1-11	5.6	5
105	NK cell recruitment limits tissue damage during an enteric helminth infection. <i>Mucosal Immunology</i> , 2020 , 13, 357-370	9.2	11
104	Detangling red hair from pain: phenotype-specific contributions from different genetic variants in melanocortin-1 receptor. <i>Pain</i> , 2020 , 161, 938-948	8	1
103	A functional polymorphism in the ATP-Binding Cassette B1 transporter predicts pharmacologic response to combination of nortriptyline and morphine in neuropathic pain patients. <i>Pain</i> , 2020 , 161, 619-629	8	7
102	A genetic polymorphism that is associated with mitochondrial energy metabolism increases risk of fibromyalgia. <i>Pain</i> , 2020 , 161, 2860-2871	8	4

(2017-2020)

101	Premorbid and concurrent predictors of TMD onset and persistence. <i>European Journal of Pain</i> , 2020 , 24, 145-158	3.7	12
100	A functional substitution in the L-aromatic amino acid decarboxylase enzyme worsens somatic symptoms via a serotonergic pathway. <i>Annals of Neurology</i> , 2019 , 86, 168-180	9.4	5
99	Stabilization of Eppioid receptor facilitates its cellular translocation and signaling. <i>Proteins: Structure, Function and Bioinformatics</i> , 2019 , 87, 878-884	4.2	4
98	Cartilage-binding antibodies induce pain through immune complex-mediated activation of neurons. Journal of Experimental Medicine, 2019 , 216, 1904-1924	16.6	34
97	Profound analgesia is associated with a truncated peptide resulting from tissue specific alternative splicing of DRG CA8-204 regulated by an exon-level cis-eQTL. <i>PLoS Genetics</i> , 2019 , 15, e1008226	6	3
96	Alternative Splicing of the Delta-Opioid Receptor Gene Suggests Existence of New Functional Isoforms. <i>Molecular Neurobiology</i> , 2019 , 56, 2855-2869	6.2	13
95	Disentangling the genetics of lean mass. American Journal of Clinical Nutrition, 2019, 109, 276-287	7	24
94	CACNG2 polymorphisms associate with chronic pain after mastectomy. <i>Pain</i> , 2019 , 160, 561-568	8	11
93	Genome-wide association reveals contribution of MRAS to painful temporomandibular disorder in males. <i>Pain</i> , 2019 , 160, 579-591	8	26
92	Genetic pathway analysis reveals a major role for extracellular matrix organization in inflammatory and neuropathic pain. <i>Pain</i> , 2019 , 160, 932-944	8	31
91	A study in scarlet: MC1R as the main predictor of red hair and exemplar of the flip-flop effect. <i>Human Molecular Genetics</i> , 2019 , 28, 2093-2106	5.6	4
90	Human pain genetics database: a resource dedicated to human pain genetics research. <i>Pain</i> , 2018 , 159, 749-763	8	37
89	The human pain genetics database: an interview with Luda Diatchenko. <i>Pain Management</i> , 2018 , 8, 259	-263	
88	Genetic studies of human neuropathic pain conditions: a review. <i>Pain</i> , 2018 , 159, 583-594	8	38
87	Low back pain. Nature Reviews Disease Primers, 2018, 4, 52	51.1	118
86	Human carbonic anhydrase-8 AAV8 gene therapy inhibits nerve growth factor signaling producing prolonged analgesia and anti-hyperalgesia in mice. <i>Gene Therapy</i> , 2018 , 25, 297-311	4	4
85	N6-methyladenosine alters RNA structure to regulate binding of a low-complexity protein. <i>Nucleic Acids Research</i> , 2017 , 45, 6051-6063	20.1	339
84	Genetic variant rs3750625 in the 3SUTR of ADRA2A affects stress-dependent acute pain severity after trauma and alters a microRNA-34a regulatory site. <i>Pain</i> , 2017 , 158, 230-239	8	10

83	Car8 dorsal root ganglion expression and genetic regulation of analgesic responses are associated with a cis-eQTL in mice. <i>Mammalian Genome</i> , 2017 , 28, 407-415	3.2	5
82	miR-183 cluster scales mechanical pain sensitivity by regulating basal and neuropathic pain genes. <i>Science</i> , 2017 , 356, 1168-1171	33.3	80
81	The more you test, the more you find: The smallest P-values become increasingly enriched with real findings as more tests are conducted. <i>Genetic Epidemiology</i> , 2017 , 41, 726-743	2.6	О
80	T-Cell Mediation of Pregnancy Analgesia Affecting Chronic Pain in Mice. <i>Journal of Neuroscience</i> , 2017 , 37, 9819-9827	6.6	33
79	Post-concussion symptoms and chronic pain after mild traumatic brain injury are modulated by multiple locus effect in the gene through the expression of antisense: A pilot prospective control study <i>Canadian Journal of Pain</i> , 2017 , 1, 112-126	1.5	2
78	Epiregulin and EGFR interactions are involved in pain processing. <i>Journal of Clinical Investigation</i> , 2017 , 127, 3353-3366	15.9	54
77	Effect of Human Genetic Variability on Gene Expression in Dorsal Root Ganglia and Association with Pain Phenotypes. <i>Cell Reports</i> , 2017 , 19, 1940-1952	10.6	34
76	N(6)-Methyladenosine Modification in a Long Noncoding RNA Hairpin Predisposes Its Conformation to Protein Binding. <i>Journal of Molecular Biology</i> , 2016 , 428, 822-833	6.5	122
75	Identification of clusters of individuals relevant to temporomandibular disorders and other chronic pain conditions: the OPPERA study. <i>Pain</i> , 2016 , 157, 1266-1278	8	62
74	Human Genetic Variability Contributes to Postoperative Morphine Consumption. <i>Journal of Pain</i> , 2016 , 17, 628-36	5.2	44
73	Agonist-dependence of functional properties for common nonsynonymous variants of human transient receptor potential vanilloid 1. <i>Pain</i> , 2016 , 157, 1515-1524	8	14
7 2	Modification of COMT-dependent pain sensitivity by psychological stress and sex. <i>Pain</i> , 2016 , 157, 858-	8 6 7	29
71	The nicotinic 8 subunit gene determines variability in chronic pain sensitivity via cross-inhibition of P2X2/3 receptors. <i>Science Translational Medicine</i> , 2015 , 7, 287ra72	17.5	44
7º	Subgrouping of low back pain patients for targeting treatments: evidence from genetic, psychological, and activity-related behavioral approaches. <i>Clinical Journal of Pain</i> , 2015 , 31, 123-32	3.5	31
69	Dual allosteric modulation of opioid antinociceptive potency by 🛮 A-adrenoceptors. <i>Neuropharmacology</i> , 2015 , 99, 285-300	5.5	12
68	Expioid receptor 6-transmembrane isoform: A potential therapeutic target for new effective opioids. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015 , 62, 61-7	5.5	23
67	Molecular genetic mechanisms of allelic specific regulation of murine Comt expression. <i>Pain</i> , 2015 , 156, 1965-1977	8	5
66	Neuropathic pain phenotyping by international consensus (NeuroPPIC) for genetic studies: a NeuPSIG systematic review, Delphi survey, and expert panel recommendations. <i>Pain</i> , 2015 , 156, 2337-2	2353	54

(2013-2015)

65	COMT gene locus: new functional variants. <i>Pain</i> , 2015 , 156, 2072-2083	8	20
64	Carbonic anhydrase-8 regulates inflammatory pain by inhibiting the ITPR1-cytosolic free calcium pathway. <i>PLoS ONE</i> , 2015 , 10, e0118273	3.7	24
63	Differential Regulation of 6- and 7-Transmembrane Helix Variants of Expioid Receptor in Response to Morphine Stimulation. <i>PLoS ONE</i> , 2015 , 10, e0142826	3.7	13
62	Genome-wide association meta-analyses to identify common genetic variants associated with hallux valgus in Caucasian and African Americans. <i>Journal of Medical Genetics</i> , 2015 , 52, 762-9	5.8	13
61	Differences in the Antinociceptive Effects and Binding Properties of Propranolol and Bupranolol Enantiomers. <i>Journal of Pain</i> , 2015 , 16, 1321-1333	5.2	21
60	Quantitative H2S-mediated protein sulfhydration reveals metabolic reprogramming during the integrated stress response. <i>ELife</i> , 2015 , 4, e10067	8.9	113
59	Structural and functional interactions between six-transmembrane Expioid receptors and 2 -adrenoreceptors modulate opioid signaling. <i>Scientific Reports</i> , 2015 , 5, 18198	4.9	25
58	Epistasis between polymorphisms in COMT, ESR1, and GCH1 influences COMT enzyme activity and pain. <i>Pain</i> , 2014 , 155, 2390-9	8	45
57	Complex multilocus effects of catechol-O-methyltransferase haplotypes predict pain and pain interference 6 weeks after motor vehicle collision. <i>NeuroMolecular Medicine</i> , 2014 , 16, 83-93	4.6	26
56	Facial pain with localized and widespread manifestations: separate pathways of vulnerability. <i>Pain</i> , 2013 , 154, 2335-2343	8	27
55	Signs and symptoms of first-onset TMD and sociodemographic predictors of its development: the OPPERA prospective cohort study. <i>Journal of Pain</i> , 2013 , 14, T20-32.e1-3	5.2	114
54	Pain modality- and sex-specific effects of COMT genetic functional variants. <i>Pain</i> , 2013 , 154, 1368-76	8	69
53	Multisystem dysregulation in painful temporomandibular disorders. Journal of Pain, 2013, 14, 983-96	5.2	43
52	Multivariable modeling of phenotypic risk factors for first-onset TMD: the OPPERA prospective cohort study. <i>Journal of Pain</i> , 2013 , 14, T102-15	5.2	58
51	Clinical orofacial characteristics associated with risk of first-onset TMD: the OPPERA prospective cohort study. <i>Journal of Pain</i> , 2013 , 14, T33-50	5.2	102
50	Psychological factors associated with development of TMD: the OPPERA prospective cohort study. <i>Journal of Pain</i> , 2013 , 14, T75-90	5.2	212
49	Study protocol, sample characteristics, and loss to follow-up: the OPPERA prospective cohort study. <i>Journal of Pain</i> , 2013 , 14, T2-19	5.2	47
48	Summary of findings from the OPPERA prospective cohort study of incidence of first-onset temporomandibular disorder: implications and future directions. <i>Journal of Pain</i> , 2013 , 14, T116-24	5.2	128

47	The phenotypic and genetic signatures of common musculoskeletal pain conditions. <i>Nature Reviews Rheumatology</i> , 2013 , 9, 340-50	8.1	152
46	Large candidate gene association study reveals genetic risk factors and therapeutic targets for fibromyalgia. <i>Arthritis and Rheumatism</i> , 2012 , 64, 584-93		67
45	Genetically determined P2X7 receptor pore formation regulates variability in chronic pain sensitivity. <i>Nature Medicine</i> , 2012 , 18, 595-9	50.5	265
44	Excess risk of temporomandibular disorder associated with cigarette smoking in young adults. Journal of Pain, 2012 , 13, 21-31	5.2	28
43	Relationship between temporomandibular disorders, widespread palpation tenderness, and multiple pain conditions: a case-control study. <i>Journal of Pain</i> , 2012 , 13, 1016-27	5.2	47
42	Construction of a global pain systems network highlights phospholipid signaling as a regulator of heat nociception. <i>PLoS Genetics</i> , 2012 , 8, e1003071	6	15
41	Elucidation of mu-Opioid Gene Structure: How Genetics Can Help Predict Responses to Opioids. <i>European Journal of Pain Supplements</i> , 2011 , 5, 433-438		13
40	Catechol O-methyltransferase haplotype predicts immediate musculoskeletal neck pain and psychological symptoms after motor vehicle collision. <i>Journal of Pain</i> , 2011 , 12, 101-7	5.2	72
39	Study methods, recruitment, sociodemographic findings, and demographic representativeness in the OPPERA study. <i>Journal of Pain</i> , 2011 , 12, T12-26	5.2	100
38	Orofacial pain prospective evaluation and risk assessment studythe OPPERA study. <i>Journal of Pain</i> , 2011 , 12, T4-11.e1-2	5.2	190
37	Potential genetic risk factors for chronic TMD: genetic associations from the OPPERA case control study. <i>Journal of Pain</i> , 2011 , 12, T92-101	5.2	135
36	Summary of findings from the OPPERA baseline case-control study: implications and future directions. <i>Journal of Pain</i> , 2011 , 12, T102-7	5.2	51
35	Potential autonomic risk factors for chronic TMD: descriptive data and empirically identified domains from the OPPERA case-control study. <i>Journal of Pain</i> , 2011 , 12, T75-91	5.2	84
34	Structural basis for Eppioid receptor binding and activation. <i>Structure</i> , 2011 , 19, 1683-90	5.2	23
33	Cytokine biomarkers and chronic pain: association of genes, transcription, and circulating proteins with temporomandibular disorders and widespread palpation tenderness. <i>Pain</i> , 2011 , 152, 2802-2812	8	86
32	Disruptive mRNA folding increases translational efficiency of catechol-O-methyltransferase variant. <i>Nucleic Acids Research</i> , 2011 , 39, 6201-12	20.1	42
31	Structural mechanism of S-adenosyl methionine binding to catechol O-methyltransferase. <i>PLoS ONE</i> , 2011 , 6, e24287	3.7	23
30	Pain perception is altered by a nucleotide polymorphism in SCN9A. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 5148-53	11.5	230

(2003-2010)

29	Multiple chronic pain states are associated with a common amino acid-changing allele in KCNS1. <i>Brain</i> , 2010 , 133, 2519-27	11.2	187
28	A genome-wide Drosophila screen for heat nociception identifies 28 as an evolutionarily conserved pain gene. <i>Cell</i> , 2010 , 143, 628-38	56.2	217
27	A novel alternatively spliced isoform of the mu-opioid receptor: functional antagonism. <i>Molecular Pain</i> , 2010 , 6, 33	3.4	51
26	Low enzymatic activity haplotypes of the human catechol-O-methyltransferase gene: enrichment for marker SNPs. <i>PLoS ONE</i> , 2009 , 4, e5237	3.7	38
25	Expansion of the human mu-opioid receptor gene architecture: novel functional variants. <i>Human Molecular Genetics</i> , 2009 , 18, 1037-51	5.6	137
24	Signaling pathways mediating beta3-adrenergic receptor-induced production of interleukin-6 in adipocytes. <i>Molecular Immunology</i> , 2009 , 46, 2256-66	4.3	27
23	Homogeneous reporter system enables quantitative functional assessment of multiple transcription factors. <i>Nature Methods</i> , 2008 , 5, 253-60	21.6	66
22	Orthodontic Treatment, Genetic Factors and Risk of Temporomandibular Disorder. <i>Seminars in Orthodontics</i> , 2008 , 14, 146-156	1.2	51
21	Beta2 adrenergic receptor activation stimulates pro-inflammatory cytokine production in macrophages via PKA- and NF-kappaB-independent mechanisms. <i>Cellular Signalling</i> , 2007 , 19, 251-60	4.9	152
20	Genetic architecture of human pain perception. <i>Trends in Genetics</i> , 2007 , 23, 605-13	8.5	161
19	Catechol-O-methyltransferase inhibition increases pain sensitivity through activation of both beta2- and beta3-adrenergic receptors. <i>Pain</i> , 2007 , 128, 199-208	8	207
18	Responses to Drs. Kim and Dionne regarding comments on Diatchenko, et al. Catechol-O-methyltransferase gene polymorphisms are associated with multiple pain-evoking stimuli. Pain 2006; 125: 216-24. <i>Pain</i> , 2007 , 129, 366-370	8	17
17	Idiopathic pain disorderspathways of vulnerability. <i>Pain</i> , 2006 , 123, 226-230	8	283
16	Catechol-O-methyltransferase gene polymorphisms are associated with multiple pain-evoking stimuli. <i>Pain</i> , 2006 , 125, 216-224	8	275
15	GTP cyclohydrolase and tetrahydrobiopterin regulate pain sensitivity and persistence. <i>Nature Medicine</i> , 2006 , 12, 1269-77	50.5	435
14	Genetic basis for individual variations in pain perception and the development of a chronic pain condition. <i>Human Molecular Genetics</i> , 2005 , 14, 135-43	5.6	980
13	Identification of novel mediators of NF-kappaB through genome-wide survey of monocyte adherence-induced genes. <i>Journal of Leukocyte Biology</i> , 2005 , 78, 1366-77	6.5	37
12	Gene expression analysis of purified hematopoietic stem cells and committed progenitors. <i>Blood</i> , 2003 , 102, 94-101	2.2	177

11	Gene expression profiling in RAS oncogene-transformed cell lines and in solid tumors using subtractive suppression hybridization and cDNA arrays. <i>Advances in Enzyme Regulation</i> , 2002 , 42, 63-82		17
10	Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 16899-903	11.5	1457
9	Caveolin-1 is down-regulated in human ovarian carcinoma and acts as a candidate tumor suppressor gene. <i>American Journal of Pathology</i> , 2001 , 159, 1635-43	5.8	234
8	Use of SMART-generated cDNA for gene expression studies in multiple human tumors. <i>BioTechniques</i> , 2001 , 30, 158-63	2.5	52
7	Structure and regulation of the mouse ing1 gene. Three alternative transcripts encode two phd finger proteins that have opposite effects on p53 function. <i>Journal of Biological Chemistry</i> , 1999 , 274, 32172-81	5.4	53
6	Suppression subtractive hybridization: a versatile method for identifying differentially expressed genes. <i>Methods in Enzymology</i> , 1999 , 303, 349-80	1.7	300
5	Stress-induced secretion of growth inhibitors: a novel tumor suppressor function of p53. <i>Oncogene</i> , 1998 , 17, 1089-96	9.2	130
4	Construction of cDNA libraries from small amounts of total RNA using the suppression PCR effect. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 230, 285-8	3.4	37
3	Equalizing cDNA subtraction based on selective suppression of polymerase chain reaction: cloning of Jurkat cell transcripts induced by phytohemaglutinin and phorbol 12-myristate 13-acetate. <i>Analytical Biochemistry</i> , 1996 , 240, 90-7	3.1	223
2	Genome-wide analysis identifies significant contribution of brain-expressed genes in chronic, but not acute, back pain		2
1	Single cell transcriptomics of primate sensory neurons identifies cell types associated with human chronic pain		3